

[Available Issues](#) | [Japanese](#)>> [Publisher Site](#)

Author:  [ADVANCED](#) | Volume  Page   
 Keyword:



[TOP](#) > [Available Issues](#) > [Table of Contents](#) > [Abstract](#)

PRINT ISSN : 0040-8891

## The Bulletin of Tokyo Dental College

Vol. 47 (2006) , No. 1 :19-23

[\[PDF \(801K\)\]](#) [\[References\]](#)

### Two Cases with Supernumerary Teeth in Lower Incisor Region

[Taishi Yokose](#)<sup>1)</sup>, [Teruo Sakamoto](#)<sup>1)</sup>, [Kenji Sueishi](#)<sup>2)</sup>, [Kenichi Yatabe](#)<sup>2)</sup>, [Keiichirou Tsujino](#)<sup>3)</sup>, [Shuhei Kubo](#)<sup>4)</sup>, [Masashi Yakushiji](#)<sup>4)</sup> and [Hideharu Yamaguchi](#)<sup>1)</sup>

1) Department of Orthodontics, Tokyo Dental College

2) Division of Orthodontics, Department of Clinical Oral Health Science, Tokyo Dental College

3) Division of Pediatric Dentistry, Department of Clinical Oral Health Science, Tokyo Dental College

4) Department of Pediatric Dentistry, Tokyo Dental College

(Received February 6, 2006)

(Accepted April 19, 2006)

**Abstract:** Abnormalities in number of teeth are occasionally noted in clinical cases. Many theories have been proposed as regards the causes of the occurrence of supernumerary teeth, including atavism theory, mechanical tooth germ separation theory, tissue induction theory, and dental lamina morphological disturbance theory. However, none of these theories alone offers a sufficient explanation for this phenomenon. The incidence of supernumerary permanent teeth is approximately 1-3%. These are the maxillary anterior teeth, the maxillary molars, and the maxillo-mandibular premolars in terms of descending order of site of occurrence. On the other hand, incidence in the mandibular anterior tooth area, of which there have been few detailed reports, is about 0.01%, a markedly low value. In this paper, we report two rare cases of supernumerary teeth in the mandibular incisor area. We discuss their etiology and orthodontic treatment, and detail a differential diagnosis between the normal and supernumerary teeth. We found that it was difficult to establish a clear etiology and differentiation between the normal and supernumerary teeth.

**Key words:** [Supernumerary teeth](#), [Lower incisor](#), [Etiology](#), [Fused tooth](#), [Orthodontic treatment with supernumerary teeth](#)

[\[PDF \(801K\)\]](#) [\[References\]](#)

To cite this article:

Taishi Yokose, Teruo Sakamoto, Kenji Sueishi, Kenichi Yatabe, Keiichirou Tsujino, Shuhei Kubo, Masashi Yakushiji and Hideharu Yamaguchi: "Two Cases with Supernumerary Teeth in Lower Incisor Region". The Bulletin of Tokyo Dental College, Vol. **47**: 19-23 (2006) .

---

doi:10.2209/tdcpublication.47.19

JOI JST.JSTAGE/tdcpublication/47.19

Copyright (c) 2006 by Tokyo Dental College, Japan

---



---

[Japan Science and Technology Information Aggregator, Electronic](#)

