

[\[Print Version\]](#)
[\[PubMed Citation\]](#) [\[Related Articles in PubMed\]](#)

TABLE OF CONTENTS

[\[INTRODUCTION\]](#) [\[METHOD AND...\]](#) [\[RESULTS\]](#) [\[DISCUSSION\]](#) [\[CONCLUSION\]](#) [\[REFERENCES\]](#) [\[TABLES\]](#) [\[FIGURES\]](#)

The Angle Orthodontist: Vol. 70, No. 4, pp. 297-302.

Maxillary Arch Size and Shape in American Blacks and Whites

Benjamin G. Burris, BA,^a Edward F. Harris, PhD^b

ABSTRACT

American blacks have larger teeth than whites, but they less frequently exhibit crowding—apparently because of larger arch dimensions. This study quantified differences in arch size and shape in these 2 constituents of the US population. Eighteen dental and bony landmarks were digitized from the maxilla of each of 332 subjects with permanent, intact dentitions, proportionately divided between blacks and whites, men and women. Linear, angular, and area measurements were computer-generated. Arch widths averaged 10% greater in blacks than whites, and mesiodistal arch depths had a greater difference, at 12%. Blacks, with a more square palate and significantly larger palatal index, were distinguished from whites primarily by greater intercanine and interpremolar widths. Arch perimeter was greater in blacks by 8%, and cross-sectional area of the arch was 19% greater in blacks than whites, so blacks and whites differ substantially for these parameters not only in size, but in shape as well. These differences are relevant in prosthodontics and orthodontics since individualization of treatment leads to more effective treatment by working within the patient's natural arch form instead of making patients fit a single standard.

KEY WORDS: Arch size, Arch form, Ethnic differences, Sex differences.

Accepted: March 2000. Submitted: December 1999.



INTRODUCTION [Return to TOC](#)

American blacks have significantly larger tooth crown dimensions than American whites,^{1,2} but, ironically, blacks are less likely to exhibit dental crowding and, actually, are more likely to have spacing between anterior teeth.³ Blacks have larger arch dimensions that provide "extra" space in which teeth can erupt. Not only are arch dimensions larger on average in American blacks, but it also seems that arch form differs, being less parabolic and more square than in whites.⁴ These differences can affect clinical treatment. While it is obvious that the dentist treats the individual and not some abstraction of the population, it also holds that people from different ethnic groups present with different modal conditions, and the clinician should anticipate the differences in size and form rather than treating all cases to a single ideal.


The purpose of this study was to quantify the nature of the arch form in a sample of American blacks and whites. The intent was to characterize the difference between ethnic groups (and sexes) in order to identify ways of customizing treatment (eg, prosthodontics, orthodontics) appropriate to the patient's ethnic background.

METHOD AND MATERIALS [Return to TOC](#)

Full-mouth dental study casts were obtained from 330 routine dental patients. The sample was composed proportionately of American blacks (n = 171) and whites (n = 159) and of men (n = 164) and women (n = 166). Ethnic affiliation was self-assessed by the patient. Admixture estimates of whites with American blacks are on the order of 20% to 30% for Northern and Western samples⁵⁻⁸ but are generally below 10% for Southeastern samples.⁹⁻¹³ All patients had a full complement of natural teeth (excluding third molars), although they were not selected on the basis of occlusion. No patients had been treated orthodontically, although some were treated subsequently. The samples represent a cross-section of routine adolescent and young adult dental patients.


Eighteen buccal cusp tips and incisor line angles were marked on each maxillary dental cast ([Figure 1A](#) ) and a 1-to-1 ratio print was made, oriented perpendicular to the occlusal surface.¹⁴ Millimetric scales were included in the print to guard against magnification and parallax. The dental landmarks then were digitized as Cartesian coordinates, and mesiodistal arch depths and buccolingual arch widths were computer-generated as straight-line distances between the dental landmarks ([Figure 1B,C](#) ). Arch depths were measured parallel to the midpalatal raphe, which was defined during data entry.




The set of dental landmarks also was used to fit a curve to each dental arch. Many suggestions of a relevant curve have been put forth in the literature, such as an ellipse, a parabola, and a catenary curve. Lu¹⁵ showed that a close fit with the human dental arch could be obtained with a fourth-order polynomial. Of note, each of the 4 weighting coefficients in the equation is interpretable in terms of an arch's form, namely its left-right asymmetry (terms b_1 and b_3) and its taperedness (b_2) and squareness (b_4). The least-squares best fit was calculated for each arch form, and the 4 regression coefficients were evaluated between races and sexes.

Two other variables, arch perimeter and arch area, were also measured. Perimeter was obtained by plotting the dental coordinates for a case and fitting them with a spline function. A spline is a smooth curve that intersects all of the points. This was accomplished using commercial software (DeltaGraph, DeltaPoint, Monterey, Cal). With the spline function plotted, the base of the dental arch was defined by a line passing through the mesiobuccal cusp tips of the second molars ([Figure 1D](#) ). This outline (with appropriate scales) was transferred to an image-processing program (NIH Image, Bethesda, Md), where the cross-sectional area was calculated. The area of the arch thus encompasses the dental elements as well as the bony palate, though it does not include the distal portion adjacent to and behind the second molars.

Two-way factorial analysis of variance was used to test for black-white and sex differences in arch size.¹⁶ The distribution of differences was assessed with nonparametric tests as


RESULTS [Return to TOC](#)


Blacks had highly significant larger mean arch widths and depths than whites ([Table 1](#) ). Arch widths averaged 9% larger in black men than white men, and 11% larger in black women than white women. Indeed, by a signed-ranks test, black and white women were significantly more different than men in the 2 groups ($P = .01$). Additionally, arch depths differed relatively more between races ($\bar{X} = 12\%$) than the transverse widths ($\bar{X} = 9\%$); this length-width difference also achieved significance ($P < .01$). The greatest percentage difference, both for length and breadth, was in the canine and first premolar region, which coincides with blacks having a less-tapered arch form.


These metric differences are apparent when graphed ([Figure 2](#) ), and it is evident that the black-white difference is much greater than the level of sexual dimorphism within either ethnic group ([Figure 3](#) ). Percentage-wise, the greatest black-white difference in arch widths is at the first premolars. In blacks, the buccal cusp of the first premolar is lateral of a line defined by the canine and second premolar ([Figure 2](#) ). In whites, the buccal cusps of the canine and 2 premolars are in approximately a straight line that is more convergent mesially than in blacks. Howes¹⁹ showed that arch width at the first premolar is a practical clinical tool for assessing the taperedness of a dental arch (when the 4–4 width is narrow) vs the broad or square arch (where 4–4 width is broader). It is precisely this feature—relative width at the first premolars—that distinguishes the shape of these black and white dental arches.


Lu¹⁵ showed that a fourth-order orthogonal polynomial produces a close fit with the shape of the human dental arch. The formula is


$$Y = b_1X^1 + b_2X^2 + b_3X^3 + b_4X^4$$

with the odd-numbered terms (b_1 and b_3) defining left-right asymmetry of an arch and the even-numbered terms measuring taperedness (b_2) and squareness (b_4). Asymmetry did not differ by race or sex ([Table 1](#) ), and 1-sample t -tests showed that there was no directional asymmetry in arch form (ie, means of b_1 and b_3 did not differ significantly from zero). This indicates that neither side of the palate is systematically longer than the other. In contrast, Woo,²⁰ Livshits and Koblyansky,²¹ and others have shown that various craniofacial complexes are characteristically larger on one side of the skull than the other.

Blacks had smaller values for b_2 , reflecting their less convergent (more nearly parallel) middle and posterior arch segments. Whites, on the other hand, with a larger value for b_2 , had more rounded arch forms. This is discernible in [Figure 3](#)  where all arch widths and depths are smaller in whites, and the black-white difference in widths is greatest in the midarch. The average arch form in whites possessed greater convergence toward the front of the arch than in blacks. The shape of the arch is, however, similar in the pair of sexes within each ethnic group. An ethnic difference in arch form also is suggested by b_4 , with a tendency towards more square arches in blacks ($P = .02$).



Since the fitted polynomial indicated that arch form is more tapered in whites than blacks, the palatal index,²² which is the ratio of arch length to width, should confirm a difference. The ratio of I1-to-M2 length divided by arch width at M2 was expressed as an index by postmultiplying by 100 ([Table 2](#) ). The black-white difference was highly significant ($P = .0002$). Arch length was a smaller percentage of M2 width in whites, confirming a rounder, less-square arch form. In contrast, there was no suggestion of a sex difference in palatal index within either ethnic group.


This shape difference also can be seen in the ratio of the front of the arch (C to C) expressed as a ratio of the back of the arch (M2 to M2). This width ratio²³ was significantly greater in blacks ([Table 2](#) ). Blacks have a greater width ratio because, on the average, their arches do not taper as much as in whites. Hence, even using different criteria it is evident that, in addition to greater size, blacks characteristically possess less-tapered arch forms.

As a statistical average, American blacks have appreciably larger tooth crown diameters than American whites,²⁴ so it is not surprising to find that perimeter of the arch, defined by the buccal cusps and incisor line angles, also is larger in blacks ([Table 2](#) ). Perimeter was 8% greater in blacks and there was a significant sex difference within each ethnic group (about 3%).


Cross-sectional area of the arch showed the greatest relative black-white difference, at 19%, because it is the cumulative product of the greater arch widths and depths of blacks. Areas exceeded 2000 mm² in black men and women, but averaged 1828 mm² in white men and 1673 mm² in white women.

DISCUSSION [Return to TOC](#)

African Americans constitute 12% of the US population, but comparatively little is known about their dental characteristics. On average, blacks have significantly larger arch lengths and widths than American whites (about 10% larger; [Figure 2](#) ), but the difference is not simply one of larger isometric dimensions. Instead, there also are differences in shape—of proportionality—and the most obvious differences are in the canine-first premolar region. The buccal cusps from the canine through the second premolar define a convex curve in blacks, while the canine and premolars are in an approximately straight line in whites. [Figure 4](#)  shows the average black and white arch forms registered at the level of the canines. This highlights how the arch in whites is disproportionately narrow in the canine-first premolar area, and it also shows how these whites define a more rounded arch form since the decrease in radius from the second molar forward is steeper. In contrast, the straighter and less convergent buccal tooth rows in blacks define a more square arch form.

These findings are essentially independent of the patient's age. Teeth remain very close to their erupted positions in the absence of outside forces.^{25,26} And, while there is some growth of the intermaxillary and interpalatine sutures,^{27,28} the amount of growth is too subtle to detect without long-term longitudinal analysis. That is, in cross-sectional data as analyzed here and in the clinical setting, interindividual variation is considerable while the arch changes within an individual are very small.²⁹ Therefore, there is no detectable systematic change in arch size or form with age. Indeed, each variable in the present study was regressed on the person's age, but none exhibited anywhere near a statistically significant age effect. The clinical consequence is that the same arch form in the permanent dentition (eg, [Figure 2](#) ) is equally appropriate irrespective of the patient's age.

Transseptal fibers³⁰ and the anterior component of force^{31,32} maintain tight occlusal contacts and, over the long term, cause the dental arch to become slightly shorter and broader.^{33,34} These effects can be detected if the same individuals are followed for a long span of adulthood,²⁹ but they are obscured in cross-sectional studies where different ages are represented by different individuals.

Various classifications have been developed to characterize arch forms, such as elliptical, parabolic, square, and U-shaped.^{35,36} As Jacobson⁴ and others note, such schemes are poorly defined and cause confusion within and among studies. The palatal index provides a quantitative sense of arch form. American blacks have a significantly higher palatal index; arch length (I1 to M2) is a greater proportion of molar width (M2–M2) because the buccal tooth rows are more nearly parallel rather than converging as sharply as in whites. This difference also can be appreciated from the width ratio between the canines and second molars, which is a measure of the size gradient between widths at the front and back of the arch ([Table 2](#) ). The ratio is significantly higher in blacks, again confirming the broader, less-tapered arch form in blacks compared to whites.

CONCLUSION [Return to TOC](#)

Arch size is notably larger in American blacks than whites and arch form is squarer and less tapered in blacks in the canine-premolar region. These ethnic differences should be considered during treatment, especially in disciplines such as prosthodontics and orthodontics where arch shape can be modified appreciably.

1. Richardson ER, Malhotra SK. Mesiodistal crown dimension of the permanent dentition of American Negroes. *Am J Orthod.* 1975; 68:157–164. [[PubMed Citation](#)]
2. Macko DJ, Ferguson FS, Sonnenberg EM. Mesiodistal crown dimensions of permanent teeth of Black Americans. *J Dent Child.* 1979; 46:42–46.
3. Kelly JE, Harvey CR. An assessment of the occlusion of youths 12–17 years. *United States Public Health Service, Vital and Health Statistics. No. 162, Series 11.* Washington, DC. 1977;
4. Jacobson A. *The Dentition of the South African Negro.* Anniston, Al: Higginbotham Inc. 1982;
5. Glass HB, Li CC. The dynamics of racial intermixture—an analysis based on the American Negro. *Am J Hum Genet.* 1953; 5:1–20.
6. Glass HB. On the unlikelihood of significant admixture of genes from North American Indians in the present composition of the Negroes of the United States. *Am J Hum Genet.* 1955; 7:368–385.
7. Roberts DF. The dynamics of racial intermixture in the American Negro—some anthropological considerations. *Am J Hum Genet.* 1955; 7:361–367.
8. Roberts DF, Hiorns RW. The dynamics of racial intermixture. *Am J Hum Genet.* 1962; 14:261–277.
9. Pollitzer WS. The Negroes of Charleston (SC): a study of hemoglobin types, serology, and morphology. *Am J Phys Anthropol.* 1958; 16:241–263.
10. Workman P, Blumberg B, Cooper AJ. Selection, gene migration, and polymorphic stability in a U.S. white and Negro population. *Am J Hum Genet.* 1963; 15:429–437.
11. Pollitzer WS, Menegaz-Bock RM, Ceppellini R, Dunn LC. Blood factors and morphology of the Negroes of James Island, Charleston, SC. *Am J Phys Anthropol.* 1964; 22:393–398.
12. Pollitzer WS, Boyle E, Cornoni J, Namboodiri KK. Physical anthropology of the Negroes of Charleston, SC. *Hum Biol.* 1970; 42:265–279. [[PubMed Citation](#)]
13. Blumberg B, Hesser JE. Loci differentially affected by selection in 2 American black populations. *Proc Natl Acad Sci.* 1971; 68:2554–2558. [[PubMed Citation](#)]
14. Brown T, Abbott AH, Burgess VB. Age changes in dental arch dimensions of Australian Aborigines. *Am J Phys Anthropol.* 1983; 62:291–303. [[PubMed Citation](#)]
15. Lu KH. An orthogonal analysis of the form, symmetry and asymmetry of the dental arch. *Arch Oral Biol.* 1966; 11:1057–1069. [[PubMed Citation](#)]
16. Winer BJ. *Statistical Principles in Experimental Design. 2nd ed.* New York, NY: McGraw-Hill Inc. 1971;
17. Siegel S, Castellan NJ Jr. *Nonparametric statistics for the behavioral sciences.* 2nd ed. New York, NY: McGraw-Hill Inc. 1988;
18. Holm S. A simple sequentially rejective multiple test procedure. *Scand J Statist.* 1979; 6:65–70.
19. Howes JH. A polygon portrayal of coronal and basal arch dimensions in the horizontal plane. *Am J Orthod.* 1954; 40:811–831.
20. Woo TL. On the asymmetry of the human skull. *Biometrika.* 1931; 22:324–352.
21. Livshits G, Koblyansky E. Fluctuating asymmetry as a possible measure of developmental homeostasis in humans, a review. *Hum Biol.* 1991; 63:441–466. [[PubMed Citation](#)]
22. Martin R. *Lehrbuch der Anthropologie in Systematischer Darstellung.* 2nd ed. Jena: Gustav Fischer. 1928;
23. Smith RJ, Bailit HL. Variation in dental occlusion and arches among Melanesians of Bougainville Island, Papua New Guinea. I. Methods, age changes, sex differences and population comparisons. *Am J Phys Anthropol.* 1977; 47:195–208. [[PubMed Citation](#)]
24. Harris EF, Rathbun TA. Small tooth sizes in a 19th century South Carolina plantation slave series. *Am J Phys Anthropol.* 1989; 78:411–420. [[PubMed Citation](#)]
25. Moorrees CFA. *The dentition of the growing child: a longitudinal study of dental development between 3 and 18 years of age.* Cambridge, Mass: Harvard University Press. 1959;
26. Knott VB. Longitudinal study of dental arch widths at four stages of dentition. *Angle Orthod.* 1972; 42:387–394. [[PubMed Citation](#)]
27. Björk A, Skieller V. Growth in width of the maxilla studied by the implant method. *Scand J Plast Reconstr Surg.* 1974; 8:26–33. [[PubMed Citation](#)]
28. Mann RW, Symes SA, Bass WM. Maxillary suture obliteration: aging the human skeleton based on intact or fragmentary maxilla. *J Forensic Sci.* 1987; 32:148–157. [[PubMed Citation](#)]
29. Harris EF. A longitudinal study of arch size and form in untreated adults. *Am J Orthod Dentofacial Orthop.* 1997; 111:419–427. [[PubMed Citation](#)]
30. Picton DC, Moss JP. The part played by the trans-septal fibre system in experimental approximal drift of the cheek teeth of monkeys (*Macaca irus*). *Arch Oral Biol.* 1973; 18:669–680. [[PubMed Citation](#)]
31. Southard TE, Behrents RG, Tolley EA. The anterior component of force. Part 1. measurement and distribution. *Am J Orthod Dentofacial Orthop.* 1989; 96:493–500. [[PubMed Citation](#)]
32. Southard TE, Behrents RG, Tolley EA. The anterior component of force. Part 2. Relationship with dental malalignment. *Am J Orthod Dentofacial Orthop.* 1989; 97:41–44.
33. Sillman JH. Dimensional changes of the dental arches: longitudinal study from birth to 25 years. *Am J Orthod.* 1964; 50:824–842.
34. Lundström A. Changes in crowding and spacing of the teeth with age. *Dent Pract.* 1968; 19:218–224.
35. Izard G. New method for the determination of the normal arch by the function of the face. *Int J Orthod.* 1927; 13:582–595.
36. Battagel JM. Individualized catenary curves: their relationship to arch form and perimeter. *Br J Orthod.* 1996; 23:21–28. [[PubMed Citation](#)]

Variable	White Male		White Female		Black Male		Black Female		F Ratios		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Race	Sex	Interaction
Arch Widths											
I2 to I2	26.4	2.4	25.4	2.3	28.3	2.7	28.2	2.9	67.3*	3.3	2.4
C to C	33.8	2.7	32.6	2.4	36.7	2.8	35.8	2.7	91.9*	10.8*	0.3
P1 to P1	39.3	3.2	37.7	2.8	43.4	3.3	42.5	3.2	165.1*	13.8*	1.3
P2 to P2	45.0	3.2	43.1	2.8	48.1	3.8	46.7	3.6	75.7*	16.9*	0.5
M1 to M1	50.1	2.9	48.0	3.2	53.4	3.2	52.3	2.9	126.6*	22.4*	2.9
M2 to M2	55.7	3.1	53.7	2.7	59.0	3.7	57.4	3.3	76.3*	21.1*	0.3
Arch Depths											
I1 to I2	6.7	1.8	6.3	2.2	7.2	1.9	7.0	2.1	6.9*	2.1	0.1
I1 to C	8.4	2.3	8.0	2.2	9.6	2.8	9.3	2.5	17.6*	1.3	0.0
I1 to P1	15.8	2.5	15.1	2.3	17.7	3.2	17.1	2.9	38.1*	4.8	0.0
I1 to P2	22.1	2.8	21.3	2.4	24.6	3.2	23.9	3.0	58.2*	4.3	0.0
I1 to M1	28.2	2.8	27.2	2.4	31.2	3.5	30.4	3.3	84.9*	6.3	0.1
I1 to M2	38.7	2.9	37.9	2.8	43.0	3.5	41.7	3.6	108.2*	7.2*	0.2
Polynomial Coefficients											
$b_1 \times 10^2$	-2.8	25.4	-4.4	21.5	0.9	34.7	-5.1	31.6	0.1	0.9	0.3
$b_2 \times 10^1$	-9.9	1.6	-9.6	1.3	-11.0	1.9	-10.7	1.8	22.9*	1.2	0.0
$b_3 \times 10^0$	-0.9	3.7	0.3	3.2	-0.3	5.3	0.9	5.4	0.8	3.6	0.0
$b_4 \times 10^0$	4.2	2.3	4.0	2.0	5.0	2.9	5.0	2.6	5.8	0.1	0.1
Arch Surface											
Arch Perimeter	128.6	6.7	125.0	8.6	138.6	6.8	134.5	6.1	94.1*	14.8*	0.1
Arch Area	1,827.8	175.0	1,673.0	179.2	2,135.7	219.3	2,035.2	168.8	161.6*	23.5*	1.1


^a Variables are in millimeters, except polynomials, which are unitless, and arch area (mm²). Sample sizes are listed in the test.
* $P < .01$ (2-tailed).

TABLE 2. Descriptive Statistics and Results of Analyses of Variance of Arch Shape^a

Variable	White Male		White Female		Black Male		Black Female		F Ratios		
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Race	Sex	Interaction
Palatal index	69.3	6.2	70.2	5.1	72.6	6.6	72.5	6.4	13.8*	0.2	0.4
3-3/7-7 ratio	60.7	4.1	60.8	4.5	62.8	5.7	62.5	5.2	8.2*	0.0	0.1


^a These ratios are unitless. Sample sizes are listed in the text.
* $P < .01$ (2-tailed).

FIGURES [Return to TOC](#)




[Click on thumbnail for full-sized image.](#)

FIGURE 1. Diagrams showing maxillary dental landmarks and arch measurements. (A) Eighteen dental landmarks (buccal cusp tips and incisor line angles) were marked on each cast, along with the midpalatal raphe, prior to making an occlusal print. (B) Six arch depths were calculated (measured parallel to the midpalatal raphe) using the mesial line angle of the central incisor as the anterior limit of the arch. (C) Six arch widths were measured. (D) Cross-sectional arch area was measured by fitting a spline function through 14 points (note that the midpoints of the incisor line angles were used) and a straight line connected the mesiobuccal cusp tips of the second molars. Arch perimeter was the spline distance around the arch (omitting the M2-M2 base)




[Click on thumbnail for full-sized image.](#)

FIGURE 2. Average arch forms for black and white men (left) and women (right). Arches are registered on the mesial limit of the central incisors. From the midpoint backwards, the landmarks are: mesial line angle of the central incisor, distal line angle of the central incisor, distal line angle of the lateral incisor, and then the buccal cusp tips from the canine through the second molar



Click on thumbnail for full-sized image.

FIGURE 3. Superimposed average arch forms for black and white men and women. Registration is on the mesial line angle of the central incisor. The size difference between men and women within each ethnic group is much less than the difference between blacks and whites. Note that the greatest transverse distance between blacks and whites is at the first premolar



Click on thumbnail for full-sized image.

FIGURE 4. Average arch forms for black and white men aligned at the canines to emphasize the greater depth of the anterior segment of the arch in blacks as well as the narrower, more tapered form in whites

^aDental Student, University of Tennessee, Memphis, Tennessee.

^bProfessor, Orthodontics, Pediatric Dentistry and Community Health, University of Tennessee, Memphis, Tennessee.

Corresponding author: Edward F. Harris, PhD, Department of Orthodontics, College of Dentistry, 875 Union Avenue, University of Tennessee, Memphis, TN 38163 (E-mail: eharris@utmem.edu).