

[\[Print Version\]](#)

[\[PubMed Citation\]](#) [\[Related Articles in PubMed\]](#)

*The Angle Orthodontist*: Vol. 68, No. 1, pp. 85–90.

## Case Report: Facial asymmetry and early condylar fracture

Takashi Yamashiro, DDS, PhD;<sup>a, b</sup> Takumi Okada, DDS;<sup>c</sup> Kenji Takada, DDS, PhD<sup>d</sup>

<sup>a</sup>Takashi Yamashiro, Department of Orthodontics, Okayama University Dental School, Shikata-cho 2-5-1, Okayama City, Okayama 700, Japan, Tel: +81-86-235-6691, Fax: +81-86-223-7317, E-mail: [yamat@dent.okayama-u.ac.jp](mailto:yamat@dent.okayama-u.ac.jp)

<sup>b</sup>Takashi Yamashiro, assistant professor, Department of Orthodontics, Okayama University Dental School, Okayama, Japan.

<sup>c</sup>Takumi Okada, private practice, Neyagawa, Osaka, Japan.

<sup>d</sup>Kenji Takada, professor and chairman, Department of Orthodontics, Faculty of Dentistry, Osaka University, Osaka, Japan.

### ABSTRACT

This report describes the surgical-orthodontic treatment of two adult females who exhibited dentofacial asymmetry. Their medical histories included early condylar fractures. Growth was disturbed on the fractured side, resulting in mandibular deviation and maxillary canting. Following the completion of growth, the asymmetries were corrected by combined orthodontic-surgical treatment. The treatment planning procedures and the results are presented.

**KEY WORDS:** Facial asymmetry, Condylar fracture.

Submitted: February 1996

Accepted: August 1996.