

[\[Print Version\]](#)

[\[PubMed Citation\]](#) [\[Related Articles in PubMed\]](#)

The Angle Orthodontist: Vol. 66, No. 6, pp. 457–462.

Anterior arch circumference adjustment - how much?

C.L. Steyn, BChD, MChD;^b A.M.P. Harris, BChD, HonsBSc, MChD, FFD(SA)Orth;^a R.J. du Preez, BSc(Eng), MS, DrEng^d

^aProf A.M.P. Harris, Department of Orthodontics, Faculty of Dentistry, University of Stellenbosch, Private Bag X1, Tygerberg 7505, Republic of South Africa. A.M.P. Harris, professor and head, Department of Orthodontics, Faculty of Dentistry, University of Stellenbosch.

^bC.L. Steyn, part-time lecturer, Department of Orthodontics, Faculty of Dentistry, University of Stellenbosch, and orthodontist in private practice.

^dR.J. du Preez, professor and head, Center for Computational Mechanics, University of Pretoria

ABSTRACT

A table is presented by means of which changes in arch length and arch circumference can be more accurately predicted, thereby simplifying treatment planning for cases with special problems, e.g. a Bolton discrepancy. A fundamental condition for the use of this table is that the anterior arch from canine to canine forms a parabolic curve.

KEY WORDS: Space analysis, Interproximal stripping, Overjet correction, Bolton discrepancy, Missing teeth.

Submitted: June 1995

Accepted: August 1995.