

[\[Print Version\]](#)

[\[PubMed Citation\]](#) [\[Related Articles in PubMed\]](#)

The Angle Orthodontist: Vol. 56, No. 2, pp. 164–175.

Occurrence and Distribution of Root Resorption in Orthodontically moved Premolars in Dogs

Marie E. Follin;^a Ingvar Ericsson; Birgit Thilander

^aDr. Marie Follin, Department of Orthodontics, Faculty of Odontology, Box 33070, 5-400 33 Goteborg, SWEDEN

ABSTRACT

An experimental study in dogs, in which 15 teeth were bodily moved and 9 teeth were tipped with the same amount of horizontal force. Less resorption was found with tipping, and less was found on the mesial root surfaces moved into the extraction sites than on the furcation side of the distal roots.

Dr. Follin is a community orthodontic consultant and a postgraduate research student at the Department of Orthodontics in Göteborg. She holds a D.D.S. and an M.S. degree in Orthodontics from the University of Göteborg, Sweden

Dr. Ericsson is Associate Professor in the Department of Periodontology, University of Göteborg. He holds D.D.S., Ph.D., and M.S. degrees in Periodontology and Prosthodontics from the University of Göteborg, Sweden

Dr. Thilander is Professor and Chairman of the Department of Orthodontics, University of Göteborg. She holds D.D.S., M.S. and Ph.D. degrees from the Caroline Institute Stockholm, Sweden

KEY WORDS: ROOT RESORPTION, ORTHODONTIC MOVEMENT, DOG.