

[\[Print Version\]](#)

[\[PubMed Citation\]](#) [\[Related Articles in PubMed\]](#)

The Angle Orthodontist: Vol. 52, No. 1, pp. 11–18.

Extreme Distal Migration of the Mandibular Second Bicuspid

A variant of eruption

STEPHEN R. MATTESON;^a MEL L. KANTOR; WILLIAM R. PROFFIT

^aDr. Stephen R. Matteson, Section of Oral Radiology, School of Dentistry, University of North Carolina, Chapel Hill, N.C. 27514

ABSTRACT

Evaluation of more than 25,000 panoramic radiographs shows that distal migration of an unerupted lower second bicuspid into a first molar extraction site occurs with sufficient frequency to require consideration of this possibility in cases of early molar loss.

Dr. Matteson is Associate Professor and Head of the Section of Oral Radiology at the University of North Carolina School of Dentistry. He is a dental graduate (D.D.S.) of the University of Pennsylvania School of Dental Medicine

Dr. Kantor is a general practice resident at the Hennepin County Medical Center, Minneapolis, Minnesota. He is a dental graduate (D.D.S.) of the University of North Carolina

Dr. Proffit is Professor and Chairman of the Department of Orthodontics at the University of North Carolina. He is a dental graduate (D.D.S.) of the University of North Carolina and holds a Ph.D. degree from the Medical College of Virginia