

Journal of Andrology, Vol. 23, No. 6, November/December 2002  
Copyright © [American Society of Andrology](#)

# Inhibition of Alkaline Phosphatase Activity of Boar Semen by Pentoxifylline, Caffeine, and Theophylline

JAN GLOGOWSKI<sup>\*</sup>, DOUGLAS R. DANFORTH<sup>†</sup> AND ANDRZEJ CIERESZKO<sup>‡</sup>

*From the <sup>\*</sup> Department of Molecular Andrology, and <sup>†</sup> Department of Semen Biology, Institute of Animal Reproduction and Food Research, Polish Academy of Sciences, Olsztyn, Poland; and <sup>‡</sup> Department of Obstetrics and Gynecology, College of Medicine, The Ohio State University, Columbus, Ohio.*

Correspondence to: Andrzej Ciereszko, PhD, Institute of Animal Reproduction and Food Research, Polish Academy of Sciences, Department of Semen Biology, Tuwima 10, 10-747 Olsztyn, Poland (e-mail: [acieresz@pan.olsztyn.pl](mailto:acieresz@pan.olsztyn.pl)).

Received for publication November 1, 2001; accepted for publication April 2, 2002.

## Abstract

Methyl xanthines have been used frequently as additives to sperm suspensions in order to improve sperm characteristics. The mechanism of action on spermatozoa is generally assumed to be inhibition of sperm phosphodiesterase activity, resulting in elevation of complementary adenosine monophosphate levels in spermatozoa. The present study was designed to examine the effect of methyl xanthines (pentoxifylline, caffeine, and theophylline) on another important enzyme system, alkaline phosphatase, in boar seminal plasma and spermatozoa. Inhibition of sperm alkaline phosphatase could be distinguished from that of seminal plasma by a paradoxical stimulation by pentoxifylline at lower pH values in spermatozoa. Among the three methyl xanthines, theophylline exhibited the most dramatic inhibition of alkaline phosphatase activity and substrate inhibition was observed with increasing concentrations. Each methyl xanthine had a different action on alkaline phosphatase activity at lower pH; theophylline showed the highest inhibition, caffeine inhibition was not related to pH, and pentoxifylline did not inhibit alkaline phosphatase of seminal plasma and, in fact, it stimulated its activity (or that of a phosphatase with lower pH optimum) in spermatozoa. These results indicate another possible mechanism of action of methyl xanthines on sperm and are in agreement with data indicating that methyl xanthines are not specific inhibitors of sperm phosphodiesterase, because clearly, they inhibit alkaline phosphatase activity as well.

**Key words:** Methyl xanthines, seminal plasma, spermatozoa

### This Article

- ▶ [Abstract](#) **FREE**
- ▶ [Full Text \(PDF\)](#)
- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me if a correction is posted](#)

### Services

- ▶ [Similar articles in this journal](#)
- ▶ [Similar articles in PubMed](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Download to citation manager](#)

### Citing Articles

- ▶ [Citing Articles via Google Scholar](#)

### Google Scholar

- ▶ [Articles by Glogowski, J.](#)
- ▶ [Articles by Ciereszko, A.](#)
- ▶ [Search for Related Content](#)

### PubMed

- ▶ [PubMed Citation](#)
- ▶ [Articles by Glogowski, J.](#)
- ▶ [Articles by Ciereszko, A.](#)

### ▲ [Top](#)

- [Abstract](#)
- ▼ [Materials and Methods](#)
- ▼ [Results](#)
- ▼ [Discussion](#)
- ▼ [References](#)

Methyl xanthines are frequently used as additives to sperm suspensions in order to improve sperm characteristics. Among methyl xanthines, pentoxifylline (3,7-dimethyl-1-(5-oxo-hexyl)-xanthine),

caffeine (1,3,7-trimethylxanthine), and theophylline (1,3-dimethylxanthine) have been employed most often. In numerous studies, methyl xanthine supplementation resulted in better motility characteristics in fresh and cryopreserved spermatozoa. These include a greater percentage of motile cells and greater velocity of sperm movement, including curvilinear velocity, which often leads to a higher percentage of nonhyperactive sperm shifting to either a transitional or hyperactivated state ([Tash and Means, 1983](#); [Schoff and Lardy, 1987](#); [Rees et al, 1990](#); [Sikka and Hellstrom 1991](#); [Coward et al, 1994](#); [Lewis et al, 1994](#); [Koutsarova et al, 1997](#); [Merino et al, 1997](#); [Sharma and Agarwal, 1997](#); [Nassar et al, 1999a](#); [Ponce et al, 1999](#)).

In immature epididymal or testicular spermatozoa, addition of methyl xanthine is crucial for acquisition or improvement of sperm motility and fertilizing ability ([Gould et al, 1988](#); [Jaiswal and Majumder, 1996](#); [Mahony et al, 1996](#); [Angelopoulos et al, 1999](#)). Stimulatory effects of methyl xanthines on capacitation and the acrosome reaction have also been demonstrated ([Tesarik et al, 1992](#); [Tesarik and Mendoza, 1993](#); [DasGupta et al, 1994](#); [Jayaprakash et al, 1997](#); [Esteves et al, 1998](#); [Ain et al, 1999](#)). Apart from modulation of sperm function, a protective role on sperm membranes by pentoxifylline has been described ([Ponce et al, 1999](#)). This effect may be ascribed to neutralization of reactive oxygen species and a reduction of lipid peroxidation ([Bell et al, 1993](#); [McKinney et al, 1996](#); [Okada et al, 1997](#)). Pentoxifylline has been proposed as a cryoprotectant owing to its protective action ([Wang et al, 1993](#); Brennen and Holden, 1995). Overall, the addition of methyl xanthines to sperm suspensions seems to improve sperm function, leading to better sperm fertilizing capacity ([Fraser, 1979](#); Loughlin and Agarwal, 1992; [Nagai et al, 1994](#); [Negri et al, 1996](#); [Abeydeera and Day, 1997](#); [Chauhan et al, 1998](#); [Nassar et al, 1999a](#)). However, the beneficial effects of methyl xanthine on sperm quality have been questioned (Lewis et al, 1993, 1994; [Mathieu et al, 1994](#); [Tournaye et al, 1994](#); [Dimitriadou et al, 1995](#); [Paul et al, 1996](#)).

The generally accepted mechanism of methyl xanthines is their inhibition of sperm cyclic nucleotide phosphodiesterase. This reduces destruction of cyclic adenosine 3',5' monophosphate (cAMP), which results in elevation of cyclic adenosine monophosphate levels in spermatozoa ([Tash and Means, 1983](#)). Indeed, a rise in cAMP levels in spermatozoa after methyl xanthine treatment has been consistently demonstrated ([Garbers et al, 1971](#); [Hoskins et al, 1975](#); [Wang et al, 1993](#)). Elevated levels of cAMP enhance cAMP-dependent processes of spermatozoa, including motility, capacitation, and acrosome reaction ([Tash and Means, 1983](#); [Monks and Fraser, 1987](#); [Fraser and Monks, 1990](#); [Armstrong et al, 1994](#); [Aitken, 1997](#)). The molecular basis of cAMP regulation of these processes is based on protein phosphorylation, especially for capacitation ([Duncan and Fraser, 1993](#); [Galantino-Homer et al, 1997](#); [Flesch et al, 1999](#); [Nassar et al, 1999b](#)). An alternate way for methyl xanthines to elevate cAMP levels is through the modulation of adenosine receptors ([Vijayaraghavan and Hoskins, 1986](#); Loughlin and Agarwal, 1992). Methyl xanthines may also affect translocation of intracellular calcium (Loughlin and Agarwal, 1992; [Nagai et al, 1994](#)), neutralization of reactive oxygen species, and reduced lipid peroxidation. These observations support the hypothesis that non-cAMP-mediated events in spermatozoa may also be modulated by methyl xanthines ([Vandevoort et al, 1994](#)).

It is well known that methyl xanthines can inhibit alkaline phosphatase of somatic origin ([Vinet et al, 1978](#); [Dai and Snow, 1991](#); [Wang and Gilles-Baillien, 1992](#); [Rezende et al, 1998](#)). This effect on somatic alkaline phosphatase is employed in determining theophylline in serum (Jourquin and Kauffman, 1998). These data raise the question of whether methyl xanthines may also inhibit alkaline phosphatase in mammalian semen, where it is universally present ([Bell and Lake, 1962](#); [Jones, 1978](#); [Glogowski, 1988](#); [Tang 1998](#)). If they do, their effects exerted on sperm functions may also be mediated through modulation of alkaline phosphatase activity. In this study, we tested the effects of pentoxifylline, caffeine, and theophylline on the alkaline phosphatase activity of boar seminal plasma and spermatozoa.

### *Animals, Preparation of Seminal Plasma and Sperm Extracts, and Alkaline Phosphatase Assay*

Semen was collected from adult boars housed in the Animal Breeding and Insemination Station in Olsztyn, Poland. Semen was collected manually into a cup with a filter, and the gel fraction was discarded. Seminal plasma was obtained by centrifugation (10 000 x *g*). Plasma was diluted 100-150 times before alkaline phosphatase assay. Sperm extracts were obtained as follows: semen (8 mL) was washed 3 times with 0.055 M HEPES-NaOH, 0.7% NaCl pH 7.4, and then 2.5 mL of cold HEPES buffer (without NaCl) was added. Sperm suspensions were stored on ice for 30 minutes, sonicated for 25 seconds, and centrifuged (10 000 x *g*). Pellets were discarded and supernatants were diluted 1:1 with HEPES buffer containing 1.5% NaCl. Extracts were frozen and stored for up to 3 days at -20° C and were not further diluted for alkaline phosphatase assay. Alkaline phosphatase activity was measured with a colorimetric assay with 4-nitrophenylphosphate (Merck & Company, Whitehouse Station, NJ) as a substrate according to the method described by Bessey et al (1946). The incubation mixture contained a 1.1 mL final volume of 50 mM glycine buffer and 7.5 mM 4-nitrophenylphosphate (disodium salt). Incubations were carried out at 37° C for 30 minutes, and the reaction was terminated by adding 10 mL of 0.02 N NaOH. Amounts of liberated 4-nitrophenol were calculated from standard curves. Alkaline phosphatase activity was expressed as micromolar units of 4-nitrophenol liberated per minute. In total, seminal plasma and sperm extracts from 5 different boars were used.

### *Effects of Methyl Xanthines on Alkaline Phosphatase Activity of Seminal Plasma and Spermatozoa*

Caffeine and pentoxifylline (Sigma Chemical Company, St Louis, Mo) effects were tested in the concentration range of 0-20 mM and theophylline (Sigma) in the range of 0-2.5 mM. Substrate and inhibitors were added first and preincubated at 37° C for 5 minutes. Reactions were started by adding an enzyme source (either seminal plasma or sperm extract).

### *Effect of pH on Inhibition of Seminal Plasma and Sperm Alkaline Phosphatase Activity by Methyl Xanthines*

Alkaline phosphatase activities were tested in the presence of 10 mM caffeine and pentoxifylline, and 0.625 mM of theophylline using 0.05 M (final concentration) glycine-NaOH buffer in the pH range of 8.6-11.0.

### *Effects of Substrate Concentration on Inhibition of Seminal Plasma and Sperm Alkaline Phosphatase Activity by Methyl Xanthines*

A 4-nitrophenylphosphate concentration range of 0-5 mM was used. Alkaline phosphatase activities were measured in the presence of 10 mM caffeine and pentoxifylline, and 0.625 mM of theophylline. The pH of the reaction mixture was 10.6 for variants with caffeine and pentoxifylline, and 10.2 for theophylline (owing to greater inhibition at pH 10.2 than 10.6).

### *Statistical Analysis*

Data are expressed as means  $\pm$  SEM (*n* = 5). One-way analysis of variance (ANOVA) was used for evaluating different concentrations of methyl xanthines on alkaline phosphatase activities and two-way ANOVA for evaluation inhibition at different pHs. A Tukey test was used for post hoc comparisons. Concentrations of theophylline for 50% inhibition of alkaline phosphatase (IC<sub>50</sub>) and regressions of Lineweaver-Burk plots were calculated using the GraphPad PRISM statistical package

## Results

- [▲ Top](#)
- [▲ Abstract](#)
- [▲ Materials and Methods](#)
- Results
- [▼ Discussion](#)
- [▼ References](#)

### Effects of Methyl Xanthines on Alkaline Phosphatase Activity of Seminal Plasma and Spermatozoa

All methyl xanthines inhibited alkaline phosphatase activity, whether they were derived from seminal plasma or spermatozoa, in a concentration-dependent manner ([Figure 1](#)). Theophylline was more effective than pentoxifylline and caffeine; the dose response line for theophylline was much steeper than it was for other methyl xanthines. Alkaline phosphatase inhibition was 27% and 37% for seminal plasma and 35% and 43% for spermatozoa, at 20 mM pentoxifylline and 20 mM caffeine, respectively. In contrast, 50% alkaline phosphatase inhibition was observed at theophylline concentrations of less than 1 mM. This inhibition was more effective for seminal plasma alkaline phosphatase than it was for spermatozoa alkaline phosphatase activities. The  $IC_{50}$  for seminal plasma alkaline phosphatase activity was  $0.34 \pm 0.02$  mM, and for sperm alkaline phosphatase it was  $0.62 \pm 0.05$  mM.




Figure 1. Effects of caffeine (0-20 mM), pentoxifylline (0-20 mM), and theophylline (0-2.5 mM) on alkaline phosphatase activity of boar seminal plasma and spermatozoa ( $n = 5$ ,  $* P < .01$ ).

View larger version  
(25K):  
[\[in this window\]](#)  
[\[in a new window\]](#)

### Effect of pH on Inhibition of Seminal Plasma and Sperm Alkaline Phosphatase Activity by Methyl Xanthines

Inhibition of both seminal plasma and sperm alkaline phosphatase activities was affected by pH (Figures [2](#) and [3](#)), however, the effects were different for particular methyl xanthines (Figures [2D](#) and [3D](#)), especially at lower pH. Inhibition of alkaline phosphatase by theophylline increased with decreasing pH. In contrast, inhibition by caffeine was not affected by pH, and pentoxifylline inhibition was greatest at high pH ( $>9.6$  for seminal plasma and  $>10.3$  for spermatozoa). At a lower pH, pentoxifylline did not inhibit seminal plasma alkaline phosphatase activity, but it stimulated sperm alkaline phosphatase activity.




Figure 2. Effects of pentoxifylline (**A**, 10 mM), caffeine (**B**, 10 mM), and theophylline (**C**, 0.625 mM) on alkaline phosphatase activity of boar seminal plasma, measured at the pH range of 8.6-11.0. Percentage of inhibition at different pHs are presented in (**D**);  $n = 5$ , \*  $P < .01$ .

View larger version  
(32K):  
[\[in this window\]](#)  
[\[in a new window\]](#)




Figure 3. Effects of pentoxifylline (**A**, 10 mM), caffeine (**B**, 10 mM), and theophylline (**C**, 0.625 mM) on alkaline phosphatase activity of boar spermatozoa measured at the pH range of 8.6-11.0. Percentage of inhibition at different pHs are presented in (**D**);  $n = 5$ , \*  $P < .01$ .

View larger version  
(32K):  
[\[in this window\]](#)  
[\[in a new window\]](#)

### Effects of Substrate Concentration on Inhibition of Seminal Plasma and Sperm Alkaline Phosphatase Activity by Methyl Xanthines

Pentoxifylline and caffeine inhibited alkaline phosphatase activities of seminal plasma and spermatozoa in a non-competitive manner (Figures 4 and 5). In contrast to theophylline, a substrate inhibition of alkaline phosphatase was observed.




Figure 4. Effects of 4-nitrophenylphosphate concentrations (range 0-5 mM) in the presence of caffeine (10 mM), pentoxifylline (10 mM), and theophylline (0.625 mM) on alkaline phosphatase activity of boar seminal plasma. Double reciprocal plots are presented in right panels;  $n = 5$ .



View larger version  
(22K):

[\[in this window\]](#)  
[\[in a new window\]](#)




Figure 5. Effects of 4-nitrophenylphosphate concentrations (range 0-5 mM) in the presence of caffeine (10 mM), pentoxifylline (10 mM), and theophylline (0.625 mM) on alkaline phosphatase activity of boar spermatozoa. Double reciprocal plots are presented in right panels; n = 5.

View larger version  
(20K):

[\[in this window\]](#)  
[\[in a new window\]](#)

## Discussion

- ▲ [Top](#)
- ▲ [Abstract](#)
- ▲ [Materials and Methods](#)
- ▲ [Results](#)
- [Discussion](#)
- ▼ [References](#)

This study demonstrates that methyl xanthines can inhibit alkaline phosphatase activity in seminal plasma and spermatozoa. Our study revealed differences among inhibition modes of 2 sources of alkaline phosphatase (seminal plasma and spermatozoa) as well as among 3 methyl xanthines.

Inhibition of sperm alkaline phosphatase could be distinguished from that of seminal plasma by stimulating the activity with pentoxifylline at lower pH values. Among the 3 methyl xanthines, theophylline exhibited the most dramatic inhibition of alkaline phosphatase activity, and substrate inhibition was observed with increasing concentration. Each methyl xanthine had a different action at lower pH: theophylline showed highest inhibition, caffeine inhibition was not related to pH, and pentoxifylline did not inhibit seminal plasma alkaline phosphatase, but it stimulated spermatozoa alkaline phosphatase activity. These results indicate another possible action of methyl xanthines on sperm and are in agreement with previous studies indicating that methyl xanthines are not pure inhibitors of phosphodiesterase.

There are some similarities between semen alkaline phosphatase and phosphodiesterase. Like alkaline phosphatase, phosphodiesterase is present in spermatozoa and seminal plasma, it occurs in multiple molecular forms, and it is differentially inhibited by methyl xanthines ([Tash, 1976](#)). Vijayaraghavan and Hoskins ([1986](#)) found that the theophylline  $IC_{50}$  for inhibiting bovine sperm phosphodiesterase was 0.7 mM, which is within the same range as inhibition obtained in this work for boar alkaline phosphatase. As with boar sperm alkaline phosphatase inhibition, theophylline is a more effective inhibitor of phosphodiesterase found in the sperm of sea urchins ([Wells and Garbers, 1976](#)) and buffalo ([Bhatangar et al, 1979](#)). On the other hand, some species-specific differences were reported: in ram semen, for example, caffeine was a more potent inhibitor of phosphodiesterase than theophylline, and both methyl xanthines inhibited phosphodiesterase by 33%-79% at concentrations of

5 mM ([Tash, 1976](#)). Therefore, interspecies comparisons of the effects of methyl xanthine on phosphodiesterase and alkaline phosphatase activity should be interpreted cautiously.

It is difficult to translate the alkaline phosphatase inhibition by methyl xanthines described here into specific clinical applications. Concentrations of methyl xanthines reported in our work were similar to those used for enhancement of sperm quality. For example, enhancement of semen characteristics has been reported at concentrations of caffeine and pentoxifylline up to 20 mM ([Stachecki et al, 1994](#)), and theophylline has been used at concentrations of 10 mM and 20 mM ([Loughlin and Agarwal, 1992](#); [Coward et al, 1994](#)) or even 30 mM ([Jaiswal and Majumder, 1996](#)). If sperm alkaline phosphatase of other mammalian species is similarly inhibited by methyl xanthines, as we have demonstrated for boar alkaline phosphatase in the present study, these data suggest that beneficial effects of these substances may be partially related to their action on sperm alkaline phosphatase. This inhibition may not always produce better sperm characteristics because a decrease in sperm quality at high concentrations of methyl xanthine has been reported ([Brennan and Holden, 1995](#)). These authors reported the detrimental effect of a high dose of pentoxifylline (10 mM) on acrosome morphology. Also, Armstrong et al ([1994](#)) found that 5 mM caffeine or pentoxifylline stimulated the motility of rat sperm more than 10 mM did. In contrast, improvement of sperm characteristics has been reported at concentrations of methyl xanthines that, in the present study, do not significantly affect boar sperm alkaline phosphatase. For example, common doses of pentoxifylline used for supplementation of human sperm suspensions are 1 mM or 3-4 mM ([Wang et al, 1993](#); [Lewis et al, 1994](#); [Brennan and Holden, 1995](#); [McKinney et al, 1996](#)) doses lower than those that were inhibitory in the present study. In such cases, inhibition of alkaline phosphatase may not be involved in the beneficial effects of pentoxifylline.

We recognize that the experiments included in the present study were performed at artificially elevated (non-physiological) pH values owing to the alkaline optimum of alkaline phosphatase activity. For this reason, extrapolation of alkaline phosphatase inhibition at more physiological (neutral or slightly alkaline) pHs requires caution. Our data indicate that theophylline is the most effective inhibitor of alkaline phosphatase activity at lower pH values, whereas pentoxifylline stimulates alkaline phosphatase activity (or a different phosphatase with an optimum pH near neutral pH [see below]). Most experiments investigating methyl xanthines have been performed using human sperm, and there are substantial differences in seminal plasma alkaline phosphatase activities among mammals. The highest levels of alkaline phosphatase enzyme activities are found in boars and the lowest are found in humans ([Bell and Lake, 1962](#); [Jones, 1978](#)). Thus, if methyl xanthines act on sperm function through modulation of alkaline phosphatase, effects on human sperm may not be as profound as those observed in the present study using boar sperm.

Despite extensive research on alkaline phosphatases in the male reproductive tract, their role in reproductive physiology is not clear. Difficulties in understanding the role of alkaline phosphatase originate from the universal presence of alkaline phosphatase in reproductive tissue and its wide substrate specificity. Alkaline phosphatase is present in both seminal plasma and spermatozoa. In seminal plasma, it is present in multiple forms that share similar kinetic properties, and it generally originates from epididymal fluid ([Strzezek and Glogowski, 1979](#); [Glogowski and Strzezek, 1980](#); [Frenette et al, 1986](#); [Glogowski, 1988](#); [Iyer et al, 1988](#); [Tang, 1998](#)). Alkaline phosphatase is present in prostasomes ([Fabiani and Ronquist, 1995](#)) and is a component of sperm plasma membranes ([Soucek and Vary, 1984](#); [Parks et al, 1987](#)). In addition, it is associated with cytoplasmic droplets and acrosomes ([Moniem and Glover, 1972](#); [Bavdek and Glover, 1970](#); [Yuan et al, 1995](#)). Alkaline phosphatase derived from semen can hydrolyze phosphate esters of various mononucleotides, sugars, glycerophosphate ([Glogowski, 1988](#)), and pyridoxal 5'-phosphate ([Glogowski, 1988](#); [Ciereszko et al, 1994](#)), as well as adenosine triphosphate ([Glogowski, 1988](#)). Minelli et al ([1995](#)) suggest a possible

role for alkaline phosphatase in the dephosphorylation of adenosine monophosphate. Tang (1998) suggested that alkaline phosphatase may inhibit the glycosylation of sperm surface glycoproteins through the hydrolysis of nucleotide sugars. The existence of multiple possible modes of alkaline phosphatase enzymatic actions makes it difficult to identify the specific functions of alkaline phosphatase in semen and to relate inhibition by methyl xanthine to specific sperm functions. In other tissues, membrane-bound alkaline phosphatase has been associated with chloride channels (Becq et al, 1993). Such channels are present in spermatozoa and are presumably involved in the acrosome reaction (Meizel, 1997). The possible link between sperm alkaline phosphatase and chloride channels deserves further study.

Our study revealed considerable differences between theophylline and 2 other methyl xanthines in their ability to inhibit alkaline phosphatase. This finding suggests that if inhibition of alkaline phosphatase is important for normal boar sperm physiology, a difference in mechanism of action on sperm should be seen between theophylline and caffeine or pentoxifylline. Such differences have not been reported for other species. However, to our knowledge, no comparative studies have been performed on the effects of all 3 methyl xanthines on boar sperm physiology. For this reason, it is difficult at present to relate inhibition of alkaline phosphatase to sperm function in the boar.

Differences in methyl xanthine action may be related to differences in the structures of a particular methyl xanthine. Theophylline lacks a methyl group at carbon 7, and this absence may play a critical role in its having the highest effectiveness among methyl xanthines in inhibiting alkaline phosphatase. On the other hand, it is possible that methyl xanthine in sperm suspensions may induce changes in membrane structure that may stimulate conformational changes of membrane proteins and possibly by modulating their function. Sato et al (1991) indicated that pentoxifylline can change the fluidity of erythrocyte membranes. It is interesting that caffeine and theophylline did not induce such fluidity changes. Because of the critical role of changes in sperm membranes in sperm functions, it needs to be established whether pentoxifylline may act by inducing changes in the fluidity of sperm membranes, and whether these changes may modulate sperm alkaline phosphatase. We used unpurified material for our analysis, and it is possible that stimulation by pentoxifylline might be due to the presence of a different phosphatase with a pH maximum near neutral pH, and that differences between sperm and seminal plasma could thus be explained by differing amounts of this phosphatase. For this reason, further studies using a purified preparation of phosphatases are necessary to define the target phosphatase for pentoxifylline. Such studies would also further our understanding of the mechanisms of alkaline phosphatase inhibition by methyl xanthines, especially theophylline.

Our data demonstrate that methyl xanthines can inhibit alkaline phosphatase activity of seminal plasma and spermatozoa. Methyl xanthine inhibition of alkaline phosphatase activity was specific to pH, substrate, and tissue. These data suggest that, in addition to the classical effect on phosphodiesterase, methyl xanthines may also modulate alkaline phosphatase activity in spermatozoa and seminal plasma.

## ***Acknowledgments***

We thank Mrs Dorota Kubiak for excellent technical assistance and the personnel of the Animal Breeding Station in Olsztyn for providing boar semen.

## ***References***



<a href="#">▲ Top</a>
<a href="#">▲ Abstract</a>
<a href="#">▲ Materials and Methods</a>
<a href="#">▲ Results</a>
<a href="#">▲ Discussion</a>
▪ <a href="#">References</a>

Abeydeera LR, Day BN. In vitro penetration of pig oocytes in a modified tris-buffered medium: effect of BSA, caffeine and calcium. *Theriogenology*.1997; 48:537 -544.

Ain R, Uma-Devi K, Shivaji S, Seshagiri PB. Pentoxifylline-stimulated capacitation and acrosome reaction in hamster spermatozoa: involvement of intracellular signaling molecules. *Mol Hum Reprod*. 1999;5:618 -626. [\[Abstract/Free Full Text\]](#)

Aitken RJ. Molecular mechanisms regulating sperm function. *Mol Hum Reprod*1997; 3:169 -173. [\[Free Full Text\]](#)

Angelopoulos T, Adler A, Krey L, Liccardi F, Noyes N, McCullough A. Enhancement or initiation of testicular sperm motility by in vitro culture of testicular tissue. *Fertil Steril*.1999; 71:240 -243. [\[Medline\]](#)

Armstrong VL, Clulow J, Murdoch RN, Jones RC. Intracellular signal transduction mechanisms of rat epididymal spermatozoa and their relationship to motility and metabolism. *Mol Reprod Dev*.1994; 38:77 -84. [\[Medline\]](#)

Bavdek S, Glover TD. Alkaline phosphatase in the cytoplasmic droplet of rabbit spermatozoa. *J Reprod Fertil*.1970; 22:371 -373.

Becq F, Fanjul M, Merten M, Figarella C, Hollande E, Gola M. Possible regulation of CFTR-chloride channels by membrane-bound phosphatase in pancreatic duct cells. *FEBS Lett*.1993; 327:337 -342. [\[Medline\]](#)

Bell JD, Lake PE. A comparison of phosphomonoesterase activities in the seminal plasmas of domestic cock, turkey tom, boar, bull, buck rabbit and of man. *J Reprod Fertil*.1962; 3:363 -368.

Bell M, Wang R, Hellstrom WJG, Sikka SC. Effect of cryoprotective additives and cryopreservation protocol on sperm membrane lipid peroxidation and recovery of motile spermatozoa. *J Androl*.1993; 14:472 -478. [\[Abstract/Free Full Text\]](#)

Bessey OH, Lowry OH, Brock MJ. A method for rapid determination of alkaline phosphatase with five cubic millimeters of serum. *J Biol Chem*. 1946;164:321 -323. [\[Free Full Text\]](#)

Bhatnagar SK, Chaudhry PS, Anand SR. Adenosine 3',5'-monophosphate phosphodiesterase of buffalo spermatozoa. *J. Reprod Fertil*.1979; 56:133 -139.

Brennan AP, Holden CA. Pentoxifylline-supplemented cryoprotectant improves human sperm motility after cryopreservation. *Hum Reprod*. 1995;10:2308 -2312. [\[Abstract/Free Full Text\]](#)

Chauhan MS, Singla SK, Palta P, Manik RS, Madan ML. Influence of theophylline on cleavage rate and embryonic development following in vitro fertilization of buffalo oocytes. *Indian J Anim Sci*.1998; 68:920 -922.

Ciereszko A, Glogowski J, Demianowicz W, Strzezek J. Stimulation of aspartate aminotransferase from farm animal semen by pyridoxal 5'-phosphate. *Anim Reprod Sci*.1994; 34:327 -341.

Cowart CL, London SN, Vernon MW, Pedigo NG. The effects of cyclic adenosine monophosphate, forskolin, and theophylline on motility parameters in gossypol-treated human sperm. *Fertil Steril*.1994; 61:929 -934. [\[Medline\]](#)

Dai X, Snow D. Differential theophylline inhibition of alkaline phosphatase and 5'-nucleotidase of bovine milk fat globule membranes. *Int J Biochem*.1991; 23:743 -747. [\[Medline\]](#)

DasGupta S, O'Toole C, Mills CL, Fraser LF. Effect of pentoxifylline and progesterone on human sperm capacitation and acrosomal reaction. *Hum Reprod.* 1994; 9:2103 -2109. [\[Abstract/Free Full Text\]](#)

Dimitriadou FD, Rizos D, Mantzavinos T, Arvaniti K, Voutsina K, Prapa A, Kanakas N. The effect of pentoxifylline on sperm motility, oocyte fertilization, embryo quality, and pregnancy outcome in an in vitro fertilization program. *Fertil Steril.* 1995; 63:880 -886. [\[Medline\]](#)

Duncan AE, Fraser LR. Cyclic AMP-dependent phosphorylation of epididymal mouse sperm proteins during capacitation in vitro: identification of an Mr 95,000 phosphotyrosine-containing protein. *J Reprod Fertil.* 1993; 97:287 -299.

Esteves SC, Sharma RK, Thomas AJ, Agarwal A. Cryopreservation of human spermatozoa with pentoxifylline improves the post-thaw agonist-induced acrosome reaction rate. *Hum Reprod.* 1998; 13:3384 -3389. [\[Abstract/Free Full Text\]](#)

Fabiani R, Ronquist G. Association of some hydrolytic enzymes with the prostasome membrane and their differential responses to detergent and PIPLC treatment. *Prostate.* 1995; 27:95 -101. [\[Medline\]](#)

Flesch FM, Colenbrander B, van Golde LMG, Gadella BM. Capacitation induces tyrosine phosphorylation of proteins in the boar sperm plasma membranes. *Biochem Biophys Res Commun.* 1999; 262:787 -792. [\[Medline\]](#)

Fraser LR. Accelerated mouse sperm penetration in vitro in the presence of caffeine. *J Reprod Fertil.* 1979; 57:377 -384.

Fraser LR, Monks NJ. Cyclic nucleotides and mammalian sperm capacitation. *J Reprod Fertil Suppl.* 1990; 42:9 -21. [\[Medline\]](#)

Frenette G, Dube JY, Tremblay RR. Origin of alkaline phosphatase of canine seminal plasma. *Arch Androl.* 1986; 16:235 -241. [\[Medline\]](#)

Galantino-Homer HL, Visconti PE, Kopf GS. Regulation of protein tyrosine phosphorylation during bovine sperm capacitation by a cyclic adenosine 3',5'-monophosphate-dependent pathway. *Biol Reprod.* 1997; 56:707 -719. [\[Abstract\]](#)

Garbers DL, Lust WD, First NL, Lardy HA. Effects of phosphodiesterase inhibitors and cyclic nucleotides on sperm respiration and motility. *Biochemistry.* 1971; 10:1825 -1831.

Glogowski J. Alkaline phosphatase of boar reproductive tract. *Acta Academiae Agriculturae ac Technicae Olstenensis.* 1988; 31(suppl B):1 -53.

Glogowski J, Strzezek J. Molecular forms of alkaline phosphatase of ram seminal plasma. *Anim Reprod Sci.* 1980; 3:307 -323.

Gould KG, Young LG, Hinton BT. Alterations in primate sperm motility with maturation and during exposure to theophylline. *Am J Primatol.* 1988; 15:325 -336.

Hoskins DD, Hall ML, Munsterman D. Induction of motility of immature bovine spermatozoa by cyclic AMP phosphodiesterase inhibitors and seminal plasma. *Biol Reprod.* 1975; 13:168 -176. [\[Abstract\]](#)

Iyer SK, Daron HH, Aull JL. Purification and properties of alkaline phosphatase from boar seminal plasma. *J Reprod Fertil.* 1988; 82:657 -664.

Jaiswal BS, Majumder GC. In-vitro initiation of forward motility in testicular spermatozoa. *Int J Androl.* 1996; 19:97 -102. [\[Medline\]](#)

Jayaprakash D, Kumar KS, Shivaji S, Seshagiri PB. Pentoxifylline induces hyperactivation and

acrosome reaction in spermatozoa of golden hamsters changes in motility kinematics. *Hum Reprod.* 1997; 12: 2192 -2199. [\[Abstract/Free Full Text\]](#)

Jones R. Comparative biochemistry of mammalian epididymal plasma. *Comp Biochem Physiol B.* 1978; 61: 365 -370. [\[Medline\]](#)

Jourquin G, Kauffmann J-M. Fluorometric determination of theophylline in serum by inhibition of bovine alkaline phosphatase in AOT based water/oil microemulsion. *J Pharm Biomed Anal.* 1998; 18: 585 -596. [\[Medline\]](#)

Koutsarova N, Todorov P, Koutsarov G. Effect of pentoxifylline on motility and longevity of fresh and thawed dog spermatozoa. *J Reprod Fertil Suppl.* 1997; 51: 117 -121. [\[Medline\]](#)

Lewis SEM, Moohan JM, Thompson W. Effects of pentoxifylline on human sperm motility in normospermic individuals using computer-assisted analysis. *Fertil Steril.* 1993; 59: 418 -423. [\[Medline\]](#)

Lewis SEM, McKinney KA, Thompson W. Influence of pentoxifylline on human sperm motility in asthenozoospermic individuals using computer-assisted analysis. *Arch Androl.* 1994; 32: 175 -183. [\[Medline\]](#)

Loughlin KR, Agarwal A. The use of theophylline to enhance sperm function. *Arch Androl.* 1992; 28: 99 -103. [\[Medline\]](#)

Mahony MC, Lazendorf S, Gordon K, Hodgen GD. Effects of caffeine and dbcAMP on zona pellucida penetration by epididymal spermatozoa of cynomolgus monkeys (*Macaca fascicularis*). *Mol Reprod Dev.* 1996; 43: 530 -535. [\[Medline\]](#)

Mathieu C, Ecochard R, Lorange J, Cordonier H, Guerin JF. Variability of the response to pentoxifylline in vitro in infertile normozoospermic and asthenozoospermic patients. *Arch Androl.* 1994; 33: 39 -49. [\[Medline\]](#)

McKinney KA, Lewis SEM, Thompson W. The effects of pentoxifylline on the generation of reactive oxygen species and lipid peroxidation in human spermatozoa. *Andrologia.* 1996; 28: 15 -20.

Meizel S. Amino acid neurotransmitter receptor/chloride channels of mammalian sperm and the acrosome reaction. *Biol Reprod.* 1997; 56: 569 -574. [\[Abstract\]](#)

Merino G, Chequer JCM, Barahona E, Bermudez JA, Carranza-Lira S. Effects of pentoxifylline on sperm motility in normogonadotropic asthenozoospermic men. *Arch Androl.* 1997; 39: 65 -69. [\[Medline\]](#)

Minelli A, Miscetti P, Proietti A, Luizi L, Mezzasoma I. Adenosine triphosphate catabolism in bovine spermatozoa. *Comp Biochem Physiol B.* 1995; 110: 605 -611. [\[Medline\]](#)

Moniem KA, Glover TD. Alkaline phosphatase in the cytoplasmic droplet of mammalian spermatozoa. *J Reprod Fertil.* 1972; 29: 65 -69.

Monks NJ, Fraser LR. Phosphodiesterase activity of mouse sperm incubated under conditions that modulate fertilizing potential in vitro. *Gamete Res.* 1987; 18: 85 -97. [\[Medline\]](#)

Nagai T, Taneka A, Mori T, Hirayama M. Effects of caffeine and casein phosphopeptides on fertilization in vitro of pig oocytes matured in culture. *Mol Reprod Dev.* 1994; 37: 452 -456. [\[Medline\]](#)

Nassar A, Morshedi M, Mahony M, Srisombut C, Lin MH, Oehninger S. Pentoxifylline stimulates various sperm motion parameters and cervical mucus penetrability in patients with asthenozoospermia. *Andrologia.* 1999a; 31: 9 -15. [\[Medline\]](#)

Nassar A, Mahony M, Morshedi M, Lin MH, Srisombut C, Oehninger S. Modulation of sperm tail protein

tyrosine phosphorylation by pentoxifylline and its correlation with hyperactivated motility. *Fertil Steril*. 1999b;71: 919 -923. [\[Medline\]](#)

Negri P, Grechi E, Tomasi A, Fabbri E, Capuzzo A. Effectiveness of pentoxifylline in semen preparation for intrauterine insemination. *Hum Reprod*. 1996; 11:1236 -1239. [\[Abstract/Free Full Text\]](#)

Okada H, Tatsumi N, Kanzaki M, Fujisawa M, Arakawa S, Kamidono S. Formation of reactive oxygen species by spermatozoa from asthenospermic patients: response to treatment with pentoxifylline. *J Urol*. 1997;157:2140 -2146. [\[Medline\]](#)

Parks JE, Arion JW, Foote RH. Lipids of plasma membrane and outer acrosomal membrane from bovine spermatozoa. *Biol Reprod*. 1987;37:1249 -1258. [\[Abstract\]](#)

Paul M, Sumpter JP, Lindsay KS. Factors affecting pentoxifylline stimulation of sperm kinematics in suspensions. *Hum Reprod*. 1996;11:1929 -1935. [\[Abstract/Free Full Text\]](#)

Ponce AA, de Cuneo MF, Ruiz RD, Vincenti LM, Santilli ME, Stutz G, Lacuara JL. Influence of pentoxifylline on sperm membrane functional integrity. *Arch Androl*. 1999; 43:77 -84. [\[Medline\]](#)

Rees JM, Ford WCL, Hull MGR. Effect of caffeine and of pentoxifylline on the motility and metabolism of human spermatozoa. *J Reprod Fertil*. 1990; 90:147 -156.

Rezende LA, Ciancaglini P, Pizauro JM, Leonoe FA. Inorganic pyrophosphate-phosphohydrolytic activity associated with rat osseous plate alkaline phosphatase. *Cell Mol Biol*. 1998; 44:293 -302.

Sato Y, Tadayoshi M, Suzuki Y. Interaction of pentoxifylline with human erythrocytes. III. Comparison of fluidity changes of erythrocyte membrane caused by S-adenosyl-L-methionine with that by pentoxifylline. *Chem Pharm Bull (Tokyo)*. 1991; 39:468 -473.

Schoff PK, Lardy HA. Effects of fluoride and caffeine on metabolism and motility of ejaculated bovine spermatozoa. *Biol Reprod*. 1987;37:1037 -1046. [\[Abstract\]](#)

Sharma RK, Agarwal A. Influence of artificial stimulation on unprocessed and Percoll-washed cryopreserved sperm. *Arch Androl*. 1997;38:173 -179. [\[Medline\]](#)

Sikka SC, Hellstrom WJG. The application of pentoxifylline in stimulation of sperm motion in man undergoing electroejaculation. *J Androl*. 1991;12:165 -170. [\[Abstract/Free Full Text\]](#)

Soucek DA, Vary JC. Some properties of acid and alkaline phosphatases from boar sperm plasma membranes. *Biol Reprod*. 1984;31:687 -693. [\[Abstract\]](#)

Stachecki JJ, Kinsburg KA, Armant DR. Stimulation of cryopreserved epididymal spermatozoa of the domestic cat using the motility stimulants caffeine, pentoxifylline, and 2'-deoxyadenosine. *J Androl*. 1994;15:157 -164. [\[Abstract/Free Full Text\]](#)

Strzezek J, Glogowski J. Molecular forms of alkaline phosphatase in bull seminal plasma. I. Isolation and characterization of two forms. *Int J Biochem*. 1979; 10:135 -146. [\[Medline\]](#)

Tang Y. Galactosyltransferase, pyrophosphatase and phosphatase activities in luminal plasma of the cauda epididymidis and in the rete testis fluid of some mammals. *J Reprod Fertil*. 1998; 114:277 -285.

Tash JS. Investigations on adenosine 3',5'-monophosphate phosphodiesterase in ram semen and initial characterization of a sperm-specific isoenzyme. *J Reprod Fertil*. 1976;47:63 -72.

Tash JS, Means AR. Cyclic adenosine 3',5' monophosphate, calcium and protein phosphorylation in flagellar motility. *Biol Reprod*. 1983; 28:75 -104. [\[Abstract\]](#)

Tesarik J, Mendoza C, Carreras A. Sperm treatment with pentoxifylline improves fertilizing ability in patients with acrosome reaction insufficiency. *Fertil Steril*.1993; 60:141 -146. [\[Medline\]](#)

Tesarik J, Mendoza C, Carreras A. Effects of phosphodiesterase inhibitors caffeine and pentoxifylline on spontaneous and stimulus-induced acrosome reaction in human sperm. *Fertil Steril*.1992; 58:1185 -1190. [\[Medline\]](#)

Tournaye H, Janssens R, Verheyen G, Devroey P, Van Steirteghem A. In vitro fertilization in couples with previous fertilization failure using sperm incubated with pentoxifylline and 2-deoxyadenosine. *Fertil Steril*. 1994;62:574 -579. [\[Medline\]](#)

Vandevoort CA, Tollner TL, Overstreet JW. Separate effects of caffeine and dbcAMP on macaque sperm motility and interaction with the zona pellucida. *Mol Reprod Dev*.1994; 37:299 -304. [\[Medline\]](#)

Vijayaraghavan S, Hoskins DD. Regulation of bovine sperm motility and cyclic adenosine 3',5'-monophosphate by adenosine and its analogs. *Biol Reprod*.1986; 34:468 -477. [\[Abstract\]](#)

Vinet B, Zizan L, Gauthier B. Characteristics of the inhibition of serum alkaline phosphatase by theophylline. *Clin Biochem*. 1978;11:57 -61. [\[Medline\]](#)

Wang H, Gilles-Baillien M. Alkaline phosphatase and ATPases in brush-border membranes of rat jejunum: distinct effect of divalent cations and of some inhibitors. *Arch Int Physiol Biochim Biophys*.1992; 100:289 -294. [\[Medline\]](#)

Wang R, Sikka SC, Veeragaven K, Bell M, Hellstrom WJG. Platelet activating factor and pentoxifylline as human sperm cryoprotectant. *Fertil Steril*.1993; 60:711 -715. [\[Medline\]](#)

Wells JN, Garbers DL. Nucleoside 3',5'-monophosphate phosphodiesterases in sea urchin sperm. *Biol Reprod*.1976; 15:46 -53. [\[Abstract\]](#)

Yuan YY, Shi QX, Srivastava PN. Inhibition of rabbit sperm acrosomal enzymes by gossypol. *Mol Reprod Dev*.1995; 40:228 -232. [\[Medline\]](#)

#### This Article

- ▶ [Abstract](#) **FREE**
- ▶ [Full Text \(PDF\)](#)
- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me if a correction is posted](#)

#### Services

- ▶ [Similar articles in this journal](#)
- ▶ [Similar articles in PubMed](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Download to citation manager](#)

#### Citing Articles

- ▶ [Citing Articles via Google Scholar](#)

#### Google Scholar

- ▶ [Articles by Glogowski, J.](#)
- ▶ [Articles by Ciereszko, A.](#)
- ▶ [Search for Related Content](#)

#### PubMed



▶ [PubMed Citation](#)

▶ [Articles by Glogowski, J.](#)

▶ [Articles by Cierieszko, A.](#)

---

[HOME](#) [HELP](#) [FEEDBACK](#) [SUBSCRIPTIONS](#) [ARCHIVE](#) [SEARCH](#) [TABLE OF CONTENTS](#)