

Published-Ahead-of-Print November 1, 2006, DOI:10.2164/jandrol.106.000729

Journal of Andrology, Vol. 28, No. 2, March/April 2007

Copyright © [American Society of Andrology](#)

DOI: 10.2164/jandrol.106.000729

Trends in Semen Parameters in the Northeast of Scotland

SREEBALA SRIPADA^{*}, SOFIA FONSECA[†], AMANDA LEE[‡], KIRSTEN HARRILD[†],
DIMITRIOS GIANNARIS[§], EILEEN MATHERS^{*} AND SILADITYA BHATTACHARYA^{*}

From the ^{*} Aberdeen Fertility Centre, Aberdeen Maternity Hospital, Aberdeen, Scotland; the [†] Department of Public Health and the [‡] Department of General Practice and Primary Care, University of Aberdeen, Scotland; and the [§] Department of Obstetrics & Gynaecology, University of Athens, Greece.

Correspondence to: Dr Sreebala Sripada, Assisted Reproduction Unit, Aberdeen Maternity Hospital, Foresterhill, Aberdeen AB25 2ZD, United Kingdom (e-mail: ogy302{at}abdn.ac.uk).

Received for publication May 18, 2006; accepted for publication October 30, 2006.

This Article

- ▶ [Abstract](#) **FREE**
- ▶ [Full Text \(PDF\)](#)
- ▶ All Versions of this Article: 28/2/313 *most recent*
[Author Manuscript \(PDF\)](#) **FREE**
- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me if a correction is posted](#)

Services

- ▶ [Similar articles in this journal](#)
- ▶ [Similar articles in PubMed](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Download to citation manager](#)

Citing Articles

- ▶ [Citing Articles via HighWire](#)
- ▶ [Citing Articles via Google Scholar](#)

Google Scholar

- ▶ [Articles by Sripada, S.](#)
- ▶ [Articles by Bhattacharya, S.](#)
- ▶ [Search for Related Content](#)

PubMed

- ▶ [PubMed Citation](#)
- ▶ [Articles by Sripada, S.](#)
- ▶ [Articles by Bhattacharya, S.](#)

Abstract

The data on trends in semen quality are conflicting and sensitive to geographical variations. Although previous British surveys on semen quality indicate a decline, the northeast of Scotland has never been included in these surveys. This is an area with low out migration rates where andrology services for a population of 500 000 are centralized within a single laboratory, thus providing a unique opportunity to study population-based trends in semen quality over time. We investigated trends in semen parameters between 1994–2005, in a cohort of 4832 men attending for routine semen analysis at the Aberdeen Fertility Centre who had a sperm density of greater than 20 million per mL. The main outcome measures were trends in sperm density, sperm motility and motile density in the first semen sample. Linear regression and time series analysis were used to examine trends over time in the semen parameters. The mean and standard deviation (SD) age of all men (n = 5204) in the study was 34 (6) years. The median (inter quartile range) for sperm density and motile density for the study population were 61 (40–91) million/mL and 99 (47–181) million. The mean (SD) sperm motility was 49 (19)%. Among 4832 men (with sperm count >20 million per mL), data adjusted for age and period of abstinence showed a decreasing trend for sperm density over time, $R^2 = 0.45$ ($P = .017$). There was no such trend in sperm motility and motile density. However, this

- ▲ [Top](#)
- [Abstract](#)
- ▼ [Materials and Methods](#)
- ▼ [Results](#)
- ▼ [Discussion](#)
- ▼ [References](#)

trend has to be interpreted with caution due to fluctuations in semen parameters, population bias and the retrospective nature of the analysis.

Key words: Fertility, infertility, sperm

The last two decades have seen the publication of several reports that suggest a global decline in semen quality ([Bostofte et al, 1983](#); [Carlsen et al, 1992](#); [Auger et al, 1995](#); [Irvine et al, 1996](#); [Swan et al, 1997](#); [Bilotta et al, 1999](#)). Within Western Europe, a four-centre study on fertile couples ([Jensen et al, 2001](#)) has shown substantial variations in semen parameters. However, the data on trends in semen quality are conflicting and sensitive to geographical variations ([Bujan et al, 1996](#); [Fisch et al, 1996a](#); [Benshushan et al, 1997](#); [Costello et al, 2002](#); [Pal et al, 2006](#)). It has also been difficult to prove an association between deterioration in semen quality and male infertility. The quality of existing studies has been questioned on the basis of population and selection bias, variations in laboratory standards, and statistical methodologies used for the analysis. Although previous British surveys on semen quality indicate a decline, the northeast of Scotland has never been included in these surveys. Earlier studies from France indicate variations in different cities within that country ([Auger and Jouannet, 1997](#)).

The northeast of Scotland is an area with low outward migration rates ([Hall et al, 1989](#); [Batty et al, 2004](#); [General Register Office of Scotland, 2006](#)), where the andrology services for a population of 500 000 are centralized within a single laboratory. This provides a unique opportunity to study population-based trends in semen quality over a period of time. In the present study, we investigated the trends in semen parameters over time in a cohort of men who attended for routine semen analysis with sperm densities greater than $20 \times 10^6/\text{mL}$.

► **Materials and Methods**

Study Subjects

This was a retrospective, observational study based on data from the Aberdeen Andrology Database. This is a secure database into which data on men who have attended for semen analysis are entered prospectively. The subjects included the male partners of subfertile couples in the Grampian region of Scotland who attended the Aberdeen Fertility Centre between 1994 and 2005. Although initially it appeared that more men attended in the period 2001 to 2005 than during the period 1994 to 1999, further analysis showed no changes in referring practice over this period. Oligozoospermic men, azoospermic men, and men with a history of vasectomy were excluded from the current analyses. The etiologies and severities of infertility were not analyzed. In order to minimize selection bias, the first semen sample from each man was used for the present study (by year of donation), and the values of sperm concentration (density), motility, and motile density were analyzed. Data on sperm morphology were excluded due to significant variations in laboratory techniques and criteria for normality over the study period. During the study period, routine semen analysis was performed using the Hamilton Thorne Sperm Motility Analyzer (Beverly, Mass) according to the guidelines of the World Health Organization (WHO). The National External Quality Assessment Service (NEQAS) oversaw the external quality controls for the laboratory four times per year. No nonconformities were reported. Internal audits for adherence to standard operating procedures during sample preparation and analysis, as well as weekly comparisons between technicians were additional internal quality control measures.

- ▲ [Top](#)
- ▲ [Abstract](#)
- [Materials and Methods](#)
- ▼ [Results](#)
- ▼ [Discussion](#)
- ▼ [References](#)

Semen Analysis


The laboratory instructed the men to abstain from ejaculation for 3–7 days before providing a sample by masturbation into a sterile plastic container. The sample was allowed to liquefy at 37° C for an initial period of 30–60 minutes, and all analyses were performed within 90 minutes of ejaculation. The volume of the ejaculate was determined by aspirating the liquefied sample into a graduated disposable pipette. To determine the concentration of sperm ($10^6/\text{mL}$), and sperm motility, a 10- μL drop of the semen sample was placed on a commercially produced, fixed-depth capillary fill chamber. The chamber used until 2002 was the Conception Technology Microcell MC-20-2 (San Diego, Calif). This was replaced by the Leja Standard Count 20- μm Analysis Chamber (Nieuw-Vennep, The Netherlands) in the subsequent years. When the chamber was full, the slide was placed in a heated stage at 37° C. The stage was placed on the microscope and the sample was assessed using the Hamilton Thorne HTM-S Semen Analyzer until the year 2000, after which the Hamilton Thorne Version 10 HTM-CEROS was used. A minimum of 200 sperm or 2 frames were counted. Local comparisons of changes in chamber and software were performed. These showed no significant differences with respect to sperm densities or motilities. Progressive motility was determined as the proportion of sperm that showed evidence of movement (WHO grades a and b) to the total number of spermatozoa counted (WHO grades a, b, c, and d) (WHO, 1993, [1999](#)).

Statistical Analysis

The software packages SPSS (13.0) (SPSS Inc, Chicago, Ill) and Microsoft Excel 2000 (Redmond, Wash) were used for statistical analysis. Linear regression and time series analyses were used to examine trends over time in the three semen parameters of interest (ie, density, motility, and motile density).

As all our men were partners in a subfertile relationship, we analyzed the trend in semen parameters for men with a sperm density greater than $20 \times 10^6/\text{mL}$, to minimize selection bias. In the time trends analysis, only men with a period of abstinence of 1–10 days and who were aged 22–56 years were included, since the consistent linear relationships between mean sperm parameters and these two confounders were no longer evident beyond these limits ([Figure 1a and b](#)). Initially, analysis of variance (ANOVA) and Kruskal-Wallis tests were carried out to see if there were any overall differences in sperm density, motility or motile density, for each year of the period 1994 to 2005 (Table 1). Examination of the mean values for the sperm parameters of these men for each year of donation from 1994 to 2005 showed that these values were normally distributed. Thus, log transformation of these data, as previously suggested ([Berman et al, 1996](#)), was not required. A linear trend was fitted to the mean of each variable of interest (sperm density, sperm motility, and motile density) across the period of abstinence from 1 to 10 days and across age from 22 to 56 years. The residuals of the fitted linear models were then utilized as the age-adjusted and abstinence-adjusted data for the follow-up period.

Figure 1. (a) Relationship between sperm parameters and period of abstinence. (b) Relationship between sperm parameters and age.



View larger version
(23K):

[\[in this window\]](#)
[\[in a new window\]](#)

Results

- ▲ [Top](#)
- ▲ [Abstract](#)
- ▲ [Materials and Methods](#)
- [Results](#)
- ▼ [Discussion](#)
- ▼ [References](#)

Data on semen samples from a total of 6761 men were available for analysis ([Figure 2](#)). Of the 6761 men, 5204 men were normal in terms of sperm density (ie, had densities greater than $20 \times 10^6/\text{mL}$). The raw data on age and sperm motility were normally distributed, while sperm density and motile density were positively skewed. The mean \pm SD age of the men ($n = 5204$) in the present study was 34 ± 6 years. The median (interquartile range) values for sperm density and motile density for the study population were 61 (40–91) $\times 10^6/\text{mL}$ and 99 (47–181) $\times 10^6/\text{mL}$. The mean \pm SD for sperm motility was $49 \pm 19\%$.




Figure 2. Characteristics of the men included in the analysis.

View larger version (9K):

[\[in this window\]](#)
[\[in a new window\]](#)

Since length of abstinence showed strong linear relationships with each of the three sperm parameters ([Figure 1a](#)), for the analysis of trend over time we considered the data for men whose period of abstinence was between 1 and 10 days. As shown in [Figure 1B](#), men younger than 22 years and those older than 56 years had large fluctuations in sperm parameters, and since they constituted

only a small proportion of the population studied, these men were excluded from any subsequent analysis. Of the 5204 men who had a sperm count of more than $20 \times 10^6/\text{mL}$, 103 men had incomplete data for age and abstinence, 219 men did not abstain at all or abstained for more than 10 days and 50 men were either younger than 22 years or older than 56 years ([Figure 2](#)). These subjects were excluded from the current study, which left 4832 men for analysis. Descriptive statistics for the unadjusted data for each of the sperm parameters for each year from 1994 to 2005 are listed in the Table, along with the results of the ANOVA and Kruskal-Wallis tests. There were significant differences in the averaged annual sperm density and motile density values across the 12-year period. [↗](#)

View this table:
[\[in this window\]](#)
[\[in a new window\]](#)

*Semen parameters for 4832 men with normal sperm densities**

For the linear regression analysis, the data were adjusted for age and abstinence by first considering the residuals of the fitted linear model, which included abstinence, and second, utilizing these data for fitting a subsequent linear model for age. The residuals of the latter fitted models were then considered as being adjusted for age and abstinence. [Figure 3](#) illustrates the variations in mean sperm density, motility, and motile density in 4832 men with sperm count of more than $20 \times 10^6/\text{mL}$, across the 12-year period between 1994 and 2005. The unadjusted and adjusted means for each year for each variable were calculated. As shown in [Figure 3](#), decreasing trends over time were noted for sperm density and motile density before the data were adjusted for age and abstinence. The adjusted data show decreased sperm density ($R^2 = 0.45$, $P = .017$), while there was no evidence for similar trends in sperm motility and motile density. There was no significant difference in the mean period of abstinence over this time period.




Figure 3. Variations over time in sperm parameters in normozoospermic men. **(A)** Unadjusted data (data before adjusting for age and abstinence). **(B)** Adjusted data (data after adjusting for age and abstinence).

View larger version
(32K):
[\[in this window\]](#)
[\[in a new window\]](#)

▲ Top
▲ Abstract
▲ Materials and Methods
▲ Results
▪ Discussion
▼ References

Our results suggest a decline in sperm counts over a 12-year period among men with a sperm density greater than $20 \times 10^6/\text{mL}$ in the northeast of Scotland. Similar trends in sperm motility and motile density values were not observed.

The relatively stable population, large sample size, and the use of a single laboratory to process all the samples are the main strengths of the present study. The availability of data regarding age and abstinence allowed adjustment for these important confounders. One of the main weaknesses of the present study is that all the subjects were men from subfertile partnerships. However, to minimize selection bias, we analyzed the trends only among those men who had a normal sperm density.

Studies on semen parameters should acknowledge the large variation in samples from the same individual and between individuals ([Heuchel et al, 1983](#); [Gyllenborg et al, 1999](#)). In order to minimize intraindividual bias, we used the first semen sample from each man. Circannual variation is a controversial topic ([Tjoa et al, 1982](#); [Carlsen et al, 2004](#); [Malm et al, 2004](#)). However, as our data were collected over all the 12 months in each year, the impact of seasonal variation was minimized by using the mean annual values. The reasons for fluctuations in semen parameters for men of less than 22 years of age and more than 56 years of age ([Figure 1b](#)) are unclear, although these fluctuations may be due to the lower numbers of men in these age groups.

While most studies ([Auger et al, 1995](#); [Irvine et al, 1996](#)) have used the conventional method for semen analysis, in the present study, the samples were analyzed using the Hamilton Thorne Method, which gives higher values for sperm density ([Neuwinger et al, 1990](#)) and lower values for sperm motility ([Sukcharoen and Aribarg, 1995](#)). Therefore, methodological differences do not explain the decreasing trend seen in the present study, as the use of computer-assisted semen analysis has the opposite effect.

This study has the largest sample size of any reported from the United Kingdom. While previous studies have used linear regression analyses, some have taken no account of confounders, such as age and abstinence. Although we considered using multiple linear regression, examination of the residuals from these models indicated high correlation levels, which violated the assumption of independent residuals. Therefore, we used the time series approach. Our study was population-based and adjusted for the confounders of age and abstinence. Our results are in line with those of an earlier study from the United Kingdom ([Irvine et al, 1996](#)). Geographical factors may well be responsible for the conflicting results reported in many other studies ([Bujan et al, 1996](#); [Fisch et al, 1996a](#); [Benshushan et al, 1997](#); [Costello et al, 2002](#); [Pal et al, 2006](#)).

In the present study, the sperm densities for subfertile men were lower than those reported in Danish ([Rasmussen et al, 1997](#)), French ([Auger et al, 1995](#)), and Finnish ([Vierula et al, 1996](#)) studies but were similar to the mean sperm concentrations in the preliminary unpublished results of the CHAPS-UK study. This suggests that British men have a lower average sperm concentration than men in most parts of Europe. There is a wide geographical variation in the mean levels of semen parameters. A Danish study ([Rasmussen et al, 1997](#)) from a similar subfertile population during the same time interval reported a high mean sperm density of $183.7 \times 10^6/\text{mL}$, while the mean sperm concentration of subfertile men in Korea was low at $60.5 \times 10^6/\text{mL}$. It is of note that the mean sperm density for all 1055 men from subfertile partnerships in the Danish study ($183.7 \times 10^6/\text{mL}$) was well above the maximum sperm density ($98 \times 10^6/\text{mL}$) in the Scottish study ([Irvine et al, 1996](#)) of fertile men who were semen donors for the Gamete Biology Research study.

Fisch et al (1996b) have highlighted the variation in mean sperm concentrations in fertile men in different countries. Within the United States, the mean concentrations of sperm in fertile populations were as high as $134 \times 10^6/\text{mL}$ in New York and as low as $48 \times 10^6/\text{mL}$ in Iowa (Fisch et al, 1996b). In addition, the reported mean sperm concentrations in developing countries have been reported as being low (Aribarg et al, 1986; Osegbe et al, 1986).

Various explanations have been put forward for the observed decline in sperm counts. General lifestyle factors, such as smoking (Storgaard et al, 2003), alcohol, drug use, and obesity may have etiological roles. In Scotland (Boyle et al, 1987), as in many other parts of the world (Bergstrom et al, 1996; dos Santos Silva et al, 1999), the incidence of testicular cancer, particularly in the age group of 15–44 years, has increased by about 50%. There have also been increases in other male genital abnormalities, such as cryptorchidism and hypospadias, in many populations. This highlights the probable role of intrauterine etiological mechanisms. Endocrine-disrupting chemicals may play a causal role (Murray et al, 2001). Exposure to estrogens during fetal life has been implicated (Storgaard et al, 2006). However, the much higher estrogen levels noted in twin pregnancies, despite these being more potent forms of estrogen, do not appear to result in lower sperm counts in adulthood (Storgaard et al, 2002). Some of the studies reporting declines in sperm counts have also noticed qualitative differences in semen parameters. Whether this is due to infections, such as with *Chlamydia epididymoorchitis*, remains to be investigated (Eley et al, 2005). The current literature highlights the increasing prevalence of genital infections among men and women (LaMontagne et al, 2004). Many factors may be operational, and it was not possible to assess the effects of causative factors in the present study.

Although there is some suggestion that sperm concentrations may be decreasing in some parts of the world, there is no conclusive evidence of decreased fertility potential in the human male population. It may be that semen parameters are poor predictive indicators of fertility potential. Alternatively, undetected compensatory mechanisms may be functioning.

We have described how Scottish men who attended a fertility clinic show a decline in semen density over a 12-year period. However, this trend has to be interpreted with caution due to fluctuations in semen parameters, population bias, and the retrospective nature of the analysis. More prospective studies are required to investigate this phenomenon and to investigate the roles of contributory factors.

References

Aribarg A, Kenkeerati W, Vorapaiboonsak V, Leepipatpai boon S, Farley TM. Testicular volume, semen profile and serum hormone levels in fertile Thai males. *Int J Androl*. 1986; 9: 170 – 180. [\[Medline\]](#)

Auger J, Jouannet P. Evidence for regional differences of semen quality among fertile French men. Federation Francaise des Centres d'Etude et de Conservation des Oeufs et du Sperme humains. *Hum Reprod*. 1997;12: 740 – 745. [\[Abstract/Free Full Text\]](#)

Auger J, Kunstmann JM, Czyglik F, Jouannet P. Decline in semen quality among fertile men in Paris during the past 20 years. *N Engl J Med*. 1995;332: 281 – 285. [\[Abstract/Free Full Text\]](#)

Batty GD, Morton SM, Campbell D, Clark H, Smith GD, Hall M, Macintyre S, Leon DA. The Aberdeen Children of the 1950s cohort study: background, methods and follow-up information on a new resource for the study of life course and intergenerational influences on health. *Paediatr Perinat Epidemiol*.

- [▲ Top](#)
- [▲ Abstract](#)
- [▲ Materials and Methods](#)
- [▲ Results](#)
- [▲ Discussion](#)
- [References](#)

Benshushan A, Shoshani O, Paltiel O, Schenker JG, Lewin A. Is there really a decrease in sperm parameters among healthy young men? A survey of sperm donations during 15 years. *J Assist Reprod Genet.* 1997;14: 347 – 353. [\[Medline\]](#)

Bergstrom R, Adami HO, Mohner M, Zatonski W, Storm H, Ekblom A, Tretli S, Teppo L, Akre O, Hakulinen T. Increase in testicular cancer incidence in six European countries: a birth cohort phenomenon. *J Natl Cancer Inst.* 1996;88: 727 – 733. [\[Abstract/Free Full Text\]](#)

Berman NG, Wang C, Paulsen CA. Methodological issues in the analysis of human sperm concentration data. *J Androl.* 1996; 17: 68 – 73. [\[Abstract/Free Full Text\]](#)

Bilotta P, Guglielmo R, Steffe M. Analysis of decline in seminal fluid in the Italian population during the past 15 years. *Minerva Ginecol.* 1999;51: 223 – 231. [\[Medline\]](#)

Bostofte E, Serup J, Rebbe H. Has the fertility of Danish men declined through the years in terms of semen quality? A comparison of semen qualities between 1952 and 1972. *Intl J Fertil.* 1983; 28: 91 – 95.

Boyle P, Kaye SB, Robertson AG. Changes in testicular cancer in Scotland. *Eur J Cancer Clin Oncol.* 1987; 23: 827 – 830. [\[CrossRef\]](#)[\[Medline\]](#)

Bujan L, Mansat A, Pontonnier F, Mieusset R. Time series analysis of sperm concentration in fertile men in Toulouse, France between 1977 and 1992. *BMJ.* 1996; 312: 471 – 472. [\[Abstract/Free Full Text\]](#)

Carlsen E, Giwercman A, Keiding N, Skakkebaek NE. Evidence for decreasing quality of semen during past 50 years. *BMJ.* 1992; 305: 609 – 613. [\[Medline\]](#)

Carlsen E, Petersen JH, Andersson AM, Skakkebaek NE. Effects of ejaculatory frequency and season on variations in semen quality. *Fertil Steril.* 2004; 82: 358 – 366. [\[CrossRef\]](#)[\[Medline\]](#)

Costello MF, Sjoblom P, Haddad Y, Steigrad SJ, Bosch EG. No decline in semen quality among potential sperm donors in Sydney, Australia, between 1983 and 2001. *J Assist Reprod Genet.* 2002; 19: 284 – 290. [\[CrossRef\]](#)[\[Medline\]](#)

dos Santos Silva I, Swerdlow AJ, Stiller CA, Reid A. Incidence of testicular germ-cell malignancies in England and Wales: trends in children compared with adults. *Int J Cancer.* 1999; 83: 630 – 634. [\[CrossRef\]](#)[\[Medline\]](#)

Eley A, Pacey AA, Galdiero M, Galdiero M, Galdiero F. Can *Chlamydia trachomatis* directly damage your sperm? *Lancet Infect Dis.* 2005;5: 53 – 57. [\[CrossRef\]](#)[\[Medline\]](#)

Fisch H, Goluboff ET, Olson JH, Feldshuh J, Broder SJ, Barad DH. Semen analyses in 1,283 men from the United States over a 25-year period: no decline in quality. *Fertil Steril.* 1996a; 65: 1009 – 1014. [\[Medline\]](#)

Fisch H, Ikeguchi EF, Goluboff ET. Worldwide variations in sperm counts. *Urology.* 1996b; 48: 909 – 911. [\[CrossRef\]](#)[\[Medline\]](#)

General Register Office for Scotland. Comparative population profile: Aberdeenshire council area Scotland. 2006. Available at: <http://www.scrol.gov.uk/scrol/browser/profile.jsp?profile=Population&mainArea=Aberdeenshire&mainLevel=CouncilArea>. Accessed 28 December 2006.

Gyllenberg J, Skakkebaek NE, Nielsen NC, Keiding N, Giwercman A. Secular and seasonal changes in semen quality among young Danish men: a statistical analysis of semen samples from 1927 donor candidates during 1977-1995. *Int J Androl.* 1999; 22: 28 – 36. [\[CrossRef\]](#)[\[Medline\]](#)

Hall MH, Campbell DM, Fraser C, Lemon J. Mode of delivery and future fertility. *Br J Obstet Gynaecol.* 1989; 96: 1297 – 1303. [\[Medline\]](#)

Heuchel V, Schwartz D, Czyglik F. Between and within subject correlations and variances for certain semen characteristics in fertile men. *Andrologia.* 1983; 15: 171 – 176. [\[Medline\]](#)

Irvine S, Cawood E, Richardson D, MacDonald E, Aitken J. Evidence of deteriorating semen quality in the United Kingdom: birth cohort study in 577 men in Scotland over 11 years. *BMJ.* 1996; 312: 467 – 471. [\[Abstract/Free Full Text\]](#)

Jensen TK, Slama R, Ducot B, Suominen J, Cawood EH, Andersen AG, Eustache F, Irvine S, Auger S, Jouannet P, Vierula M, Jorgensen N, Toppari J, Skakkebaek NE, Keiding N, Spira A. Regional differences in waiting time to pregnancy among fertile couples from four European cities. *Hum Reprod.* 2001;16: 2697 – 2704. [\[Abstract/Free Full Text\]](#)

LaMontagne DS, Fenton KA, Randall S, Anderson S, Carter P. Establishing the National Chlamydia Screening Programme in England: results from the first full year of screening. *Sex Transm Infect.* 2004;80: 335 – 341. [\[Abstract/Free Full Text\]](#)

Malm G, Haugen TB, Henriksen T, Bjorsvik C, Grotmol T, Saether T, Malm J, Figenschau Y, Hagmar L, Rylander L, Levine RJ, Giwercman A. Reproductive function during summer and winter in Norwegian men living north and south of the Arctic circle. *J Clin Endocrinol Metab.* 2004;89: 4397 – 4402. [\[Abstract/Free Full Text\]](#)

Murray TJ, Lea RG, Abramovich DR, Haites NE, Fowler PA. Endocrine disrupting chemicals: effects on human male reproductive health. *Early Pregnancy [Electronic Resource]*. 2001; 5: 80 – 112.

Neuwinger J, Behre HM, Nieschlag E. External quality control in the andrology laboratory: an experimental multicenter trial. *Fertil Steril.* 1990;54: 308 – 314. [\[Medline\]](#)

Osegbe DN, Amaku EO, Nnatu SN. Are changing semen parameters a universal phenomenon? *European Urology.* 1986; 12: 164 – 168. [\[Medline\]](#)

Pal PC, Rajalakshmi M, Manocha M, Sharma RS, Mittal S, Rao DN. Semen quality and sperm functional parameters in fertile Indian men. *Andrologia.* 2006; 38: 20 – 25. [\[CrossRef\]](#) [\[Medline\]](#)

Rasmussen PE, Erb K, Westergaard LG, Laursen SB. No evidence for decreasing semen quality in four birth cohorts of 1,055 Danish men born between 1950 and 1970. *Fertil Steril.* 1997; 68: 1059 – 1064. [\[CrossRef\]](#) [\[Medline\]](#)

Storgaard L, Bonde JP, Ernst E, Andersen CY, Kyvik KO, Olsen J. Effect of prenatal exposure to oestrogen on quality of semen: comparison of twins and singleton brothers. *BMJ.* 2002; 325: 252 – 253. [\[Free Full Text\]](#)

Storgaard L, Bonde JP, Ernst E, Spano M, Andersen CY, Frydenberg M, Olsen J. Does smoking during pregnancy affect sons' sperm counts? *Epidemiology.* 2003; 14: 278 – 286. [\[CrossRef\]](#) [\[Medline\]](#)

Storgaard L, Bonde JP, Olsen J. Male reproductive disorders in humans and prenatal indicators of estrogen exposure. A review of published epidemiological studies. *Reprod Toxicol.* 2006; 21: 4 – 15. [\[CrossRef\]](#) [\[Medline\]](#)

Sukcharoen N, Aribarg A. Evaluation of the Hamilton-Thorn Motility Analyzer for routine semen analysis. *J Med Assoc Thai.* 1995; 78: 182 – 190. [\[Medline\]](#)

Swan SH, Elkin EP, Fenster L. Have sperm densities declined? A reanalysis of global trend data. *Environ Health Perspect.* 1997;105: 1228 – 1232. [\[Medline\]](#)

Tjoa WS, Smolensky MH, Hsi BP, Steinberger E, Smith KD. Circannual rhythm in human sperm count revealed by serially independent sampling. *Fertil Steril*. 1982; 38: 454 – 459. [\[Medline\]](#)

Vierula M, Niemi M, Keiski A, Saaranen M, Saarikoski S, Suominen J. High and unchanged sperm counts of Finnish men. *Int J Androl*. 1996;19: 11 – 17. [\[Medline\]](#)

World Health Organization. *WHO Laboratory Manual for the Examination of Human Semen and Sperm-Cervical Mucus Interactions*. 4th ed. Cambridge, United Kingdom: Cambridge University Press; 1999 .

This article has been cited by other articles:



HUMAN REPRODUCTION UPDATE

▶ HOME

A. Nyboe Andersen, E. Carlsen, and A. Loft

Trends in the use of intracytoplasmic sperm injection marked variability between countries

Hum. Reprod. Update, November 1, 2008; 14(6): 593 - 604.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

This Article

- ▶ [Abstract](#) **FREE**
- ▶ [Full Text \(PDF\)](#)
- ▶ All Versions of this Article:
28/2/313 *most recent*
[Author Manuscript \(PDF\)](#) **FREE**
- ▶ [Alert me when this article is cited](#)
- ▶ [Alert me if a correction is posted](#)

Services

- ▶ [Similar articles in this journal](#)
- ▶ [Similar articles in PubMed](#)
- ▶ [Alert me to new issues of the journal](#)
- ▶ [Download to citation manager](#)

Citing Articles

- ▶ [Citing Articles via HighWire](#)
- ▶ [Citing Articles via Google Scholar](#)

Google Scholar

- ▶ [Articles by Sripada, S.](#)
- ▶ [Articles by Bhattacharya, S.](#)
- ▶ [Search for Related Content](#)

PubMed

- ▶ [PubMed Citation](#)
- ▶ [Articles by Sripada, S.](#)
- ▶ [Articles by Bhattacharya, S.](#)