



HOME HELP FEEDBACK SUBSCRIPTIONS ARCHIVE SEARCH TABLE OF CONTENT

Journal of Andrology, Vol 4, Issue 2 131-135, Copyright © 1983 by The American Society of Andrology

JOURNAL ARTICLE

# Effect of cryptorchidism and orchidopexy on inhibin secretion by rat Sertoli cells

L. Seethalakshmi and A. Steinberger

The present study explored the effects of experimental bilateral cryptorchidism (of 21-, 28-, and 35-days duration) and orchidopexy (14 and 42 days) in the adult rat on the secretion of inhibin by cultures of isolated Sertoli cells. Changes in serum levels of gonadotropins, testis weight, and spermatogenesis also were assessed to verify the effectiveness of the surgical procedures. Cryptorchidy resulted in a progressive decline in testicular weight and a loss of germ cells, associated with increasing serum levels of FSH and LH.

#### This Article

- ▶ Full Text (PDF)
- Alert me when this article is cited
- Alert me if a correction is posted

#### Services

- ▶ Similar articles in this journal
- ▶ Similar articles in PubMed
- ▶ Alert me to new issues of the journal
- ▶ <u>Download to citation manager</u>

## Citing Articles

Liting Articles via Google Scholar

## Google Scholar

- Articles by Seethalakshmi, L.
- Articles by Steinberger, A.
- Search for Related Content

### PubMed

- ▶ PubMed Citation
- Articles by Seethalakshmi, L.
- Articles by Steinberger, A.

Inhibin secretion in vitro became nondetectable by 28 days after surgery. At 42 days after orchidopexy, spermatogenesis showed qualitative recovery, with a small increase in testes weight. Levels of LH in the circulation declined, but only to twice the intact control levels. However, inhibin secretion and serum FSH levels returned to nearly normal values. These results indicate that bilateral cryptorchidism severely impairs the secretion of inhibin and possibly other Sertoli cell functions which may account, at least partly, for the increase in circulating FSH levels and the arrest of spermatogenesis. The effects of cryptorchidism on these parameters can be reversed to a large degree by orchidopexy.

HOME HELP FEEDBACK SUBSCRIPTIONS ARCHIVE SEARCH TABLE OF CONTENTS

Copyright © 1983 by The American Society of Andrology.