



 **Current Issue**

 **Browse Issues**

 **Search**



 **About this Journal**

 **Instruction to Authors**

 **Online Submission**

 **Subscription**

 **Contact Us**



 **RSS Feed**

## Acta Medica Iranica

2009;47(4) : 24-25

"Tc-99m MIBI imaging in lymphomas: Comparison with Tl-201 and Ga-67 scintigraphy "

"Dadparvar S, Damsker JL, Asrari F, Dou Y, Patil S, Brodsky I "

### Abstract:

Tc-99m MIBI has recently been used in the functional imaging of various tumors. This prospective study was performed to evaluate the role of Tc-99m MIBI imaging at the time of the initial staging, assessment of treatment response, follow-up studies and surveillance in Hodgkin's and non-Hodgkin's Lymphoma. 25 patients (14 with Hodgkin's and 11 with non-Hodgkin's Lymphoma) underwent 32 studies. The patient's age range was 19-73 years, with a mean age of 41.7 years. All patients underwent whole body planar imaging following intravenous injection of 20-25 mCi Tc-99m MIBI. Comparative imaging with 3 mCi Tl-201 and 10 mCi Ga-67 injection was done. Clinical, radiographic and biopsy correlation was obtained in all cases.

TUMS ID: 1330

Full Text HTML  Full Text PDF  425 kB

top ▲

[Home](#) - [About](#) - [Contact Us](#)

TUMS E. Journals 2004-2009  
Central Library & Documents Center  
Tehran University of Medical Sciences

Best view with Internet Explorer 6 or Later at 1024\*768 Resolutions