



 **Current Issue**

 **Browse Issues**

 **Search**



 **About this Journal**

 **Instruction to Authors**

 **Online Submission**

 **Subscription**

 **Contact Us**



 **RSS Feed**

Acta Medica Iranica

2009;47(4) : 21-27

Ecology of Anopheles dthali Patton in Bandar Abbas District, Hormozgan Province, Southern Iran

H Vatandoost, M Shahi, AA Hanafi-Bojd, MR Abai, MA Oshaghi, F Raffi

Abstract:

Ecology of Anopheles dthali was studied in Bandar Abbas County, where there is indigenous malaria. Anopheles dthali plays as a secondary malaria vector in the region. It is active throughout the year in mountainous area with two peaks of activity, whereas in coastal area it has one peak. There is no report of hibernation or aestivation for this species in the region. Precipitin tests on specimens from different parts showed that 15.6-20.8% were positive for human blood. This species usually rests outdoors. It has different larval habitats. Insecticides susceptibility tests on adult females exhibited susceptibility to all insecticides recommended by WHO. LT50 for the currently used insecticide, lambda-cyhalothrin, is measured less than one minute. The irritability tests to pyrethroid insecticides, showed that permethrin and lambda-cyhalothrin had more irritancy compared to deltamethrin and cyfluthrin. Larval bioassay using malathion, chlorpyrifos, temephos and fenithrothion did not show any sign of resistance to these larvicides at the diagnostic dose. It is recommended that all the decision makers should consider the results of our study for any vector control measures in the region.

Keywords:

Malaria vector

TUMS ID: 3589

Full Text HTML  Full Text PDF  142 KB

top ▲

[Home](#) - [About](#) - [Contact Us](#)

TUMS E. Journals 2004-2009
Central Library & Documents Center
Tehran University of Medical Sciences

Best view with Internet Explorer 6 or Later at 1024*768 Resolutions