

Received: July 5, 2001

Accepted: September 15, 2001

Ref: Eke F.U., Frank-Briggs A., Ottor J. Childhood Mortality In Port Harcourt, Nigeria. *Anil Aggrawal's Internet Journal of Forensic Medicine and Toxicology*, 2001; Vol. 2, No. 2 (July-Dec 2001): ; Published September 15, 2001, ([Accessed:](#)

[Email Felicia Eke by clicking here](#)



EMBASE Accession Number: 2004204927



Felicia Eke
(Click to enlarge
in a different
window)

Childhood Mortality in Port Harcourt, Nigeria

-By:

*Professor Felicia U Eke , FRCP,
FRCPCH, MD, FWACP*

Dr. Angela Frank-Briggs MB BS

Dr. Jonathan Ottor MB BS

Department of Pediatrics

*University of Port Harcourt Teaching
Hospital*

Box 141, Uniport

University of Port Harcourt

Port Harcourt , Nigeria

Abstract

Childhood mortality is much higher in developing countries than in developed ones. A study of 98 children aged 0 to 16 years who died while on admission in the University of Port Harcourt Teaching Hospital between 1st January and 31st December 2000 was undertaken to determine the all-cause mortality. Out of a total admission of 1777 children, there were 98 deaths (5.1%). Most of the causes of death were preventable. The largest single cause was birth asphyxia (20.4%) followed by malaria (10.2%), anemia (10.2%) and tetanus (9.2%). Neonates predominated accounting for 62 deaths (63.3%). This study shows that although the percentage of children who die on admission in our hospital is high there has been a fall in the death rate compared to an earlier study when the mortality was 11.6%. Emphasis should be laid on prevention and early treatment of the causes as most of these are preventable.

Key Words

Mortality, childhood, Africa, Nigeria

Introduction

The prevalence of childhood mortality in third world countries is very high. In Africa and South Asia alone, it accounts for 90% of the world total¹. A previous study in Nigeria² showed an overall childhood mortality of 11.6%. In the majority of developed countries, the mortality among children is estimated to be 0.02 per 1000 childhood population³. Most African countries record an all-cause childhood mortality of around 26.5%⁴ with diseases such as malaria and HIV (AIDS) predominating.



Figure 1



Figure 2



Figure 3



Figure 4

Figure 1 shows a 14 years old boy unconscious from cerebral malaria. Figure 2 shows a 6 years old girl unconscious with flaccid paraplegia from meningococcal meningitis. Figure 3 shows a 9 years old with Burkitt's Lymphoma of the jaw. Figure 4 shows a 4 years old who had

The following study was undertaken to ascertain the all-cause mortality among children in the urban city of Port Harcourt.

PATIENTS AND METHODS

This study took place at the University of Port Harcourt Teaching Hospital a tertiary institution which serves a population of 10 million.

From January to December 2000, photographs of severely ill children admitted into the Children's wards and Children's emergency rooms were taken with the parents' consent. The case notes of all admitted children were assembled for analysis at the end of the study period. Data extracted from the case notes included name, hospital number, age, sex, address, date of admission, investigations, diagnosis on admission, treatment, date of death, diagnosis at death, and post mortem findings. The various causes of death were expressed as the percentage of the total number of deaths.

RESULTS

Figures 1 to 11 are photographs of severely

There were 98 deaths (5.1%) out of a total of 1777 admissions between January to December 2000. The male to female ratio was 1:1.04. Table 1 outlines the various causes of death. The largest single cause was birth asphyxia (20.4%) followed by malaria (10.2%), anemia (10.2%), and tetanus (9.2%). The age range was 1 day-16 yrs, (mean 5.3 yrs). Neonates predominated accounting for 62 deaths (63.3%), (Figure 12). A post mortem was done in only one child and the final diagnosis was leukemia.

DISCUSSION

Port Harcourt, one of the fastest developing cities in the country is the capital of the Rivers State which is considered one of the richest Oil Producing States in Nigeria. An estimated 40% of the population are aged 0-16 years.

This study shows a mortality rate of 5.1% which is much higher than the 0.02 per 1000 children seen in developed countries³ but comparatively lower than that of 11.6% recorded in Ilorin - western Nigeria². It is also much lower than a community based study done in Kenya which showed an all cause mortality rate of 26.5%⁴. In Ghana, another community based study recorded a rate of 23.9% per one thousand children per year⁵. These mortality rates in under developed countries remain unacceptably high. The discrepancies in the two Nigerian studies may reflect better facilities in the health care delivery system since 1987 when the earlier study was done. It may also indicate a better standard of living in this oil rich city compared to the middle belt of Nigeria which is considered less wealthy.

The
commonest
causes
of
death
were
birth
asphyxia
(20.4%),
malaria



Figure 5




Figure 6

(10.2%),
anemia
(10.2%),
and
tetanus
(9.2%).
These
are



Figure 7

preventable.

Our
findings
contrast
sharply
with
the
common
causes
of
admission
and
deaths



Figure 9

in
developed

countries.

A
study
done

Figure 5 shows a 5 years old who had acute renal failure secondary to black water fever (severe malaria). Beside him is a bottle containing his bloody urine. He is on blood transfusion as his Haemoglobin had dropped to 2Gms%. Figure 6 shows an 11years old with tuberculous lymphadenitis. She had disseminated tuberculosis. Figure 7 shows severe burns of the sole of the foot in a 3 years old who had febrile convulsions. The parents had deliberately cast his foot into the fire to arrest the convulsion. Figure 8 is a post mortem finding of haemorrhagic brain of a child who died from leukemia. Figure 9 shows a 3month old baby with subcostal indrawing secondary to bronchopneumonia. Figure 10 shows a neonate with tetanus. The spasms draw the hands across the chest.

in Nottingham, UK showed that the commonest presenting problems to an accident and emergency department in a children's hospital were trauma or surgical (69%) as opposed to medical (27%)⁶. In the United States of America the major causes of death in infancy and childhood are congenital heart defects, whose mortality rate has decreased to 0.02 per 1000 population³. In Holland there is an increased mortality risk in children with epilepsy (0.5 per 1000 children) compared with the general population of children⁷.

There was no death from trauma in this study. This was a surprise finding. The explanation is unclear. It is however possible that most childhood deaths from road traffic accidents occur at the site of the accident and the victims do not therefore reach the hospital.



Figure 8



Figure 10

The prevalence rates and deaths from tetanus remain disturbing. In 1990, an immunization coverage of 95% was achieved in Nigeria with the help of UNICEF and Nigerian Federal Ministry of Health. This has fallen to 40%⁸. Factors contributing to the low immunization coverage include ignorance, low literacy rate, poverty and poor accessibility to Health Centers by the majority of the population who live around the creeks and islands of the delta basin area of the Rivers State⁹.

Pediatric HIV/AIDS infections accounted for only 1% of deaths in this study. There is an increased number of pediatric HIV (AIDS) cases in Nigeria in recent times¹⁰. Most of the cases abscond as soon as the diagnosis is made because of the associated social taboo coupled with a lack of anti retroviral drugs whose prices are not within the reach of most Nigerians. Another factor in the low prevalence of HIV/AIDS in this study is the fact that serology for HIV is not done routinely and even when indicated relations who have to pay for the test find it too expensive. It is therefore likely that the contribution of HIV/AIDS to the mortality rate in this study is underestimated.

Harmful practices such as burning a child to abort a febrile convulsion or forcing a child to drink cow's urine¹¹ or other concoctions whose chemistry is unknown, still abound. Specific educative programmes should be targeted at these.

Post mortem examination was performed in only one child. There is a worrying dearth of post mortem examinations in this institution and in other parts of Nigeria¹². Parents refuse post mortem examinations as being a waste of time. This is similar to findings in Zambia and other parts of Africa¹³.

In summary, although the percentage deaths on admission among children in Nigeria is less than it was over a decade ago, it remains high. The majority of the causes of death are preventable. Efforts to further reduce the mortality should be directed at prevention and early treatment.

REFERENCES

1. Costello A White H. Reducing global inequalities in child health. *Archives of Disease in Childhood* 2001; **84**:98-102
2. Fagbule D and Joiner KT. Pattern of Childhood Mortality at the University of Ilorin Teaching Hospital. *Nigerian Journal of Pediatrics* 1987; **14**: 1-5.
3. Boneva R S, Botto LD, Moore CA, Yang Q, Correa A, Erickson JD. Mortality associated with congenital heart defects in the United States: trends and racial disparities, 1979-1997. *Circulation* 2001;**103**:2376-81.
4. McElroy PD, Kuile FO ter, Hightower AW, Hawly WA, Phillipshoward PA, Oloo AJ, Lal AA, Nahlen BL. All-cause mortality among young children in western Kenya VI: the Asembo Bay Cohort Project. *American Journal of Tropical Medicine and Hygiene* 2001; **64**(1-2 Suppl): 18.
5. Allotey P, Reidpath D. International Programs, Key Centre for Women's Health, Faculty of Medicine and Dentistry. *Social and Scientific Medicine* 2001; **52**: 1007-12
6. Armon K, Stephen T, Gabriel V, Macfaul R, Eccletus P, Werneke U, Smith S, 2001 Determining the common medical presenting problems to an accident and emergency department. *Archives of Disease in Childhood* 2001; **84**:390-392
7. Callenbach PM, Westendorp RG, Geerts AT, Arts WF, Peeters EA, Van, Donselaar CA, Peters AC, Stroink H, Brouwer OF. Mortality risk in children with epilepsy: the dutch study of epilepsy in childhood. *Pediatrics* 2001;**107**: 1259-63.
8. UNICEF. The progress of nations. UNICEF 1998
9. Ogunbekun IO Financing medical care in Nigeria: Whither social insurance? *Nigerian Medical Practitioner* 1990;**19**:63-65
10. Bello CSS, Mawak JD, Otubu JAM. Human Immunodeficiency virus (HIV) and treponema pallidum infections in mothers and their babies at delivery in Jos. *Tropical Journal of Obstetrics and Gynaecology* 1997; **14**: 46-50
11. Ayorinde F O, Avery JW, Adekile AD, Ojewole JA, Odebiyi OO. Chemistry of a Nigerian Herbal Preparation (Cow's Urine

- 12. Onuigbo MAC. Acute renal failure in south-eastern Nigeria : A twenty-four months prospective analysis
- 13. Lishimpi K, Chintu C, Lucas S, Mudenda V, Kaluwaji J, Story A, Maswahu D, Bhat G, Nunn AJ, Zumla A. Necropsies in African children: consent dilemmas for parents and guardians. Archives of Disease in Childhood 2001; 84:463-467



Figure 11

Figure 11 shows a 2 years old with severe measles.




Figure 12: Histogram showing the number of deaths in the various age groups. Click to enlarge

In Association with **amazon.co.uk** SEARCH GO! All Products

No image available	Transactions of the First Annual Mee... American Association for the Study &... Buy Now	No image available	The healthy start initiative available from National Maternal and... Buy Now		The Late Life Legacy of Very Early L... Gabriele Dobhammer Our Price: £54.00 Buy Now
	Infant Feeding and Its Influence on... Charles Henry Felix Routh Our Price: £15.83 Buy Now		Birth, Poverty And Wealth - A Study... Richard M. Titmuss Our Price: £14.45 Buy Now	No image available	Mortality Statistics Population Censuses & Surveys Office... Buy Now
No image available	Infant Mortality of Indian Muslims Mohammed Omer Bajkhaif, Kuttan Mahad... Buy Now		Nutrition and Health in Developing C... P. Piot, Richard D. Semba, Martin W... Our Price: £107.50 Buy Now	No image available	Mortality Statistics Population Censuses & Surveys Office... Buy Now

[Books](#) | [Electronics & Photo](#) | [Music](#) | [DVD](#) | [Video](#) | [Software](#) | [PC & Video Games](#) | [Kitchen & Home](#) | [Toys](#)

Purchase these excellent books on Infant and child mortality at a discount from Amazon, UK through this online journal

● *N.B. It is essential to read this journal - and especially this paper as it contains tables and high resolution pictures - under a screen resolution of 1600 x 1200 dpi or more. If the resolution is less than this, you may see broken or overlapping tables, pictures overlapping text or other anomalies. It is strongly advised to switch over to this resolution to read this journal - and especially this paper. These pages are viewed best in Netscape Navigator 4.7 and above. All pictures appearing in this paper were submitted to this journal in very high resolution. However because of space constraints, the pictures have been reduced in resolution. Readers wanting to see the pictures in high resolution, may want to contact the author herself*

*Corresponding author and requests for reprints (and high resolution pictures):


Professor Felicia U. Eke
Department of Pediatrics, Box 141, Uniport, University of Port Harcourt Port Harcourt, NIGERIA
Tel/Fax: 234 84 231 337

● [Email Felicia Eke by clicking here](#)

You've been on Professor Felicia U. Eke's Paper for seconds.

● *N.B. It is essential to read this journal - and especially this paper as it contains several tables and high resolution graphics - under a screen resolution of 1600 x 1200 dpi or more, and preferably on a 17" or bigger monitor. If the resolution is less than this, you may see broken or overlapping tables/graphics, graphics overlying text or other anomalies. It is strongly advised to switch over to this resolution to read this journal - and especially this paper. These pages are viewed best in Netscape Navigator 4.7 and above.*

-Anil Aggrawal



j Web j www.geradts.com

[Sign My Guestbook](#)



[View My Guestbook](#)

● Click [here](#) to contact us.



You are Visitor No:



since July 22, 2000
when this page was created.

● This page has been constructed and maintained by Dr. Anil Aggrawal, Professor of Forensic Medicine, at the Maulana Azad Medical College, New Delhi-110002. You may want to give me the feedback to make this pages better. Please be kind enough to write your comments in the guestbook maintained above. These comments would help me make these pages better.

🌐 **IMPORTANT NOTE: ALL PAPERS APPEARING IN THIS ONLINE JOURNAL ARE COPYRIGHTED BY "ANIL AGGRAWAL'S INTERNET JOURNAL OF FORENSIC MEDICINE AND TOXICOLOGY" AND MAY NOT BE REPOSTED, REPRINTED OR OTHERWISE USED IN ANY MANNER WITHOUT THE WRITTEN PERMISSION OF THE WEBMASTER**

