



 **Current Issue**


 **Browse Issues**

 **Search**




 **About this Journal**

 **Instruction to Authors**

 **Online Submission**

 **Subscription**

 **Contact Us**



 **RSS Feed**

Acta Medica Iranica

2009;47(4) : 102-108

Simultaneous determination of metoprolol, propranolol and phenol red in samples from rat in situ intestinal perfusion studies

Parvin Zakeri-Milani, Hadi Valizadeh,, Yadollah Azarmi, Mohammad Barzegar Jalali, Hosniyeh Tajerzadeh

Abstract:

Single-pass intestinal perfusion technique (SPIP) is the most used classic technique employed in the study of intestinal absorption of compounds in which a non-absorbable marker such as phenol red is used to correct the water flux. A simple and rapid reversed-phase high performance liquid chromatographic method with UV detection at 227 nm was developed for simultaneous quantitation of propranolol and metoprolol along with phenol red for in-situ permeability studies. The mobile phase was a mixture of 55% methanol, 45% of 0.05 M KH₂PO₄ aqueous solution (adjusted to pH 6) and 0.2 % (v/v) triethylamine. Analysis was run at a flow rate of 1 ml/min with a 9 min run time. The calibration curves were linear for all three compounds ($r > 0.999$) across the concentration range of 7.5-125 µg/ml with a limit of detection of 4.24, 2.18 and 8.57 ng/ml and limit of quantification of 14, 7.2 and 28.3 ng/ml for metoprolol, propranolol and phenol red respectively. The coefficient of variation for intra-assay and inter-assay precision was less than 8% and the accuracy was between 93.6-107%. Using the SPIP technique and the suggested HPLC method for sample analysis, the mean values of 0.49×10^{-4} (± 0.19) cm/sec and 0.32×10^{-4} (± 0.09) cm/sec were obtained for propranolol and metoprolol intestinal permeability coefficients respectively.

Keywords:

Metoprolol . Phenol red . liquid chromatography . Permeability

TUMS ID: 2383

Full Text HTML  Full Text PDF  240 KB

top ▲

[Home](#) - [About](#) - [Contact Us](#)

TUMS E. Journals 2004-2009
Central Library & Documents Center
Tehran University of Medical Sciences

Best view with Internet Explorer 6 or Later at 1024*768 Resolutions