



 **Current Issue**

 **Browse Issues**

 **Search**



 **About this Journal**

 **Instruction to Authors**

 **Online Submission**

 **Subscription**

 **Contact Us**



 **RSS Feed**

Acta Medica Iranica

2009;47(4) : 25-26

CONSTITUENTS OF THE ESSENTIAL OIL OF CHIMONANTHUS ERA GRANCE LINDLE

KATAYOUN JAVIDN1A, RAMTN MIRI, MARYAM CHERIKI ABBAS SHAFIEE

Abstract:

The constituent of the essential oil of *Chimonanthus fragrance* Lindle (Calycanthaceae) were characterized by GLC and GC/MS. Twenty components representing 97% of the oil composition of which fifteen were sesquiterpenes were identified. The major components were p-Elementene, p-Caryophyllene, γ -Cadinene, γ -Bisabolene, p-Elementone and a-Eudesmol.

Keywords:

[Chimonanthus fragrance](#)

TUMS ID: 4347

Full Text HTML  Full Text PDF  192 kB

top ▲

[Home](#) - [About](#) - [Contact Us](#)

TUMS E. Journals 2004-2009
Central Library & Documents Center
Tehran University of Medical Sciences

Best view with Internet Explorer 6 or Later at 1024*768 Resolutions