

异常黑胆质成熟剂合用5-氟尿嘧啶对移植性EAC肿瘤的增效减毒作用

热娜古丽·艾则孜¹, 帕尔哈提·塔衣尔², 艾尼瓦尔·艾克木³, 艾山江·阿帕尔⁴, 祖拉亚提⁵, 陆明¹, 塔依尔江·吐尔逊⁶, 哈木拉提·吾甫尔⁶

1. 新疆医科大学附属中医医院, 乌鲁木齐 830000;
2. 新疆医科大学第一附属医院, 乌鲁木齐 830054;
3. 新疆医科大学药学院, 乌鲁木齐 830011;
4. 喀什地区第一人民医院, 喀什 844000;
5. 长春师范大学化学学院, 长春 130000;
6. 新疆医科大学维吾尔医学院, 乌鲁木齐 830011

Synergistic Attenuation of Abnormal Savda Munziq Combination with 5-Fluorouracil on Transplantated EAC Tumor

AI ZEZI Renaguli¹, TAYIR Parhat², AIKEMU Ainiwaer³, APER Aishanjan⁴, Zuliayati⁵, LU Ming¹, TUERXUN Tayierjiang⁶, UPUR Halmurat⁶

1. The Affiliated Traditional Chinese Medicine Hospital, Xinjiang Medical University, Urumqi 830000, China;
2. The First Affiliated Hospital, Xinjiang Medical University, Urumqi 830054, China;
3. College of Pharmacy, Xinjiang Medical University, Urumqi 830011, China;
4. The First People's Hospital of Kashgar, Kashgar 840000, China;
5. College of Chemistry, Changchun Normal University, Changchun 130000, China;
6. Faculty of Traditional Uighur Medicine, Xinjiang Medical University, Urumqi 830011, China

[摘要](#) [图/表](#) [参考文献](#) [相关文章 \(6\)](#)