

NAVIGATING IN THE CAYLEY GRAPH OF $SL_2(\mathbb{F}_p)$ AND APPLICATIONS TO HASHING

LISA BROMBERG, VLADIMIR SHPILRAIN, AND ALINA VDOVINA

ABSTRACT. Cayley hash functions are based on a simple idea of using a pair of (semi)group elements, A and B , to hash the 0 and 1 bit, respectively, and then to hash an arbitrary bit string in the natural way, by using multiplication of elements in the (semi)group. In this paper, we focus on hashing with 2×2 matrices over \mathbb{F}_p . Since there are many known pairs of 2×2 matrices over \mathbb{Z} that generate a free monoid, this yields numerous pairs of matrices over \mathbb{F}_p , for a sufficiently large prime p , that are candidates for collision-resistant hashing. However, this trick can “backfire”, and lifting matrix entries to \mathbb{Z} may facilitate finding a collision. This “lifting attack” was successfully used by Tillich and Zémor in the special case where two matrices A and B generate (as a monoid) the whole monoid $SL_2(\mathbb{Z}_+)$. However, in this paper we show that the situation with other, “similar”, pairs of matrices from $SL_2(\mathbb{Z})$ is different, and the “lifting attack” can (in some cases) produce collisions in the *group* generated by A and B , but not in the positive *monoid*. Therefore, we argue that for these pairs of matrices, there are no known attacks at this time that would affect security of the corresponding hash functions. We also give explicit lower bounds on the length of collisions for hash functions corresponding to some particular pairs of matrices from $SL_2(\mathbb{F}_p)$.

1. INTRODUCTION

Hash functions are easy-to-compute compression functions that take a variable-length input and convert it to a fixed-length output. Hash functions are used as compact representations, or digital fingerprints, of data and to provide message integrity. Basic requirements are well known:

- (1) *Preimage resistance* (sometimes called *non-invertibility*): it should be computationally infeasible to find an input which hashes to a specified output;
- (2) *Second pre-image resistance*: it should be computationally infeasible to find a second input that hashes to the same output as a specified input;
- (3) *Collision resistance*: it should be computationally infeasible to find two different inputs that hash to the same output.

A challenging problem is to determine mathematical properties of a hash function that would ensure (or at least, make it likely) that the requirements above are met.

Research of the second author was partially supported by the NSF grant CNS-1117675.

Early suggestions (especially the SHA family) did not really use any mathematical ideas apart from the Merkle-Damgard construction for producing collision-resistant hash functions from collision-resistant compression functions (see e.g. [9]); the main idea was just to “create a mess” by using complex iterations (this is not meant in a derogatory sense, but just as an opposite of using mathematical structure one way or another).

An interesting direction worth mentioning is constructing hash functions that are provably as secure as underlying assumptions, e.g. as discrete logarithm assumptions; see [2] and references therein. These hash functions however tend to be not very efficient. For a general survey on hash functions we refer to [9].

Another direction, relevant to the present paper, is using a pair of elements, A and B , of a semigroup S , such that the Cayley graph of the semigroup generated by A and B is expander, in the hope that such a graph would have a large girth and therefore there would be no short relations. Probably the most popular implementation of this idea so far is the Tillich-Zémor hash function [16]. We refer to [11] and [13] for a more detailed survey on Cayley hash functions.

The Tillich-Zémor hash function, unlike functions in the SHA family, is *not* a block hash function, i.e., each bit is hashed individually. More specifically, the “0” bit is hashed to a particular 2×2 matrix A , and the “1” bit is hashed to another 2×2 matrix B . Then a bit string is hashed simply to the product of matrices A and B corresponding to bits in this string. For example, the bit string 1000110 is hashed to the matrix BA^3B^2A .

Tillich and Zémor use matrices A, B from the group $SL_2(R)$, where R is a commutative ring (actually, a field) defined as $R = \mathbb{F}_2[x]/(p(x))$. Here \mathbb{F}_2 is the field with two elements, $\mathbb{F}_2[x]$ is the ring of polynomials over \mathbb{F}_2 , and $(p(x))$ is the ideal of $\mathbb{F}_2[x]$ generated by an irreducible polynomial $p(x)$ of degree n (typically, n is a prime, $127 \leq n \leq 170$); for example, $p(x) = x^{131} + x^7 + x^6 + x^5 + x^4 + x + 1$. Thus, $R = \mathbb{F}_2[x]/(p(x))$ is isomorphic to \mathbb{F}_{2^n} , the field with 2^n elements.

Then, the matrices A and B are:

$$A = \begin{pmatrix} \alpha & 1 \\ 1 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} \alpha & \alpha+1 \\ 1 & 1 \end{pmatrix},$$

where α is a root of $p(x)$.

This particular hash function was successfully attacked in [4] and [12]; see also [10] for a more general attack approach.

Another idea for hashing with matrices is to use a pair of 2×2 matrices, A and B , over \mathbb{Z} that generate a free monoid, and then reduce the entries modulo a large prime p to get matrices over \mathbb{F}_p . Since there cannot be an equality of two different products of positive powers of A and B unless at least one of the entries in at least one of the products is $\geq p$, this gives a lower bound on the minimum length of bit strings where a collision may occur. This lower bound is going to be on the order of $\log p$; we give more precise bounds for some particular examples of A and B in our Section 3.

The first example of a pair of matrices over \mathbb{Z} that generate a free monoid is:

$$A(1) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad B(1) = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}.$$

These matrices are obviously invertible, so they actually generate the whole group $SL_2(\mathbb{Z})$. This group is not free, but the *monoid* generated by $A(1)$ and $B(1)$ is free, and this is what matters for hashing because only positive powers of $A(1)$ and $B(1)$ occur in hashing. However, the fact that these two matrices generate the whole $SL_2(\mathbb{Z})$ yields an attack on the corresponding hash function (where the matrices $A(1)$ and $B(1)$ are considered over \mathbf{F}_p , for a large p), see [15], where a collision is found by using Euclidean algorithm on the entries of a matrix.

At this point, we note that a pair of matrices

$$A(x) = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}, \quad B(y) = \begin{pmatrix} 1 & 0 \\ y & 1 \end{pmatrix}$$

generate a free subgroup of $SL_2(\mathbb{Z})$ if $xy \geq 4$.

In Section 2, we consider the following pair of matrices:

$$A(2) = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}, \quad B(2) = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}.$$

By using a result from an old paper of Sanov [14] and combining it with the attack on hashing with $A(1)$ and $B(1)$ offered in [15], we show that there is an efficient heuristic algorithm that finds circuits of length $O(\log p)$ in the Cayley graph of the *group* generated by $A(2)$ and $B(2)$, considered as matrices over \mathbf{F}_p . However, this has no bearing on the (in)security of the hash function based on $A(2)$ and $B(2)$ since in hashing only positive powers of $A(2)$ and $B(2)$ are used, and group relations of length $O(\log p)$ produced by the mentioned algorithm will involve negative as well as positive powers with overwhelming probability.

To conclude the Introduction, we mention that the fact that Cayley graphs of the groups generated by $A(1), B(1)$ and by $A(2), B(2)$ are expanders was known for a couple of decades (see e.g. [7] for all background facts on expanders), but the same property of the Cayley graph relevant to $A(3), B(3)$ was a famous 1-2-3 question of Lubotzky, which was settled in the positive in [1] using results from [5].

We also give a word of caution: while intuitively, we think of expander graphs as graphs with a large (“expanding”) girth, this is not necessarily the case in general. In particular, first explicit examples of expander graphs due to Margulis[8] have bounded girth.

2. HASHING WITH $A(2)$ AND $B(2)$ AND CIRCUITS IN THE CAYLEY GRAPH

In this section, motivated by hashing with the matrices $A(2)$ and $B(2)$ considered as matrices over \mathbf{F}_p , we discuss circuits in the relevant Cayley graph.

Tillich and Zémor [15] offered an attack on the hash function based on $A(1)$ and $B(1)$ (again, considered as matrices over \mathbf{F}_p). To the best of our knowledge, this is the only published attack on that hash function. In this section we explain why this particular attack should not work with the matrices $A(2)$ and $B(2)$, and this

therefore leaves the door open for using these matrices (over \mathbf{F}_p , for a sufficiently large p) for hashing.

First we explain, informally, what appears to be the reason why the attack from [15] should not work with $A(2)$ and $B(2)$. The reason basically is that, while $A(1)$ and $B(1)$ (considered over \mathbb{Z}) generate (as a monoid!) the whole monoid of 2×2 matrices over \mathbb{Z} with positive entries, with the matrices $A(2)$ and $B(2)$ the situation is much less transparent. There is a result from an old paper by Sanov [14] that says: the *subgroup* of $SL_2(\mathbb{Z})$ generated by $A(2)$ and $B(2)$ consists of *all* matrices of the form $\begin{pmatrix} 1+4m_1 & 2m_2 \\ 2m_3 & 1+m_4 \end{pmatrix}$, where all m_i are arbitrary integers. This, however, does not tell much about the *monoid* generated by $A(2)$ and $B(2)$. In fact, a generic matrix of the above form would *not* belong to this monoid. This is not surprising because: (1) $A(2)$ and $B(2)$ generate a free group, by another result of Sanov [14]; (2) the number of different elements represented by *all* freely irreducible words in $A(2)$ and $B(2)$ of length $m \geq 2$ is $4 \cdot 3^{m-1}$, whereas the number of different elements represented by *positive* words of length $m \geq 2$ is 2^m . Thus, the share of matrices in the above form representable by positive words in $A(2)$ and $B(2)$ is exponentially negligible.

What Tillich and Zémor’s “lifting attack” [15] can still give is an efficient heuristic algorithm that finds relations of length $O(\log p)$ in the *group* generated by $A(2)$ and $B(2)$, considered as matrices over \mathbf{F}_p . We describe this algorithm below because we believe it might be useful in other contexts, although it has no bearing on the security of the hash function based on $A(2)$ and $B(2)$ since in hashing only positive powers of $A(2)$ and $B(2)$ are used, and group relations of length $O(\log p)$ produced by the algorithm mentioned above will involve negative as well as positive powers with overwhelming probability, even if p is rather small. What one would need to attack the hash function corresponding to $A(2)$ and $B(2)$ is a result, similar to Sanov’s, describing all matrices in the *monoid* generated by $A(2)$ and $B(2)$.

We are now going to use a combination of the attack on $A(1)$ and $B(1)$ offered in [15] with the aforementioned result of Sanov [14] to find relations of length $O(\log p)$ in the *group* generated by $A(2)$ and $B(2)$.

Theorem 1. *There is an efficient heuristic algorithm that finds particular relations of the form $w(A(2), B(2)) = 1$, where w is a group word of length $O(\log p)$, and the matrices $A(2)$ and $B(2)$ are considered over \mathbb{F}_p .*

Proof. It was shown in [15] that:

(a) For any prime p , there is an efficient heuristic algorithm that finds positive integers k_1, k_2, k_3, k_4 such that the matrix $\begin{pmatrix} 1+k_1p & k_2p \\ k_3p & 1+k_4p \end{pmatrix}$ has determinant 1 and all k_i are of about the same magnitude $O(p^2)$.

(b) A generic matrix from part (a) has an efficient factorization (in $SL_2(\mathbb{Z})$) in a product of positive powers of $A(1)$ and $B(1)$, of length $O(\log p)$. (This obviously yields a collision in $SL_2(\mathbf{F}_p)$ since the matrix from part (a) equals the identity matrix in $SL_2(\mathbf{F}_p)$.)

Now we combine these results with the aforementioned result of Sanov the following way. We are going to multiply a matrix from (a) (call it M) by a matrix from $SL_2(\mathbb{Z})$ (call it S) with very small (between 0 and 5 by the absolute value) entries, so that the resulting matrix $M \cdot S$ has the form as in Sanov's result. Since the matrix M , by the Tillich-Zémor results, has an efficient factorization (in $SL_2(\mathbb{Z})$) in a product $w(A(1), B(1))$ of powers of $A(1)$ and $B(1)$ of length $O(\log p)$, the same holds for the matrix $M \cdot S$. Then, since the matrix $M \cdot S$ is in "Sanov's form", we know that it is, in fact, a product of powers of $A(2)$ and $B(2)$.

Now we need one more ingredient to efficiently re-write a product of $A(1)$ and $B(1)$ into a product of $A(2)$ and $B(2)$ without blowing up the length too much. This procedure is provided by Theorem 2.3.10 in [3]. We cannot explain it without introducing a lot of background material, but the fact is that, since the group $SL_2(\mathbb{Z})$ is hyperbolic (whatever that means) and the subgroup generated by $A(2)$ and $B(2)$ is quasiconvex (whatever that means), there is a quadratic time algorithm (in the length of the word $w(A(1), B(1))$) that re-writes $w(A(1), B(1))$ into a $u(A(2), B(2))$ such that $w(A(1), B(1)) = u(A(2), B(2))$ and the length of u is bounded by a constant (independent of w) times the length of w .

Thus, what is now left to complete the proof is to exhibit, for all possible matrices M as in part (a) above, particular "small" matrices S such that $M \cdot S$ is in "Sanov's form". We are therefore going to consider many different cases corresponding to possible combinations of residues modulo 4 of the entries of the matrix M (recall that M has to have determinant 1), and in each case we are going to exhibit the corresponding matrix S such that $M \cdot S$ is in "Sanov's form". Denote by \hat{M} the matrix of residues modulo 4 of the entries of M . Since the total number of cases is too large, we consider matrices \hat{M} "up to a symmetry".

$$\begin{aligned}
 \text{(1)} \quad \hat{M} &= \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, S = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \\
 \text{(2)} \quad \hat{M} &= \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}, S = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \\
 \text{(3)} \quad \hat{M} &= \begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix}, S = \begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix} \\
 \text{(4)} \quad \hat{M} &= \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}, S = \begin{pmatrix} 1 & 3 \\ 3 & 2 \end{pmatrix} \\
 \text{(5)} \quad \hat{M} &= \begin{pmatrix} 2 & 3 \\ 3 & 1 \end{pmatrix}, S = \begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix} \\
 \text{(6)} \quad \hat{M} &= \begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix}, S = \begin{pmatrix} 0 & 1 \\ 3 & 2 \end{pmatrix} \\
 \text{(7)} \quad \hat{M} &= \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix}, S = \begin{pmatrix} 1 & 3 \\ 3 & 2 \end{pmatrix} \\
 \text{(8)} \quad \hat{M} &= \begin{pmatrix} 2 & 3 \\ 3 & 3 \end{pmatrix}, S = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \\
 \text{(9)} \quad \hat{M} &= \begin{pmatrix} 3 & 3 \\ 0 & 1 \end{pmatrix}, S = \begin{pmatrix} 3 & 3 \\ 0 & 1 \end{pmatrix}
 \end{aligned}$$

$$(10) \hat{M} = \begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}, S = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$(11) \hat{M} = \begin{pmatrix} 3 & 0 \\ 1 & 3 \end{pmatrix}, S = \begin{pmatrix} -1 & 0 \\ 1 & -1 \end{pmatrix}$$

$$(12) \hat{M} = \begin{pmatrix} 3 & 0 \\ 2 & 3 \end{pmatrix}, S = \begin{pmatrix} -1 & 0 \\ 2 & -1 \end{pmatrix}$$

$$(13) \hat{M} = \begin{pmatrix} 3 & 0 \\ 3 & 3 \end{pmatrix}, S = \begin{pmatrix} -1 & 0 \\ 3 & -1 \end{pmatrix}$$

$$(14) \hat{M} = \begin{pmatrix} 3 & 2 \\ 2 & 3 \end{pmatrix}, S = \begin{pmatrix} -1 & 2 \\ 2 & -5 \end{pmatrix}$$

This completes the proof. □

To conclude this section, we point out an example of re-writing a word in $A(1)$ and $B(1)$ into a word in $A(2)$ and $B(2)$. All matrices here are considered over \mathbb{Z} .

$$A(1)B(1)A(1)B(1)A(1)B(1) = A(2)A(2)B(2)^{-1}B(2)^{-1}A(2)^{-1}A(2)^{-1}B(2)B(2).$$

We see that even in this simple example, both positive and negative powers of $A(2)$ and $B(2)$ are required.

3. GIRTH OF THE CAYLEY GRAPH RELEVANT TO $A(k)$ AND $B(k)$

Our starting point here is the following observation: the entries of matrices that are products of length n of positive powers of $A(k)$ and $B(k)$ exhibit the fastest growth (as functions of n) if $A(k)$ and $B(k)$ alternate in the product: $A(k)B(k)A(k)B(k) \cdots$. More formally:

Proposition 1. *Let $w_n(a, b)$ be an arbitrary positive word of even length n , and let $W_n = w_n(A(k), B(k))$, with $k \geq 2$. Let $C_n = (A(k) \cdot B(k))^{\frac{n}{2}}$. Then: (a) the sum of entries in any row of C_n is at least as large as the sum of entries in any row of W_n ; (b) the largest entry of C_n is at least as large as the largest entry of W_n .*

Proof. First note that multiplying a matrix X by $A(k)$ on the right amounts to adding to the second column of X the first column multiplied by k . Similarly, multiplying X by $B(k)$ on the right amounts to adding to the first column of X the second column multiplied by k . This means, in particular, that when we build a word in $A(k)$ and $B(k)$ going left to right, elements of the first row change independently of elements of the second row. Therefore, we can limit our considerations to pairs of positive integers, and the result will follow from the following

Lemma 1. *Let (x, y) be a pair of positive integers and let $k \geq 2$. One can apply transformations of the following two kinds: (1) transformation R takes (x, y) to $(x, y + kx)$; (2) transformation L takes (x, y) to $(x + ky, y)$. Among all sequences of these transformations of the same length, the sequence where R and L alternate results in: (a) the largest sum of elements in the final pair; (b) the largest maximum element in the final pair.*

Proof. We are going to prove (a) and (b) simultaneously using induction by the length of a sequence of transformations. Suppose our lemma holds for all sequences of length at most $m \geq 2$, with the same initial pair (x, y) . Suppose the final pair after m alternating transformations is (X, Y) . Without loss of generality, assume that $X < Y$. That means the last applied transformation was R . Now applying L to (X, Y) gives $(X + kY, Y)$, while applying R to (X, Y) gives $(X, Y + kX)$. Since $X + kY > Y + kX$, applying L results in a larger sum of elements as well as in a larger maximum element. Thus, we have a sequence of $(m + 1)$ alternating transformations, and now we have to consider one more case.

Suppose some sequence of m transformations applied to (x, y) results in a pair (X', Y') with $X' + Y' < X + Y$, $Y' < Y$, but $X' > X$. Then applying L to this pair gives $(X' + kY', Y')$, and the sum is $X' + Y' + kY' < X + Y + kY$ since $X' + Y' < X + Y$ and $Y' < Y$. The maximum element of the pair $(X' + kY', Y')$ is $X' + kY' = X' + Y' + (k - 1)Y'$. Again, since $X' + Y' < X + Y$ and $Y' < Y$, we have $X' + kY' < X + kY$. This completes the proof of the lemma and the proposition. \square

\square

\square

This motivates us to consider powers of the matrix $C(k) = A(k)B(k)$ to get to entries larger than p “as quickly as possible”.

3.1. Powers of $C(2) = A(2)B(2)$. The matrix $C(2)$ is $\begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$. If we denote $(C(2))^n = \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix}$, then the following recurrence relations are easily proved by induction on n :

$$a_n = 5a_{n-1} + 2b_{n-1}; \quad b_n = c_n = 2a_{n-1} + b_{n-1}; \quad d_n = a_{n-1}.$$

Combining the recurrence relations for a_n and b_n , we get $2b_n = a_n - a_{n-1}$, so $2b_{n-1} = a_{n-1} - a_{n-2}$. Plugging this into the displayed recurrence relation for a_n gives

$$a_n = 6a_{n-1} - a_{n-2}.$$

Similarly, we get

$$b_n = 6b_{n-1} - b_{n-2}.$$

Solving these recurrence relations (with appropriate initial conditions), we get

$$a_n = \left(\frac{1}{2} + \frac{1}{\sqrt{8}}\right)(3 + \sqrt{8})^n + \left(\frac{1}{2} - \frac{1}{\sqrt{8}}\right)(3 - \sqrt{8})^n, \quad b_n = \frac{1}{\sqrt{8}}(3 + \sqrt{8})^n - \frac{1}{\sqrt{8}}(3 - \sqrt{8})^n.$$

Thus, a_n is the largest entry of $(C(2))^n$, and we conclude that no entry of $(C(2))^n$ is larger than p as long as $n < \log_{3+\sqrt{8}} p$. Since $C(2) = A(2)B(2)$ is a product of two generators, $(C(2))^n$ has length $2n$ as a word in the generators $A(2)$ and $B(2)$.

Therefore, no two positive words of length $\leq m$ in the generators $A(2)$ and $B(2)$ (considered as matrices over \mathbb{F}_p) can be equal as long as

$$m < 2 \log_{3+\sqrt{8}} p = \log_{\sqrt{3+\sqrt{8}}} p,$$

so we have the following

Corollary 1. *There are no collisions of the form $u(A(2), B(2)) = v(A(2), B(2))$ if positive words u and v are of length less than $\log_{\sqrt{3+\sqrt{8}}} p$. In particular, the girth of the Cayley graph of the semigroup generated by $A(2)$ and $B(2)$ (considered as matrices over \mathbb{F}_p) is at least $\log_{\sqrt{3+\sqrt{8}}} p$.*

The base of the logarithm here is $\sqrt{3+\sqrt{8}} \approx 2.4$. Thus, for example, if p is on the order of 2^{256} , then there are no collisions of the form $u(A(2), B(2)) = v(A(2), B(2))$ if positive words u and v are of length less than 203.

We also note, in passing, that our Proposition 1 also holds without the assumption on the words $w_n(a, b)$ to be positive if we consider the absolute values of the matrix entries and their sums. Our lower bound on the girth of the Cayley graph of the group generated by $A(2)$ and $B(2)$ therefore improves (in this particular case) the lower bound given in [5], where the base of the logarithm in the lower bound is 3.

3.2. Powers of $C(3) = A(3)B(3)$. The matrix $C(3)$ is $\begin{pmatrix} 10 & 3 \\ 3 & 1 \end{pmatrix}$. If we let $(C(3))^n = \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix}$, and use the fact that $(C(3))^n = C(3) \cdot (C(3))^{n-1}$, we get the following recurrence relations:

$$\begin{aligned} a_n &= 10a_{n-1} + 3b_{n-1} \\ b_n &= 3a_{n-1} + b_{n-1} = c_n \\ d_n &= a_{n-1} \end{aligned}$$

Combining these recurrence relations for a_n and b_n , we get

$$a_n = 11a_{n-1} - a_{n-2}, \quad b_n = 11b_{n-1} - b_{n-2}.$$

Solving these recurrence relations with the initial conditions $a_1 = 10, a_2 = 109$ and $b_1 = 3, b_2 = 33$, we get

$$\begin{aligned} a_n &= \left(\frac{9}{2\sqrt{117}} + \frac{1}{2} \right) \left(\frac{11 + \sqrt{117}}{2} \right)^n + \left(\frac{1}{2} - \frac{9}{2\sqrt{117}} \right) \left(\frac{11 - \sqrt{117}}{2} \right)^n, \\ b_n &= \frac{3}{\sqrt{117}} \left(\frac{11 + \sqrt{117}}{2} \right)^n - \frac{3}{\sqrt{117}} \left(\frac{11 - \sqrt{117}}{2} \right)^n. \end{aligned}$$

From this we see that a_n is the largest entry of $(C(3))^n$, so no entry of $(C(3))^n$ is larger than p if $n < \log_{\frac{11+\sqrt{117}}{2}} p$. Since $C(3)$ is a product of two generators, $A(3)$ and $B(3)$, we have:

Corollary 2. *There are no collisions of the form $u(A(3), B(3)) = v(A(3), B(3))$ if positive words u and v are of length less than $2 \log_{\frac{11+\sqrt{117}}{2}} p = \log_{\sqrt{\frac{11+\sqrt{117}}{2}}} p$. In particular, the girth of the Cayley graph of the semigroup generated by $A(3)$ and $B(3)$ (considered as matrices over \mathbb{F}_p) is at least $\log_{\sqrt{\frac{11+\sqrt{117}}{2}}} p$.*

The base of the logarithm here is $\sqrt{\frac{11+\sqrt{117}}{2}} \approx 3.3$. For example, if p is on the order of 2^{256} , then there are no collisions of the form $u(A(2), B(2)) = v(A(2), B(2))$ if positive words u and v are of length less than 149.

4. CONCLUSIONS

We have analyzed the girth of the Cayley graph of the group and the monoid generated by pairs of matrices $A(k)$ and $B(k)$ (considered over \mathbb{F}_p), for various $k \geq 1$. Our conclusions are:

- The “lifting attack” by Tillich and Zémor [15] that produces explicit relations of length $O(\log p)$ in the monoid generated by $A(1)$ and $B(1)$, can be used in combination with an old result by Sanov [14] and some results from the theory of automatic groups [3] to efficiently produce explicit relations of length $O(\log p)$ in the group generated by $A(2)$ and $B(2)$.
- Generically, relations produced by this method will involve negative as well as positive powers of $A(2)$ and $B(2)$, and therefore will *not* produce collisions for the corresponding hash function.
- In the absence of a result for $A(3)$ and $B(3)$ similar to Sanov’s result for $A(2)$ and $B(2)$, at this time there is no known efficient algorithm for producing explicit relations of length $O(\log p)$ even in the group generated by $A(3)$ and $B(3)$, let alone in the monoid generated by this pair. At the same time, such relations do exist by the pigeonhole principle.
- We have computed an explicit lower bound for the length of relations in the monoid generated by $A(2)$ and $B(2)$; the lower bound is $\log_b p$, where the base b of the logarithm is approximately 2.4. For the monoid generated by $A(3)$ and $B(3)$, we have a similar lower bound, with base b of the logarithm approximately equal to 3.3.
- We conclude that at this time, there are no known attacks on hash functions corresponding to the pair $A(2)$ and $B(2)$ or $A(3)$ and $B(3)$ and therefore no visible threat to their security.

Acknowledgement. We are grateful to Ilya Kapovich for helpful comments, in particular for pointing out the relevance of some results from [3] to our work. We are also grateful to Harald Helfgott for useful discussions.

REFERENCES

- [1] J. Bourgain, A. Gamburd, *Uniform expansion bounds for Cayley graphs of $SL_2(\mathbf{F}_p)$* . Ann. of Math. (2) **167** (2008), 625–642.
- [2] S. Contini, A. K. Lenstra and R. Steinfeld, *VSH, an Efficient and Provable Collision Resistant Hash Function*, in: Eurocrypt 2006, Lecture Notes Comp. Sci. **4004** (2006), 165–182.

- [3] D. B. A. Epstein, J. Cannon, D. F. Holt, S. V. F. Levy, M. S. Paterson, W. P. Thurston, *Word processing in groups*. Jones and Bartlett Publishers, Boston, MA, 1992.
- [4] M. Grassl, I. Ilić, S. Magliveras, R. Steinwandt, *Cryptanalysis of the Tillich-Zémor hash function*, J. Cryptology **24** (2011), 148-156.
- [5] H. A. Helfgott, *Growth and generation in $SL_2(\mathbb{Z}/p\mathbb{Z})$* Ann. of Math. (2) **167** (2008), 601–623.
- [6] M. Larsen, *Navigating the Cayley graph of $SL_2(\mathbb{F}_p)$* , Int. Math. Res. Notes **27** (2003), 1465-1471.
- [7] A. Lubotzky, *Discrete groups, expanding graphs and invariant measures*, Progress in Mathematics **125**, Birkhäuser Verlag, Basel, 1994.
- [8] G. A. Margulis, *Explicit constructions of concentrators*, Problems of Information Transmission **9** (1973), no. 4, 325–332.
- [9] A. Menezes, P. van Oorschot and S. Vanstone, *Handbook of Applied Cryptography*, CRC Press, 1997.
- [10] C. Mullan and B. Tsaban, *Short collision search in arbitrary SL_2 homomorphic hash functions*, preprint, <http://arxiv.org/abs/1306.5646>
- [11] C. Petit, *On graph-based cryptographic hash functions*, PhD thesis, 2009.
- [12] C. Petit, J. Quisquater, *Preimages for the Tillich-Zémor hash function*, in: SAC 10, Lecture Notes Comp. Sci. **6544** (2010), 282-301.
- [13] C. Petit and J.-J. Quisquater, *Rubik's for cryptographers*, Notices Amer. Math. Soc. **60** (2013), 733–739.
- [14] I. N. Sanov, *A property of a representation of a free group* (Russian), Doklady Akad. Nauk SSSR (N. S.) **57** (1947), 657–659.
- [15] J.-P. Tillich and G. Zémor, *Group-theoretic hash functions*, in Proceedings of the First French-Israeli Workshop on Algebraic Coding, Lecture notes Comp. Sci. **781** (1993), 90–110.
- [16] J.-P. Tillich and G. Zémor, *Hashing with SL_2* , in CRYPTO 1994, Lecture Notes Comp. Sci. **839** (1994), 40–49.

GRADUATE CENTER, CITY UNIVERSITY OF NEW YORK
E-mail address: lisa.bromberg@gmail.com

DEPARTMENT OF MATHEMATICS, THE CITY COLLEGE OF NEW YORK, NEW YORK, NY
 10031
E-mail address: shpil@groups.sci.ccny.cuny.edu

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEWCASTLE, NEWCASTLE
 UPON TYNE, NE1 7RU, U.K.
E-mail address: alina.vdovina@ncl.ac.uk