

A Non-Markovian Process with Unbounded p-Variation

Martynas Manstavicius, *University of Connecticut, USA*

Abstract

A recent theorem by M. Manstavicius (2004) provided a link between a certain function of transition probabilities of a strong Markov process and the boundedness of the p-variation of its trajectories. Here one assumption of that theorem is relaxed and an example is constructed to show that the Markov property cannot be easily dispensed with.

Full text: [PDF](#) | [PostScript](#)

Pages: 17-28

Published on: February 25, 2005

Bibliography

1. J. R. Kinney. Continuity properties of sample functions of Markov processes. *Trans. Amer. Math. Soc.* 74 (1953), 280--302. [Math. Review MR0053428](#)
2. E. B. Dynkin. Criteria of continuity and of absence of discontinuities of the second kind for trajectories of a Markov random process. *Izv. Akad. Nauk SSSR Ser. Mat.* 16 (1952), 563--572, (in Russian). [Math. Review MR0052055](#)
3. M. Manstavicius. The p-variation of strong Markov processes. *Ann. Probab.* 32 (2004), no. 3A, 2053-2066. [Math. Review MR2073185](#)
4. R. M. Dudley and R. Norvaiša. An Introduction to p-Variation and Young Integrals. Maphysto Lecture notes 1 (1998, revised 1999) ([pdf file](#)) *Math.* Review number not available.

Research Support Tool

[Capture Cite](#)
[View Metadata](#)
[Printer Friendly](#)

▼ [Context](#)

[Author Address](#)

▼ [Action](#)

[Email Author](#)
[Email Others](#)

