

A Note on Reflecting Brownian Motions

Florin Soucaliuc, *Université Paris-Sud*
Wendelin Werner, *Université Paris-Sud and IUF*

Abstract

We give another proof of the following result from a joint paper with Bálint Tóth: *A Brownian motion reflected on an independent time-reversed Brownian motion is a Brownian motion.*

Full text: [PDF](#) | [PostScript](#)

Pages: 117-122

Published on: January 17, 2002

Bibliography

1. K. Burdzy, D. Nualart (2000): Brownian motion reflected on Brownian motion, *Probab. Th. Rel. Fields*, to appear.
2. J. Pitman, M. Yor (1996): Decomposition at the maximum for excursions and bridges of one-dimensional diffusions, in *Ito's stochastic calculus and probability theory*, Springer, 293-310. [Math. Review 98f:60153](#)
3. F. Soucaliuc (2001): Réflexion, coalescence et retournement du temps pour certaines familles de diffusions, PhD Thesis, Université Paris-Sud.
4. F. Soucaliuc, B. Tóth, W. Werner (2000): Reflection and coalescence between one-dimensional Brownian paths, *Ann. Inst. Henri Poincaré Probab. Statist.* 36, 509-536, [Math. Review 2002a:60139](#)
5. B. Tóth, W. Werner (1998): The true self-repelling motion, *Probab. Th. Rel. Fields*, 111, 375-452 [Math. Review 99i:60092](#)

Research Support Tool

[Capture Cite](#)
[View Metadata](#)
[Printer Friendly](#)

▼ [Context](#)

[Author Address](#)

▼ [Action](#)

[Email Author](#)
[Email Others](#)

