

## $L^1$ -Norm of Infinitely Divisible Random Vectors and Certain Stochastic Integrals

Michael B. Marcus, *The City College of CUNY*  
Jan Rosinski, *University of Tennessee*

### Abstract

Equivalent upper and lower bounds for the  $L^1$  norm of Hilbert space valued infinitely divisible random variables are obtained and used to find bounds for different types of stochastic integrals.

Full text: [PDF](#) | [PostScript](#)

Pages: 15-29

Published on: January 10, 2001

### Bibliography

1. De la Pena, V.H. and Gine, E. (1999), *Decoupling. From dependence to independence*. Springer-Verlag, New York. [Math. Review 99k:60044](#)
2. Dunford, N and Schwartz, J.T. (1958), *Linear Operators, Part I: General Theory. Decoupling*. Wiley, New York. [Math. Review 90g:47001a](#)
3. I.I. Gihman and A.V. Skorohod. (1975), *The theory of Stochastic Processes, Vol. 2*. Springer, Berlin. [Math. Review 51:11656](#)
4. Klass, M.J. (1980), *Precision bounds for the relative error in the approximation of  $E|S_n|$  and extensions*. Ann. Probab. 8, 350--367. [Math. Review 82h:60097](#)
5. Kwapien, S. and Woyczynski, W.A. (1992), *Random Series and Stochastic Integrals: Single and Multiple*. Birkh"auser, Boston. [Math. Review 94k:60074](#)
6. Marcus, M.B. and Rosinski, J. *Continuity of stochastic integrals with respect to infinitely divisible random measures*. 2000 (preprint) Math. Review number not available.
7. Parthasarathy, K.R. (1967), *Probability measures on metric spaces*. Academic Press, New York. [Math. Review 37:2271](#)

### Research Support Tool

[Capture Cite](#)  
[View Metadata](#)  
[Printer Friendly](#)

▼ [Context](#)

[Author Address](#)

▼ [Action](#)

[Email Author](#)  
[Email Others](#)

