



A note on global Markov properties for mixed graphs

[Michael Eichler](#)

(Submitted on 15 Jul 2011 ([v1](#)), last revised 16 Nov 2011 (*this version*, [v2](#)))

Global Markov properties in mixed graphs are usually formulated in terms of the path-oriented m-separation or by use of augmented graphs (similar to moral graphs in the case of directed acyclic graphs). We provide an alternative characterization that can be easily implemented.

Comments: 5 pages, 1 figure

Subjects: **Methodology (stat.ME)**

Cite as: [arXiv:1107.3036](#) [stat.ME]

(or [arXiv:1107.3036v2](#) [stat.ME] for this version)

Submission history

From: Michael Eichler [[view email](#)]

[\[v1\]](#) Fri, 15 Jul 2011 10:29:07 GMT (14kb)

[\[v2\]](#) Wed, 16 Nov 2011 13:41:39 GMT (125kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

stat.ME

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[stat](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

