

Prevention of Oxidative Stress Injury to Sperm

ASHOK AGARWAL, SUSHIL A. PRABAKARAN,
AND TAMER M. SAID

From the Center for Advanced Research in Human Reproduction, Infertility, and Sexual Function, Glickman Urological Institute and Department of Obstetrics-Gynecology, The Cleveland Clinic Foundation, Cleveland, Ohio.

The greatest paradox of aerobic respiration is that oxygen, which is essential for energy production, may also be detrimental because it leads to the production of reactive oxygen species (ROS) (Saleh and Agarwal, 2002). When levels of reactive oxygen species (ROS) overwhelm the body's antioxidant defense system, oxidative stress (OS) occurs. OS is a condition in which the elevated levels of ROS damage cells, tissues, or organs (Moller et al, 1996; Sharma and Agarwal, 1996; Saleh et al, 2003).

ROS are free radicals that play a significant role in many of the sperm physiological processes such as capacitation, hyperactivation, and sperm-oocyte fusion (Aitken et al, 2004; Allamaneni et al, 2004; de Lamirande et al, 1998). However, they also trigger many pathological processes in the male reproductive system, and these processes have been implicated in cancers of the bladder and prostate, as well as in male infertility (Bankson et al, 1993; Hietanen et al, 1994; Agarwal and Saleh, 2002).

Spermatozoa are sensitive to OS because they lack cytoplasmic defenses (Donnelly et al, 1999; Saleh and Agarwal, 2002). Moreover, the sperm plasma membrane contains lipids in the form of polyunsaturated fatty acids, which are vulnerable to attack by ROS. ROS, in the presence of polyunsaturated fatty acids, triggers a chain of chemical reactions called lipid peroxidation (Agarwal et al, 1994; Kobayashi et al, 2001; Zalata et al, 2004). ROS can also damage DNA by causing deletions, mutations, and other lethal genetic effects (Moustafa et al, 2004; Tominaga et al, 2004). It is difficult to block the OS-

Correspondence to: Dr Ashok Agarwal, Professor of Surgery, Lerner College of Medicine and Director, Center for Advanced Research in Human Reproduction, Infertility and Sexual Function, Glickman Urological Institute and Department of Obstetrics-Gynecology, The Cleveland Clinic Foundation, 9500 Euclid Avenue, Desk A19.1, Cleveland, Ohio 44195 (e-mail: agarwaa@ccf.org).

Received for publication January 25, 2005; accepted for publication June 9, 2005.

DOI: 10.2164/jandrol.05016

Minireview

induced injury to cells or tissues because ROS are continuously produced by cellular aerobic metabolism (Davies, 2000). Several clinical trials are currently attempting to minimize the toxic effects of OS on human spermatozoa (Agarwal et al, 2004). In this review, we highlight the various protective measures available to minimize OS-induced injury to spermatozoa.

Sources of ROS in Semen

There are two main sources of ROS in semen: leukocytes and immature spermatozoa (Garrido et al, 2004). Of these, leukocytes are considered to be the primary source (Aitken et al, 1992). Leukocytes, particularly neutrophils and macrophages, have been associated with excessive ROS production that ultimately leads to sperm dysfunction (Aitken and Baker, 1995; Aitken et al, 1997; Hendin et al, 1999; Ochsendorf, 1999; Pasqualotto et al, 2000; Saleh et al, 2002; Shalika et al, 1996; Sharma et al, 2001).

Spermatozoa produce ROS mainly when a defect occurs during spermiogenesis that results in retention of cytoplasmic droplets (Gomez et al, 1996; Zini et al, 1993). A strong positive correlation exists between immature spermatozoa and ROS production, which in turn negatively affects the sperm quality (Gil-Guzman et al, 2001; Said et al, 2004).

The two main sites of ROS production are the mitochondrion and the sperm plasma membrane. The mitochondrion is the powerhouse of respiration. Hence, it is the major site of ROS generation, which is produced through the nicotinamide adenine dinucleotide-dependent oxido-reductase pathway (Gavella and Lipovac, 1992). In contrast, the sperm plasma membrane produces ROS through the nicotinamide adenine dinucleotide phosphate-dependent oxidase system (Aitken et al, 1992; Agarwal et al, 2003). Xanthine oxidase—a key enzyme in purine catabolism—is also involved in the production of ROS in spermatozoa (Aitken et al, 1993; Sanocka et al, 1996).

Oxidative Damage to Spermatozoa by Life-Style Behaviors

Damage Caused by Pollution—Environmental pollution and radiation can generate various ROS such as hydrogen peroxide (H_2O_2), superoxide anion ($O_2^{\bullet-}$), and hydroxyl radical ($\bullet OH$) (Gate et al, 1999). According to an Italian study on tollgate workers, traffic pollution damages sperm and may reduce fertility in young and middle-aged men

(De Rosa et al, 2003). The workers in that study showed poorer sperm quality than age-matched men living in the same area who were not exposed to the same level of automobile pollution. In addition, sperm function tests revealed that the workers had less active spermatozoa. In the subset of workers with abnormal sperm parameters, sperm count and viability were inversely correlated with lead (Pb) levels, whereas sperm motility, viability, membrane function, nuclear DNA integrity, linearity, and amplitude of sperm lateral movement were inversely correlated with methemoglobin levels—a marker for nitrogen oxide (De Rosa et al, 2003).

Damage Caused by Smoking—Smoking induces OS either by increasing levels of oxidants originating from the smoke or by decreasing levels of antioxidants in the seminal plasma (Fraga et al, 1996). Therefore, smokers may need to increase their intake of antioxidants to compensate for this deficiency (Bui et al, 1992). According to a study conducted by our group, smoking greatly increases levels of leukocyte and ROS (Saleh et al, 2002). Other studies have reported that tobacco adversely affects sperm quality (ie, concentration, motility, and morphology) (Kunzle et al, 2003, 2004). Smoking also predominately affects sperm DNA, as evidenced from the increased level of 8-deoxyguanosine (8-oxodG)—a marker for oxidative DNA damage (Potts et al, 1999). Therefore, smoking may not affect the fertilization rate per se, but it will increase the risk of heritable mutations (Kunzle et al, 2003; Lewin et al, 1991; Rubes et al, 1998; Vine et al, 1996).

Oxidative Damage to Spermatozoa by Infection

Sperm damage can be caused either by the invading pathogens or by the defense mechanisms that are employed against them (Ochsendorf, 1999). For example, when microorganisms invade the human body, it produces polymorphonuclear leukocytes and macrophages, which are the major sources of ROS production (Ochsendorf, 1999; Saran et al, 1999; Zalata et al, 2004). Prostatitis and accessory gland infection increase OS, which severely damages spermatozoa (Potts and Pasqualotto, 2003). In addition, a past infection by the sexually transmitted *Neisseria gonorrhoea* is associated with leukocytospermia (Trum et al, 1998). Although there is no direct evidence that *Neisseria gonorrhoea* directly increases ROS production, the associated leukocytospermia is well known to produce ROS (Trum et al, 1998). According to a study by Depuydt et al, leukocytospermia and male accessory glands infection reduce a man's fertilizing potential by affecting sperm parameters both in vitro and in vivo (Depuydt et al, 1998).

Iatrogenic Oxidative Damage to Spermatozoa

Prolonged in vitro incubation of semen samples that contain high levels of immature spermatozoa before sperm

processing increases the risk of OS damage to mature spermatozoa (Gil-Guzman et al, 2001). In a study by Aitken and Clarkson, it was reported that repeated centrifugation mechanically injures spermatozoa and increases ROS production (Aitken and Clarkson, 1988). OS may also damage sperm during cryopreservation. A study by Bilodeau et al revealed that ROS generated during freeze-thaw cycles are detrimental to sperm function and that levels of antioxidants were diminished during each cycle (Bilodeau et al, 2000).

Strategies for OS Prevention

Modifying Life-Style Habits—As reported earlier, because smoking is known to increase OS in seminal plasma, patients should be advised to abstain from it. It is also of extreme importance to avoid excessive exposure to environmental pollution in working and living conditions, as environmental pollution increases the OS-related sperm damage.

Antioxidants—Antioxidants are the main defense against OS induced by free radicals. There are prevention antioxidants and scavenger antioxidants. Prevention antioxidants such as metal chelators and metal-binding proteins block the formation of new ROS, whereas scavenger antioxidants remove the ROS that have already formed.

Prevention Antioxidants—Transition metal ions, mainly iron, are involved in the generation of the highly reactive $\bullet\text{OH}$ by Fenton's reaction (Biemond et al, 1984; Ochsendorf, 1999). They stimulate lipid peroxidation by decomposing the peroxides into peroxy and alkoxy radicals, which in turn causes the chain reaction of lipid peroxidation. Metal chelators such as transferrin, lactoferrin, and ceruloplasmin that are present in human semen control lipid peroxidation of the sperm plasma membrane, protecting its integrity (Sanocka and Kurpisz, 2004). Other metal chelators such as ethylene diamine tetraacetic acid, 1,10-phenanthroline, and neocuproine have been shown to reduce sperm DNA damage in fishes (Bruskov et al, 2002). However, similar data are lacking in human.

In vitro supplementation of metal chelators such as DL-penicillamine, 2,3-dimercaptopropan-1 sulphonate and meso-2,3-dimercapto-succinimic acid showed enhancement of sperm quality during assisted reproductive technique (Henkel and Schill, 2003). In a study by Wroblewski et al, incubating sperm with D-penicillamine significantly increased sperm motility (Wroblewski et al, 2003).

Cadmium (Cd), a class B element, is capable of replacing zinc, thereby exerting its toxic effect in spermatogenesis. Metallothionein removes Cd from the body by binding to it, which results in improved spermatogenesis, maturation, and capacitation of spermatozoa (Omu and Fernandes, 2001).

Scavenger Antioxidants—Dietary antioxidants form an essential part of the human antioxidant defense system.

Fruits and vegetables as well as daily dietary supplements constitute the potential sources of various antioxidants. The National Academy of Sciences has recommended 60 mg vitamin C per day for an adult male. The daily requirement of vitamin E varies from 50 to 800 mg, depending on the intake of fruits, vegetables, tea, or wine. Carotenoids and selenium work synergistically with vitamin E and have a recommended dietary allowances value of 1000 and 70 micrograms per day, respectively (National Academy of Sciences, 1989).

Oxidative stress may also be limited by using chain-breaking antioxidants such as vitamin E and vitamin C as drug supplements. Vitamin C is a major chain-breaking antioxidant and is present in the extracellular fluid. It neutralizes hydroxyl, superoxide, and hydrogen peroxide radicals and prevents sperm agglutination (Agarwal et al, 2004). In addition, it also helps recycle vitamin E (Sies et al, 1992). Vitamin C is found in reduced quantity in the seminal plasma of infertile men (Lewis et al, 1997). Vitamin C increases sperm counts *in vivo* in infertile male patients with oral doses ranging from 200–1000 mg/day (Dawson et al, 1987). The combination of flavonoids and vitamin C increases the effectiveness of both of these compounds (Mathiesen et al, 1996). They conserve the alpha-tocopherol content of low-density lipoprotein and delay the onset of lipid peroxidation (de Whalley et al, 1990). Quercetin and xanthohumol constitute some of the important dietary flavonoids.

The principal chain-breaking antioxidant vitamin E is present within the cell membrane. It neutralizes H₂O₂ and protects the plasma membrane from lipid peroxidation. In a randomized cross-over study, oral administration of 600 mg/day of vitamin E improved sperm function as assessed by the zona binding test (Kessopoulou et al, 1995).

Selenium is a necessary component for the synthesis of glutathione peroxidase. It works synergistically with vitamin E (Bartfay et al, 1998). According to a study conducted by Keskes-Ammar et al, 225 µg/day of selenium and 400 mg/day of vitamin E oral supplementation for 3 months significantly decreased malondialdehyde concentrations in seminal plasma and improved sperm motility (Keskes-Ammar et al, 2003). However these findings were not confirmed in another study (Iwanier and Zachara, 1995).

Carotenoids such as beta-carotene and lycopene also form an important component of the antioxidant defense (Gupta and Kumar, 2002). Beta-carotenes protect the plasma membrane against lipid peroxidation. Lycopene is found in abundance in tomatoes (Agarwal and Rao, 2000) and has a suggested daily intake of 5–10 mg/day. Lycopene has been shown to be twice as potent as beta-carotene and 10 times more potent than vitamin E in scavenging singlet oxygen and inhibiting lipid peroxidation in serum plasma (Di Mascio et al, 1989).


Glutathione is the most abundant antioxidant found in the body. It plays an important role in protecting lipids, proteins, and nucleic acids against oxidative damage. It combines with vitamin E and selenium to form glutathione peroxidase. In a placebo-controlled, double-blind, cross-over trial, 600 mg glutathione was administered for 2 months by intramuscular injection in 20 infertile men. Glutathione therapy significantly increased sperm motility, particularly forward progression (Lenzi et al, 1993).

Coenzyme Q-10 (CoQ-10) found in the sperm mid-piece (Lewin and Lavon, 1997) recycles vitamin E, prevents its prooxidant activity (Thomas et al, 1996; Thomas et al, 1997), and is also involved in energy production. In a study conducted on 32 infertile males, CoQ-10 inhibited H₂O₂ formation in seminal fluid and in seminal plasma (Alleva et al, 1997). In an *in vitro* study on 22 semen samples of asthenozoospermic men, incubation with 50 µM of CoQ-10 significantly increased sperm motility. In the same study, *in vivo* supplementation of 60 mg of CoQ-10 in 17 infertile men improved their fertilization rate without changing their semen parameters (Lewin and Lavon, 1997).

Zinc and copper are trace metals that constitute a part of superoxide dismutase—a key enzymatic antioxidant. Their adequate intake is necessary to maintain the optimal functioning level of these enzymes. At present, it is estimated that the average daily intake in United States is 12.3 mg of zinc and 900 µg of copper per person. Caution should be exercised during intake of these trace metals because some of them can catalyze the reactions that lead to the formation of ROS. An *in vitro* study by Lloyd et al on salmon sperm DNA determined that at concentrations of 20–50 µM and above, these metal ions caused a maximum DNA strand breaks (Lloyd et al, 1998). No such studies are available in humans, and the *in vivo* dosage required for these concentrations in seminal plasma is still unknown.

Other antioxidants may also protect against OS, such as alpha lipoic acid and carnitines. Alpha lipoic acid (thiols) undergoes reduction to form dihydrolipoic acid, thereby regenerating other antioxidants such as vitamins C and E and reduced glutathione through redox cycling (Biewenga et al, 1997). In contrast, carnitines are dietary antioxidants that decrease ROS by removing excess intracellular toxic acetyl-CoA that are responsible for mitochondrial ROS production. Carnitines also improve sperm motility (Agarwal and Said, 2004; Vicari and Calogero, 2001).

However, the positive effects of antioxidants are still debatable. Multiple studies have found no improvement following antioxidant intake (Agarwal et al, 2004; Ford and Whittington, 1998; Martin-Du Pan and Sakkas, 1998; Moilanen and Hovatta, 1995; Rolf et al, 1999).



Suggested algorithm for evaluation and management of infertile patients with increased oxidative stress in their seminal plasma. Oxidative stress should be assessed as a part of the male infertility evaluation. ROS-TAC score was calculated by applying principal component analysis to normalized values of ROS and TAC in semen specimens from a population of proven fertile donors; scores below 30 indicate the presence of oxidative stress. TAC indicates total antioxidant capacity; ROS, reactive oxygen species; COMET, single-cell microgel electrophoresis; TUNEL, terminal deoxynucleotidyl transferase-mediated deoxyuridine triphosphate-nick end labeling; SCSA, sperm chromatin structure assay; and ART, assisted reproductive technology.

Prevention of Iatrogenic Oxidative Damage—The use of specific sperm preparation techniques has greatly reduced the OS associated with sperm handling and cryopreservation. Sperm separation techniques such as migration-sedimentation, density centrifugation gradient, and glass-wool filtration significantly reduce the level of ROS by removing leukocytes, which are the major source of ROS (Henkel and Schill, 2003).

In vitro supplements used during sperm preparation and assisted reproductive technique also help to protect spermatozoa against ROS. Moreover, adding antioxidants to the culture media neutralizes ROS produced by the leukocytes and immature spermatozoa and improves sperm-oocyte fusion (Irvine, 1996). In an in vitro study, rebamipide effectively scavenged ROS during sperm processing and cryopreservation (Park et al, 2003). In another study performed on samples from 25 male partners of infertile couples, in vitro supplementation with superoxide dismutase and catalase prevented lipid peroxidation

of the sperm plasma membrane by ROS and contributed to the recovery of high-quality spermatozoa after freezing-thawing procedures (Rossi et al, 2001). Similarly, it has been found that adding glutathione and hypotaurine protects spermatozoa against oxidative damage induced by H₂O₂ (Donnelly et al, 2000).

Pentoxifylline—a methylxanthine derivative that inhibits phosphodiesterase—has been approved by the US Food and Drug Administration for use in humans. It has a beneficial effect on sperm motility and acrosome reaction and reduces the O₂^{•-} release by the human spermatozoa (Agarwal et al, 2004; Henkel and Schill, 2003; McKinney et al, 1996). N-acetyl-L-cysteine—a precursor of glutathione—reduces the ROS production in human ejaculate (Agarwal et al, 2004; Oeda et al, 1997), as well as ROS-induced DNA damage (Lopes et al, 1998). The use of vitamin E in vitro has been also documented to improve sperm motility and viability (Verma and Kanwar, 1999). Hughes et al (1998) has determined that in vitro

supplementation of vitamins C, E, and urate separately has protective effects on sperm DNA integrity on irradiation (Hughes et al, 1998).

Conclusions

Spermatozoa are under a continuous influence of OS because of excessive generation of ROS. Although spermatozoa are affected in different ways by OS, there are sufficient antioxidant protections that can decrease the progression of the damage. However, when an imbalance exists between levels of ROS and the natural antioxidant defenses, various measures can be used to protect spermatozoa against the OS-induced injury (Figure). Diet forms an important component of the antioxidant protection system; it supplies the major antioxidants such as vitamin C, vitamin E, and carotenoids. Therefore, food rich in these elements should form a part of the daily diet. For those patients who are suspected to have high levels of ROS, antioxidant supplements can be considered. Nevertheless, further studies are required to validate their use in this group of patients. In certain cases, it is also essential to modify certain lifestyle behaviors because many habits and environmental factors increase the production of ROS and affect fertility.

Another important method for decreasing OS is the use of antioxidants during various sperm processing techniques. Antioxidants decrease the oxidative damage to spermatozoa induced during these techniques. There are many controversies regarding the doses, types, and combinations that could be used in different sperm-handling techniques. Future research should address these issues to develop standard and reliable protocols.

Acknowledgments

We thank the Glickman Urological Institute at the Cleveland Clinic Foundation for research support, and Mrs Robin K. Verdi for secretarial support.

References

Agarwal A, Hamamah, S, Shekarriz M. Reactive oxygen species and fertilizing capacity of spermatozoa. *Contracept Fertil Sex*. 1994;22:327–330.

Agarwal A, Nallella KP, Allamaneni SS, Said TM. Role of antioxidants in treatment of male infertility: an overview of the literature. *Reprod Biomed Online*. 2004;8:616–627.

Agarwal A, Said TM. Carnitines and male infertility. *Reprod Biomed Online*. 2004;8:376–384.

Agarwal A, Saleh RA. Role of oxidants in male infertility: rationale, significance, and treatment. *Urol Clin North Am*. 2002;29:817–827.

Agarwal A, Saleh RA, Bedaiwy MA. Role of reactive oxygen species in the pathophysiology of human reproduction. *Fertil Steril*. 2003;79:829–843.

Agarwal S, Rao AV. Tomato lycopene and its role in human health and chronic diseases. *Can Med Assoc J*. 2000;163:739–744.

Aitken RJ, Baker HW. Seminal leukocytes: passengers, terrorists or good samaritans? *Hum Reprod*. 1995;10:1736–1739.

Aitken RJ, Buckingham D, Harkiss D. Use of a xanthine oxidase free radical generating system to investigate the cytotoxic effects of reactive oxygen species on human spermatozoa. *J Reprod Fertil*. 1993;97:441–450.

Aitken RJ, Buckingham D, West K, Wu FC, Zikopoulos K, Richardson DW. Differential contribution of leucocytes and spermatozoa to the generation of reactive oxygen species in the ejaculates of oligospermic patients and fertile donors. *J Reprod Fertil*. 1992;94:451–462.

Aitken RJ, Clarkson JS. Significance of reactive oxygen species and antioxidants in defining the efficacy of sperm preparation techniques. *J Androl*. 1988;9:367–376.

Aitken RJ, Fisher HM, Fulton N, Gomez E, Knox W, Lewis B, Irvine S. Reactive oxygen species generation by human spermatozoa is induced by exogenous NADPH and inhibited by the flavoprotein inhibitors diphenylene iodonium and quinacrine. *Mol Reprod Dev*. 1997;47:468–482.

Aitken RJ, Ryan AL, Baker MA, McLaughlin EA. Redox activity associated with the maturation and capacitation of mammalian spermatozoa. *Free Radic Biol Med*. 2004;36:994–1010.

Allamaneni SS, Naughton CK, Sharma RK, Thomas AJ Jr, Agarwal A. Increased seminal reactive oxygen species levels in patients with varicoceles correlate with varicocele grade but not with testis size. *Fertil Steril*. 2004;82:1684–1686.

Alleva R, Scararmucci A, Mantero F, Bompadre S, Leoni L, Littarru GP. The protective role of ubiquinol-10 against formation of lipid hydroperoxides in human seminal fluid. *Mol Aspects Med*. 1997;18(suppl):S221–S228.

Bankson DD, Kestin M, Rifai N. Role of free radicals in cancer and atherosclerosis. *Clin Lab Med*. 1993;13:463–480.

Bartfay WJ, Hou D, Brittenham GM, Bartfay E, Sole MJ, Lehotay D, Liu PP. The synergistic effects of vitamin E and selenium in iron-overloaded mouse hearts. *Can J Cardiol*. 1998;14:937–941.

Biamond P, van Eijk HG, Swaak AJ, Koster JF. Iron mobilization from ferritin by superoxide derived from stimulated polymorphonuclear leukocytes. Possible mechanism in inflammation diseases. *J Clin Invest*. 1984;73:1576–1579.

Biewenga GP, Haenen GR, Bast A. The pharmacology of the antioxidant lipoic acid. *Gen Pharmacol*. 1997;29:315–331.

Bilodeau JF, Chatterjee S, Sirard MA, Gagnon C. Levels of antioxidant defenses are decreased in bovine spermatozoa after a cycle of freezing and thawing. *Mol Reprod Dev*. 2000;55:282–288.

Bruskov VI, Malakhova LV, Masalimov ZK, Chernikov AV. Heat-induced formation of reactive oxygen species and 8-oxoguanine, a biomarker of damage to DNA. *Nucleic Acids Res*. 2002;30:1354–1363.

Bui MH, Sauty A, Collet F, Leuenberger P. Dietary vitamin C intake and concentrations in the body fluids and cells of male smokers and non-smokers. *J Nutr*. 1992;122:312–316.

Davies KJ. Oxidative stress, antioxidant defenses, and damage removal, repair, and replacement systems. *IUBMB Life*. 2000;50:279–289.

Dawson EB, Harris WA, Rankin WE, Charpentier LA, McGanity WJ. Effect of ascorbic acid on male fertility. *Ann N Y Acad Sci*. 1987;498:312–323.

de Lamirande E, Tsai C, Harakat A, Gagnon C. Involvement of reactive oxygen species in human sperm arcsome reaction induced by A23187, lysophosphatidylcholine, and biological fluid ultrafiltrates. *J Androl*. 1998;19:585–594.

De Rosa M, Zarrilli S, Paesano L, Carbone U, Boggia B, Petretta M, Maisto A, Cimmino F, Puca G, Colao A, Lombardi G. Traffic pollutants affect fertility in men. *Hum Reprod*. 2003;18:1055–1061.

de Whalley CV, Rankin SM, Hoult JR, Jessup W, Leake DS. Flavonoids

- inhibit the oxidative modification of low density lipoproteins by macrophages. *Biochem Pharmacol.* 1990;39:1743–1750.
- Depuydt C, Zalata A, Christophe A, Mahmoud A, Comhaire F. Mechanisms of sperm deficiency in male accessory gland infection. *Andrologia.* 1998;30(suppl 1):29–33.
- Di Mascio P, Kaiser S, Sies H. Lycopene as the most efficient biological carotenoid singlet oxygen quencher. *Arch Biochem Biophys.* 1989; 274:532–538.
- Donnelly ET, McClure N, Lewis SE. Antioxidant supplementation in vitro does not improve human sperm motility. *Fertil Steril.* 1999;72:484–495.
- Donnelly ET, McClure N, Lewis SE. Glutathione and hypotaurine in vitro: effects on human sperm motility, DNA integrity and production of reactive oxygen species. *Mutagenesis.* 2000;15:61–68.
- Ford WC, Whittington K. Antioxidant treatment for male subfertility: a promise that remains unfulfilled. *Hum Reprod.* 1998;13:1416–1419.
- Fraga CG, Motchnik PA, Wyrobek AJ, Rempel DM, Ames BN. Smoking and low antioxidant levels increase oxidative damage to sperm DNA. *Mutat Res.* 1996;351:199–203.
- Garrido N, Meseguer M, Simon C, Pellicer A, Remohi J. Pro-oxidative and anti-oxidative imbalance in human semen and its relation with male fertility. *Asian J Androl.* 2004;6:59–65.
- Gate L, Paul J, Ba GN, Tew KD, Tapiero H. Oxidative stress induced in pathologies: the role of antioxidants. *Biomed Pharmacother.* 1999;53: 169–180.
- Gavella M, Lipovac V. NADH-dependent oxidoreductase (diaphorase) activity and isozyme pattern of sperm in infertile men. *Arch Androl.* 1992;28:135–141.
- Gil-Guzman E, Ollero M, Lopez MC, Sharma RK, Alvarez JG, Thomas AJ Jr, Agarwal A. Differential production of reactive oxygen species by subsets of human spermatozoa at different stages of maturation. *Hum Reprod.* 2001;16:1922–1930.
- Gomez E, Buckingham DW, Brindle J, Lanzafame F, Irvine DS, Aitken RJ. Development of an image analysis system to monitor the retention of residual cytoplasm by human spermatozoa: correlation with biochemical markers of the cytoplasmic space, oxidative stress, and sperm function. *J Androl.* 1996;17:276–287.
- Gupta NP, Kumar R. Lycopene therapy in idiopathic male infertility—a preliminary report. *Int Urol Nephrol.* 2002;34:369–372.
- Hendin BN, Kolettis PN, Sharma RK, Thomas AJ Jr, Agarwal A. Varicocele is associated with elevated spermatozoal reactive oxygen species production and diminished seminal plasma antioxidant capacity. *J Urol.* 1999;161:1831–1834.
- Henkel RR, Schill WB. Sperm preparation for ART. *Reprod Biol Endocrinol.* 2003;1:108.
- Hietanen E, Bartsch H, Bereziat JC, Camus AM, McClinton S, Eremin O, Davidov L, Boyle P. Diet and oxidative stress in breast, colon and prostate cancer patients: a case-control study. *Eur J Clin Nutr.* 1994; 48:575–586.
- Hughes CM, Lewis SE, McKelvey-Martin VJ, Thompson W. The effects of antioxidant supplementation during Percoll preparation on human sperm DNA integrity. *Hum Reprod.* 1998;13:1240–1247.
- Irvine DS. Glutathione as a treatment for male infertility. *Rev Reprod.* 1996;1:6–12.
- Iwanier K, Zachara BA. Selenium supplementation enhances the element concentration in blood and seminal fluid but does not change the spermatozoal quality characteristics in subfertile men. *J Androl.* 1995; 16:441–447.
- Keskes-Ammar L, Feki-Chakroun N, Rebai T, Sahnoun Z, Ghazzi H, Hammami S, Zghal K, Fki H, Damak J, Bahloul A. Sperm oxidative stress and the effect of an oral vitamin E and selenium supplement on semen quality in infertile men. *Arch Androl.* 2003;49:83–94.
- Kessopoulou E, Powers HJ, Sharma KK, Pearson MJ, Russell JM, Cooke ID, Barratt CL. A double-blind randomized placebo cross-over controlled trial using the antioxidant vitamin E to treat reactive oxygen species associated male infertility. *Fertil Steril.* 1995;64:825–831.
- Kobayashi H, Gil-Guzman E, Mahran AM, Rakesh, Nelson DR, Thomas AJ Jr, Agarwal A. Quality control of reactive oxygen species measurement by luminol-dependent chemiluminescence assay. *J Androl.* 2001;22:568–574.
- Kunzle R, Mueller MD, Hanggi W, Birkhauser MH, Drescher H, Bersinger NA. Semen quality of male smokers and nonsmokers in infertile couples. *Fertil Steril.* 2003;79:287–291.
- Kunzle R, Mueller MD, Huber AW, Drescher H, Bersinger NA. Seasonality in human semen quality of smokers and non-smokers: effect of temperature. *Asian J Androl.* 2004;6:243–247.
- Lenzi A, Culasso F, Gandini L, Lombardo F, Dondero F. Placebo-controlled, double-blind, cross-over trial of glutathione therapy in male infertility. *Hum Reprod.* 1993;8:1657–1662.
- Lewin A, Gonen O, Orvieto R, Schenker JG. Effect of smoking on concentration, motility and zona-free hamster test on human sperm. *Arch Androl.* 1991;27:51–54.
- Lewin A, Lavon H. The effect of coenzyme Q10 on sperm motility and function. *Mol Aspects Med.* 1997;18(suppl):S213–S219.
- Lewis SE, Sterling ES, Young IS, Thompson W. Comparison of individual antioxidants of sperm and seminal plasma in fertile and infertile men. *Fertil Steril.* 1997;67:142–147.
- Lloyd DR, Carmichael PL, Phillips DH. Comparison of the formation of 8-hydroxy-2'-deoxyguanosine and single- and double-strand breaks in DNA mediated by fenton reactions. *Chem Res Toxicol.* 1998;11:420–427.
- Lopes S, Jurisicova A, Sun JG, Casper RF. Reactive oxygen species: potential cause for DNA fragmentation in human spermatozoa. *Hum Reprod.* 1998;13:896–900.
- Martin-Du Pan RC, Sakkas D. Is antioxidant therapy a promising strategy to improve human reproduction? Are anti-oxidants useful in the treatment of male infertility? *Hum Reprod.* 1998;13:2984–2985.
- Mathiesen L, Wang S, Halvorsen B, Malterud KE, Sund RB. Inhibition of lipid peroxidation in low-density lipoprotein by the flavonoid myricetin and ascorbic acid. *Biochem Pharmacol.* 1996;51:1719–1725.
- McKinney KA, Lewis SE, Thompson W. The effects of pentoxifylline on the generation of reactive oxygen species and lipid peroxidation in human spermatozoa. *Andrologia.* 1996;28:15–20.
- Moilanen J, Hovatta O. Excretion of alpha-tocopherol into human seminal plasma after oral administration. *Andrologia.* 1995;27:133–136.
- Moller P, Wallin H, Knudsen LE. Oxidative stress associated with exercise, psychological stress and life-style factors. *Chem Biol Interact.* 1996;102:17–36.
- Moustafa MH, Sharma RK, Thornton J, Mascha E, Abdel-Hafez MA, Thomas AJ Jr, Agarwal A. Relationship between ROS production, apoptosis and DNA denaturation in spermatozoa from patients examined for infertility. *Hum Reprod.* 2004;19:129–138.
- National Academy of Sciences. *Recommended Dietary Allowances.* 10th ed. National Academy Press. 1989.
- Ochsendorf FR. Infections in the male genital tract and reactive oxygen species. *Hum Reprod Update.* 1999;5:399–420.
- Oeda T, Henkel R, Ohmori H, Schill WB. Scavenging effect of N-acetyl-L-cysteine against reactive oxygen species in human semen: a possible therapeutic modality for male factor infertility? *Andrologia.* 1997;29:125–131.
- Omu EA, Fernandes S. The relationship between zinc/cadmium ratio in human semen: effect on immune response. *Kuwait Med J.* 2001;33: 38–43.
- Park NC, Park HJ, Lee KM, Shin DG. Free radical scavenger effect of rebamipide in sperm processing and cryopreservation. *Asian J Androl.* 2003;5:195–201.
- Pasqualotto FF, Sharma RK, Potts JM, Nelson DR, Thomas AJ, Agarwal

- A. Seminal oxidative stress in patients with chronic prostatitis. *Urology*. 2000;55:881–885.
- Potts JM, Pasqualotto FF. Seminal oxidative stress in patients with chronic prostatitis. *Andrologia*. 2003;35:304–308.
- Potts RJ, Newbury CJ, Smith G, Notarianni LJ, Jefferies TM. Sperm chromatin damage associated with male smoking. *Mutat Res*. 1999;423:103–111.
- Rolf C, Cooper TG, Yeung CH, Nieschlag E. Antioxidant treatment of patients with asthenozoospermia or moderate oligoasthenozoospermia with high-dose vitamin C and vitamin E: a randomized, placebo-controlled, double-blind study. *Hum Reprod*. 1999;14:1028–1033.
- Rossi T, Mazzilli F, Delfino M, Dondero F. Improved human sperm recovery using superoxide dismutase and catalase supplementation in semen cryopreservation procedure. *Cell Tissue Bank*. 2001;2:9–13.
- Rubes J, Lowe X, Moore D II, Perreault S, Slott V, Evenson D, Selevan SG, Wyrobek AJ. Smoking cigarettes is associated with increased sperm disomy in teenage men. *Fertil Steril*. 1998;70:715–723.
- Said TM, Agarwal A, Sharma RK, Mascha E, Sikka SC, Thomas AJ Jr. Human sperm superoxide anion generation and correlation with semen quality in patients with male infertility. *Fertil Steril*. 2004;82:871–877.
- Saleh RA, Agarwal A. Oxidative stress and male infertility: from research bench to clinical practice. *J Androl*. 2002;23:737–752.
- Saleh RA, Agarwal A, Kandirali E, Sharma RK, Thomas AJ, Nada EA, Evenson DP, Alvarez JG. Leukocytospermia is associated with increased reactive oxygen species production by human spermatozoa. *Fertil Steril*. 2002;78:1215–1224.
- Saleh RA, Agarwal A, Nada EA, El-Tonsy MH, Sharma RK, Meyer A, Nelson DR, Thomas AJ. Negative effects of increased sperm DNA damage in relation to seminal oxidative stress in men with idiopathic and male factor infertility. *Fertil Steril*. 2003;79(suppl 3):1597–1605.
- Saleh RA, Agarwal A, Sharma RK, Nelson DR, Thomas AJ Jr. Effect of cigarette smoking on levels of seminal oxidative stress in infertile men: a prospective study. *Fertil Steril*. 2002;78:491–499.
- Sanocka D, Kurpisz M. Reactive oxygen species and sperm cells. *Reprod Biol Endocrinol*. 2004;2:12.
- Sanocka D, Miesel R, Jędrzejczak P, Kurpisz MK. Oxidative stress and male infertility. *J Androl*. 1996;17:449–454.
- Saran M, Beck-Speier I, Fellerhoff B, Bauer G. Phagocytic killing of microorganisms by radical processes: consequences of the reaction of hydroxyl radicals with chloride yielding chlorine atoms. *Free Radic Biol Med*. 1999;26:482–490.
- Shalika S, Dugan K, Smith RD, Padilla SL. The effect of positive semen bacterial and Ureaplasma cultures on in-vitro fertilization success. *Hum Reprod*. 1996;11:2789–2792.
- Sharma RK, Agarwal A. Role of reactive oxygen species in male infertility. *Urology*. 1996;48:835–850.
- Sharma RK, Pasqualotto AE, Nelson DR, Thomas AJ Jr, Agarwal A. Relationship between seminal white blood cell counts and oxidative stress in men treated at an infertility clinic. *J Androl*. 2001;22:575–583.
- Sies H, Stahl W, Sundquist AR. Antioxidant functions of vitamins. Vitamins E and C, beta-carotene, and other carotenoids. *Ann N Y Acad Sci*. 1992;669:7–20.
- Thomas SR, Neuzil J, Stocker R. Cosupplementation with coenzyme Q prevents the prooxidant effect of alpha-tocopherol and increases the resistance of LDL to transition metal-dependent oxidation initiation. *Arterioscler Thromb Vasc Biol*. 1996;16:687–696.
- Thomas SR, Neuzil J, Stocker R. Inhibition of LDL oxidation by ubiquinol-10. A protective mechanism for coenzyme Q in atherogenesis? *Mol Aspects Med*. 1997;18(suppl):S85–S103.
- Tominaga H, Kodama S, Matsuda N, Suzuki K, Watanabe M. Involvement of reactive oxygen species (ROS) in the induction of genetic instability by radiation. *J Radiat Res (Tokyo)*. 2004;45:181–188.
- Trum JW, Mol BW, Pannekoek Y, Spanjaard L, Wertheim P, Bleker OP, van der Veen F. Value of detecting leukocytospermia in the diagnosis of genital tract infection in subfertile men. *Fertil Steril*. 1998;70:315–339.
- Verma A, Kanwar KC. Effect of vitamin E on human sperm motility and lipid peroxidation in vitro. *Asian J Androl*. 1999;1:151–154.
- Vicari E, Calogero AE. Effects of treatment with carnitines in infertile patients with prostatic-vesiculo-epididymitis. *Hum Reprod*. 2001;16:2338–2342.
- Vine MF, Tse CK, Hu P, Truong KY. Cigarette smoking and semen quality. *Fertil Steril*. 1996;65:835–842.
- Wroblewski N, Schill WB, Henkel R. Metal chelators change the human sperm motility pattern. *Fertil Steril*. 2003;79(suppl 3):1584–1589.
- Zalata AA, Ahmed AH, Allamaneni SS, Comhaire FH, Agarwal A. Relationship between acrosin activity of human spermatozoa and oxidative stress. *Asian J Androl*. 2004;6:313–318.
- Zini A, de Lamirande E, Gagnon C. Reactive oxygen species in semen of infertile patients: levels of superoxide dismutase- and catalase-like activities in seminal plasma and spermatozoa. *Int J Androl*. 1993;16:183–188.