

Spermicidal Activity of Bacterial Lipopolysaccharide Is Only Partly Due to Lipid A

HAMID HAKIMI,* IAN GEARY,* ALLAN PACEY,† AND ADRIAN ELEY*

From the *Division of Genomic Medicine, The Medical School; and the †Division of Clinical Sciences, The Jessop Wing, Central Sheffield University Hospitals Trust, The University of Sheffield, Sheffield, United Kingdom.

ABSTRACT: We have previously shown that co-incubation of *Chlamydia trachomatis* lipopolysaccharide (LPS) leads to premature sperm death by an apoptosis-like mechanism. It was always assumed that lipid A is the toxic component of LPS. Here we investigate the possible involvement of 3-deoxy-D-manno-octulosonic acid (Kdo), which is an additional component of the LPS in *C. trachomatis*. Highly motile preparations of sperm from normozoospermic patients were incubated for 6 hours with commercial sources of lipid A and Kdo. Conventional lipid A inhibitors, polymyxin B (PMB) and anti-CD14 monoclonal antibody (mAb) were used to test the ability of both lipid A and Kdo to induce an apoptotic-like response in mature sperm. Flow cytometry was used to determine apoptosis by the expression of annexin V. Caspase activity was also measured by

fluorometry and by the use of a pan-caspase inhibitor and caspase-3 inhibitor. Both lipid A and Kdo at 50 µg/mL caused significant mortality of sperm. However, although PMB and anti-CD14 mAb were inhibitory to the activity of lipid A on sperm, no such effect was seen against Kdo. In the presence of either lipid A or Kdo, sperm death was caused by an apoptotic-like effect that was caspase mediated. We conclude that Kdo shares its spermicidal properties with lipid A and seems to kill sperm in a similar manner. These results provide an explanation for higher than expected levels of spermicidal activity of LPS that are not caused by lipid A.

Key words: Apoptosis, human sperm, Kdo.

J Androl 2006;27:774–779

Introduction

Lipopolysaccharide (LPS) is a heat-stable complex and unique glycolipid that is present in all Gram-negative bacteria. It is the immunodominant antigen of most Gram-negative bacteria and is considered to be intimately associated with the virulence of the pathogen (Volk et al, 1996). LPS consists of 3 structural sections: lipid A, core polysaccharide, and O antigen. Lipid A is responsible for the endotoxin activity of LPS and binds to polymyxin B (Morrison and Jacobs, 1976). The core oligosaccharide region has 2 parts. The inner core is characterized by more unusual sugars, particularly 3-deoxy-D-manno-2-octulosonic acid (Kdo) and heptose. The former of these is seen in almost every LPS looked at to date, being α -bound to the carbohydrate backbone of lipid A in every case. The outer core typically consists of common hexose sugars such as glucose, galactose, *N*-acetyl galactosamine, and *N*-acetyl glucosamine and is generally more variable than the inner core. Interestingly, while the O antigen and the majority of the core

can be removed in some viable mutants, the Kdo residue, like lipid A, is always required for bacterial viability (Erridge et al, 2002). The interaction between LPS and the effector cell occurs via the CD14 receptor, a phosphatidylinositol-anchored molecule, in the presence of a plasma protein, the LPS binding protein which functions as an opsonin (Tobias et al, 1986). After this stage, cell signaling proceeds via Toll-like receptors and the cell is activated, which in many cases is followed by cytokine release.

Chlamydia trachomatis LPS is known to be particularly spermicidal (Galdiero et al, 1994; Hosseinzadeh et al, 2003) and is composed of a simple structure containing lipid A and Kdo (Rund et al, 1999). Our previous studies showed the toxicity of chlamydial LPS on ejaculated human spermatozoa in vitro, although inhibition by polymyxin B (PMB) was incomplete (Hosseinzadeh et al, 2003). On the basis of these considerations, the purpose of the present study was to verify the probable effect of Kdo in triggering apoptosis-like death of spermatozoa. Although the Kdo we used in the following experiments was synthetic, our aim was to assess any biologic activity of LPS that might be attributed to Kdo rather than lipid A using the specific binding of PMB and inhibitory effect of an anti-CD14 monoclonal antibody (mAb) as well as caspase inhibitors.

Correspondence to: Dr Adrian Eley, Division of Genomic Medicine, The Medical School, The University of Sheffield, Sheffield, S10 2RX, United Kingdom (e-mail: a.r.eley@sheffield.ac.uk).

Received for publication March 3, 2006; accepted for publication June 21, 2006.

DOI: 10.2164/jandrol.106.000083

Materials and Methods

Sperm Preparation

Semen samples were obtained from the andrology laboratory (Jessop Wing, Royal Hallamshire Hospital, Sheffield, United Kingdom) from normozoospermic patients by masturbation. Ethical approval for the use of semen samples in this study was granted by the South Sheffield Research Ethics Committee (project number 02/337). The ejaculates had a high sperm concentration ($>60 \times 10^6$ sperm/mL), with more than 15% ideal morphologic forms according to World Health Organization (1999) criteria. From each ejaculate, fast, progressively motile spermatozoa were purified and washed by a Percoll gradient technique (Moohan and Lindsay, 1995), and the samples which yielded more than 90% motile spermatozoa were used and adjusted to 5×10^6 /mL in Earle balanced salt solution (EBSS; Sigma Chemical Co, Poole, United Kingdom) containing 0.3% (w/v) bovine serum albumin.

Assessment of Sperm Viability

Sperm viability was determined using the hypo-osmotic swelling (HOS) test (Jeyendran et al, 1984). Briefly 20 μ L of each incubate was transferred to 200 μ L of HOS solution (tri-sodium citrate and fructose) and incubated at 37°C for 30 minutes. 10 μ L of each incubate was then spotted onto a microscope slide, and spermatozoa was classified as live or dead according to the method of Jeyendran and coworkers (1984).

Preparation of Control Populations

In the following experiments, 1 mM staurosporine-treated sperm and untreated Percoll-prepared sperm were used as positive and negative apoptosis controls, respectively. A necrosis control was produced by heat treatment of spermatozoa at 56°C for 1 hour.

Endotoxin Activity of the Kdo Preparation

The absence of endotoxin activity in the Kdo preparation was checked using a limulus amoebocyte lysate assay (Cambrex Biosciences, Wokingham, United Kingdom). The method was performed as recommended by the manufacturer.

The Effect of Lipid A and Kdo on Sperm

This experiment was conducted to assess the relative spermicidal activities of the 2 major components of bacterial LPS. Semen samples were obtained from 6 men and prepared as described above to give 4×200 - μ L aliquots of each sample. To 3 aliquots were added 50 μ g/mL of lipid A (Sigma), 50 μ g/mL of Kdo (Sigma), and a combination of both. To the remaining aliquot, 20 μ L of EBSS was added as a control. All were incubated for 6 hours at 37°C in 5% CO₂ and examined for sperm mortality.

The Effect of PMB and Anti-CD14 mAb on the Spermicidal Activity of Lipid A and Kdo

The LPS inhibitors PMB and anti-CD14 mAb, which work against lipid A, were used in experiments to confirm the ability

of lipid A and Kdo to kill spermatozoa. Semen samples were obtained from a further 6 men and prepared as described above to give 9×200 - μ L aliquots of each sample. PMB (Sigma) was used at a final concentration of 100 μ g/mL as previously reported (Hosseinzadeh et al, 2003); anti-CD14 mAb (BD PharMingen, Oxford, United Kingdom), which was free of endotoxin and sodium azide, was used at a final concentration of 10 μ g/mL. Both were incubated with spermatozoa for 30 minutes prior to adding lipid A and Kdo to the spermatozoa as described above.

Evaluation of the Apoptotic Effect of Kdo and Lipid A on Spermatozoa

Annexin V binding and use of a pan-caspase inhibitor allowed further insights into whether spermicidal activities caused by Kdo and lipid A are due to an apoptotic-like effect. Preparations of 5×10^6 sperm/mL from 6 men were incubated with either 50 μ g/mL of lipid A or Kdo at 37°C in 5% CO₂ for 6 hours. In another set of experiments, before the above was added, sperm were incubated with 200 μ M of pan-caspase inhibitor (Z-VAD-FMK; Calbiochem, Darmstadt, Germany) for 1 hour at room temperature. To prepare the samples for flow cytometry, the cells were then washed twice with PBS and resuspended in $1 \times$ annexin binding buffer (BD PharMingen) at a final concentration of 1×10^6 sperm/mL. An aliquot of 100 μ L of the solution (1×10^5 cells) was then transferred to a 12×75 -mm polystyrene test tube (Elkay, Basingstoke, United Kingdom) followed by the addition of 5 μ L of annexin V-FITC and 50 μ g/mL propidium iodide (PI; Sigma). The cells were then mixed by vortexing gently and incubated for 15 minutes at room temperature in the dark. An aliquot of 400 μ L of $1 \times$ annexin binding buffer was added to each tube, and the cells were analyzed by flow cytometry in a FACSCalibur (BD Biosciences, Oxford, United Kingdom). A minimum of 10 000 spermatozoa were examined for each test. The sperm population was gated by using forward-angle light scatter. The FITC-labeled annexin V-positive sperm cells were measured in the FL1 channel, and the PI-labeled cells were measured in the FL2 channel of the flow cytometer, as described by Eley and colleagues (2005a).

Quantification of Caspase Activity in Spermatozoa Incubated with Lipid A or Kdo

Confirmatory evidence of an apoptotic-like effect on sperm caused by Kdo and lipid A was obtained by measurement of sperm caspase activities (as described by Weng and coworkers, 2002) and the use of a caspase-3 inhibitor. Sperm lysates were prepared as described by Weng et al (2002). Briefly, 1×10^7 sperm (after a 6-hour incubation with either 50 μ g/mL lipid A or Kdo) were centrifuged to form a pellet. In another set of experiments, before incubation with lipid A, Kdo, or staurosporine, sperm were incubated with 200 μ M caspase-3 inhibitor I (Calbiochem) for 1 hour at room temperature. The pellet was then placed on dry ice for 5 to 10 minutes prior to lysis in 400 μ L of 10 mM Tris buffer with 0.5% Triton X-100 (Sigma), pH 7.5, containing general protease inhibitors 2 mM

phenylmethylsulfonyl fluoride, 10 $\mu\text{g}/\text{mL}$ leupeptin, 10 $\mu\text{g}/\text{mL}$ pepstatin, 10 mM dithiothreitol (Sigma) for 15 minutes at 4°C. The lysates were vortexed and sonicated before centrifugation (20 minutes at 4°C) to remove particulate matter. The supernatant was then removed, and an aliquot of sperm lysate (100 μg , according to the Bradford method [1976]) was diluted with 1 mL of solution containing interleukin 1 β -converting enzyme buffer (50 mM HEPES, 10% sucrose, 0.1% CHAPS [3-([3-chloamidopropyl]dimethylammonio)-1-propanesulfonate], pH 7.5) and 50 μM of fluorogenic substrate Ac-DEVD-afc (Bachem, Merseyside, United Kingdom) and incubated for 45 minutes at 37°C. The fluorescent emission (excitation 400 nm and emission 505 nm) was measured in a fluorometer (Perkin Elmer, Beaconsfield, United Kingdom). Background fluorescence was determined using blanks without sperm. Standards containing 1 to 5 μM AFC-(7-amino-4-trifluoromethyl coumarin) were used to determine the amount of fluorochrome released, and apoptotic human neutrophils treated with 1 mmol/L cycloheximide were used as a positive control.

Statistical Analysis

We used the Student's *t* test and 1-way analysis of variance (ANOVA) test as indicated in the figure legends. A *P* value of less than .001 was considered significant.

Analysis was by GraphPad Instat software version 3.0 (GraphPad Software Inc, San Diego, Calif).

Results

The Effect of Lipid A and Kdo on Sperm

Immediately after density gradient centrifugation, only 5.4% of spermatozoa in the control incubation were dead. Over the 6-hour incubation period this value did not change significantly in the control population ($P > .05$). However, the mortality of the spermatozoa that were incubated with lipid A, Kdo, and a combination of both (50 $\mu\text{g}/\text{mL}$) for 6 hours increased up to 23.6%, 20.6%, and 21%, respectively (Figure 1). These values were significantly different from the 6-hour control with a sperm mortality of only 8.8% ($P < .001$). Endotoxin activity in the Kdo preparation was shown to be less than 0.01 ng/mL.

The Effect of PMB and Anti-CD14 mAb on Lipid A and Kdo Acting Against Sperm

After a 6-hour incubation anti-CD14 mAb (8.0%) and PMB (9.8%) had no significant effect on spermatozoa death above that seen in the control population (approximately 10.2%) ($P > .05$). However, the spermicidal activity of lipid A and Kdo was significantly higher than that seen in control incubations ($P > .001$) (Figure 2). Spermatozoa in the presence of a combina-




Figure 1. The effects of lipid A, 3-deoxy-D-manno-octulosonic acid (Kdo), and the combination of lipid A and Kdo (all at 50 $\mu\text{g}/\text{mL}$) on human spermatozoa. The parameter measured is sperm mortality. The data shown are the mean of 6 experiments + SEM. *** indicates a significant difference from the control incubation using the Student's *t* test ($P < .001$).

tion of lipid A and anti-CD14 mAb (12.6%) and lipid A and PMB (9.6%) showed significantly fewer dead spermatozoa ($P < .001$). However, anti-CD14 mAb (17.6%) and PMB (17.8%) were not able to reduce the effect of Kdo on sperm ($P > .05$) (Figure 2).

Evaluation of the Apoptotic-like Effect of Kdo and Lipid A on Spermatozoa

Both lipid A and Kdo induced an apoptotic-like effect in spermatozoa (Figure 3), which was significantly inhibited by the pan-caspase inhibitor Z-VAD-FMK ($P < .001$). This suggested that the lipid A and Kdo-induced apoptotic-like effect might be caspase mediated.

Quantification of Caspase Activity in Spermatozoa Stimulated With Lipid A or Kdo

High levels of caspase activity were seen in cycloheximide-treated human neutrophils which were obtained from 6 men and gave a mean level of 129 nM/min mg (results not shown). Spermatozoa treated with staurosporine, lipid A, or Kdo showed significantly higher levels of caspase activity compared with unstimulated controls which were significantly reduced in the presence of caspase-3 inhibitor ($P < .001$), confirming that the apoptotic-like effect in these spermatozoa was caspase mediated (Figure 4).




Figure 2. The effect of anti-CD14 monoclonal antibody (mAb) (10 μ g/mL) and polymyxin B (PMB) (100 μ g/mL) on lipid A and 3-deoxy-D-manno-octulosonic acid (Kdo) (both at 50 μ g/mL) acting against human spermatozoa. Spermatozoa were incubated with lipid A or Kdo with and without anti-CD14 mAb and PMB for 6 hours at 37°C. The data shown are the mean of 6 experiments \pm SEM. *** indicates significant differences of $P < .001$ for lipid A incubated with anti-CD14 mAb and PMB. Additions to Kdo were not significant (NS) at $P > .05$. + indicates a significant difference of $P < .001$ for additions of lipid A and Kdo when compared with the control incubation using the Student's t test.

Discussion

In this study we have shown that Kdo appears to be as spermicidal as lipid A, and this may explain why PMB incompletely inhibits *C. trachomatis* LPS-induced sperm death in ejaculated human spermatozoa (Hosseinzadeh et al, 2003). In *C. trachomatis* LPS, the core region consists of the trisaccharide $Kdo_{\alpha} \rightarrow 8 Kdo_2 \rightarrow 4 Kdo$, emphasizing the importance of Kdo in this bacterium (Rund et al, 1999).

Although anti-CD14 mAb was able to inhibit the ability of lipid A to kill sperm, this was not seen with Kdo. One possible reason for this is that the α configuration of Kdo (as in the LPS molecule) may be required for CD14 binding (Cavaillon et al, 1996) but that this tautomer was not present in its synthetic form.

Although there is no additive spermicidal effect of lipid A and Kdo, this is not surprising given the findings of dose-response experiments using chlamydial LPS in which significant increases in LPS are required to see marked differences in spermicidal activity (Hosseinzadeh et al, 2003).

On further investigation of the mode of sperm death following exposure to either lipid A or Kdo, it appeared

that both induced an apoptotic-like effect (as observed through the expression of annexin V) and that caspases were activated as shown by inhibition with both the pan-caspase and caspase-3 inhibitors. These findings support the role of caspases in ejaculated sperm apoptosis as described by Weng and colleagues (2002) and Eley and coworkers (2005a).

While it has been suggested that the exposure of phosphatidylserine may occur in sperm as part of the changes that take place in the plasma membrane as a result of capacitation (de Vries et al, 2003) in a series of previous experiments (Eley et al, 2005a), those were shown to be caspase mediated; therefore, we are confident that the results described here are associated with apoptosis rather than capacitation. Recently our findings have been confirmed by Satta and collaborators (2006) who described *C. trachomatis* producing apoptosis (including expression of annexin V) in human sperm.

The spermicidal activity of Kdo was unexpected. Lack of anti-CD14 inhibition of Kdo to kill sperm suggested that a different method of cell signaling was occurring in comparison to that seen with lipid A. However, the extent and mechanisms of causation of sperm death seemed to be similar.




Figure 3. Percent spermatozoa expressing annexin V in the presence and absence of pan-caspase inhibitor (PCI) when exposed to the following over a 6-hour incubation period: control (no addition), H/S (heat-shock), S (1 mM staurosporine), 50 μ g/mL Kdo, and 50 μ g/mL lipid A. Data shown are the mean \pm SEM of incubations with sperm preparations from 6 men. *** indicates a significant difference ($P < .001$) between incubations with and without PCI using the 1-way analysis of variance test.




Figure 4. Caspase activity of spermatozoa in the presence and absence of caspase-3 inhibitor (C3I) when exposed to the following over a 6-hour incubation period: control (no addition), H/S (heat shock), S (1 mM staurosporine), 50 μ g/mL Kdo, and 50 μ g/mL lipid A. Data shown are the mean \pm SEM of incubations with sperm preparations from 6 men. *** indicates a significant difference ($P < .001$) between incubations with and without C3I with the 1-way analysis of variance test.

Recently we have hypothesized (Eley et al, 2005b) that as chlamydial LPS is the principal factor leading to sperm apoptosis following co-incubation with *C. trachomatis*, a new line of inquiry would be to measure the levels of LPS in semen and relate these to parameters of semen quality. However, now that we have shown comparative levels of spermicidal activity for Kdo as well as lipid A, measurement of LPS in semen (which only detects the lipid A portion and not Kdo) would underestimate the total spermicidal activity and is an important factor for consideration in such studies of LPS detection. In possible situations in which the lipid A may be sequestered away from semen, any remaining Kdo present would be undetectable by routine LPS detection methods and yet could be responsible for spermicidal activity, thereby complicating any Gram-negative bacterial association with semen quality.

Because of our surprising findings of the spermicidal activity of synthetic Kdo, they need to be confirmed using Kdo extracted from chlamydial LPS. However, there are difficulties in growing enough chlamydia in tissue culture to purify Kdo from sufficient LPS. We also believe that further work should be performed on

Kdo component sugars to investigate those chemical structures that are responsible for sperm death and to assess their spermicide potential.

Acknowledgments

We would like to thank Dr S Hosseinzadeh for help and advice. We also thank the Ministry of Health and Medical Education and Rafsanjan University of Medical Sciences, Iran, for financial support.

References

- Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilising the principle of protein-dye binding. *Anal Biochem.* 1976;72:248–254.
- Cavaillon J-M, Marie C, Caroff M, Ledur A, Godard I, Pulain D, Fitting C, Haeffner-Cavaillon N. CD14/LPS receptor exhibits lectin-like properties. *J Endotoxin Res.* 1996;3:471–480.
- de Vries KJ, Wiedmer T, Sims PJ, Gadella BM. Caspase-independent exposure of aminophospholipids and tyrosine phosphorylation in bicarbonate responsive human sperm cells. *Biol Reprod.* 2003;68:2122–2134.
- Eley A, Hosseinzadeh S, Hakimi H, Geary I, Pacey AA. Apoptosis of ejaculated human sperm is induced by co-incubation with *Chlamydia trachomatis* lipopolysaccharide. *Hum Reprod.* 2005a;20:2601–2607.
- Eley A, Pacey AA, Galdiero M, Galdiero M, Galdiero F. Can *Chlamydia trachomatis* directly damage your sperm? *Lancet Infect Dis.* 2005b;5:53–57.
- Erridge C, Bennett-Guerrero F, Poxton IR. Structure and function of lipopolysaccharides. *Microbes Infect.* 2002;4:837–851.
- Galdiero F, Sommese L, Gorga F, Galdiero E, Rizzo A, Ajello M. Toxic effect on human spermatozoa by *Chlamydia trachomatis* purified lipopolysaccharide. *FEMS Microbiol Lett.* 1994;115:197–200.
- Hosseinzadeh S, Pacey AA, Eley A. *Chlamydia trachomatis*-induced death of human spermatozoa is caused primarily by lipopolysaccharide. *J Med Microbiol.* 2003;52:193–200.
- Jeyendran RS, Van der Ven HH, Perez-Pelaez M, Crabo BG, Zaneveld LJD. Development of an assay to assess the functional integrity of the human sperm membrane and its relationship to the other semen characteristics. *J Reprod Fertil.* 1984;70:219–228.
- Mooohan JM, Lindsay KS. Spermatozoa selected by a discontinuous Percoll density gradient exhibit better motion characteristics, more hyperactivation, and longer survival than direct swim-up. *Fertil Steril.* 1995;64:160–165.
- Morrison DC, Jacobs DM. Binding of polymyxin B to the lipid A portion of bacterial lipopolysaccharides. *Immunochemistry.* 1976;13:813–818.
- Rund S, Lindner B, Brade H, Holst O. Structural analysis of the lipopolysaccharide from *Chlamydia trachomatis* serotype L2. *J Biol Chem.* 1999;274:16819–16824.
- Satta A, Stivala A, Garozzo A, Morello A, Perdichizzi A, Vicari E, Salmeri M, Calogero AE. Experimental *Chlamydia trachomatis* infection causes apoptosis in human sperm. *Hum Reprod.* 2006;21:134–137.
- Tobias PS, Soldau K, Ulevitch RJ. Isolation of a lipopolysaccharide-binding acute phase reactant from rabbit serum. *J Exp Med.* 1986;164:777–793.
- Volk WA, Gebhardt BM, Hammarskjold M-L. Bacterial cell structures. In Volk WA, Gebhardt BM, Hammarskjold M-L, Kadner RJ, eds. *Essentials of Medical Microbiology.* 5th ed. Philadelphia, Pa: Lippincott-Raven Publishers; 1996;240.
- Weng SL, Taylor SL, Morshedi M, Schuffner A, Duran EH, Beebe S, Oehninger S. Caspase activity and apoptotic markers in ejaculated human sperm. *Mol Hum Reprod.* 2002;8:984–991.
- World Health Organization. *WHO Laboratory Manual for the Examination of Human Semen and Semen–Cervical Mucus Interaction.* Cambridge: Cambridge University Press; 1999.