

# Calcium-Modulated Rod Outer Segment Membrane Guanylate Cyclase Type 1 Transduction Machinery in the Testes

ANNA JANKOWSKA,\* BEATA BURCZYNSKA,\* TERESA DUDA,† JERZY B. WARCHOL,\* AND RAMESHWAR K. SHARMA†

From the \*Department of Cell Biology, University of Medical Sciences, Poznan, Poland; and the †Unit of Regulatory and Molecular Biology, Departments of Cell Biology and Ophthalmology, SOM & NJMS, University of Medicine and Dentistry of New Jersey, Stratford, New Jersey.

**ABSTRACT:** The importance of the second messengers, Ca<sup>2+</sup> and cyclic GMP, for the process of fertilization is well established; the mechanisms for their intracellular regulations in the testes are, however, poorly understood. This study documents the biochemical, molecular, and functional identity of a Ca<sup>2+</sup>-modulated membrane guanylate cyclase transduction machinery in bovine testes. The machinery is both inhibited and stimulated by free Ca<sup>2+</sup> levels. The Ca<sup>2+</sup>-sensor component of the inhibitory mode of the machinery is GCAP1 (guanylate cyclase activating protein type 1) and for the

stimulatory mode is S100B. The transduction component is a Ca<sup>2+</sup>-driven rod outer segment membrane guanylate cyclase type 1, ROS-GC1. The cyclase is predominantly expressed in spermatogenic cells. GCAP1 expression is restricted to a small population of spermatogonia, whereas S100B is present in the majority of spermatocytes and spermatids. The expression of GCAP1 and S100B in spermatocytes and spermatids is mutually exclusive.

Key words: ROS-GC1, GCAP1, S100B, signal transduction.

**J Androl 2007;28:50–58**

Intracellular second messenger cyclic guanosine monophosphate (cGMP) is involved in the regulation of a broad spectrum of physiological processes ranging from peripheral to the central nervous system. In testes, it has been implicated to affect motility in spermatozoa, development of testicular germ cells, relaxation of peritubular lamina propria cells, testosterone synthesis in Leydig cells, dilatation of testicular blood vessels, and fertilization (Ong et al, 1975; Rossi et al, 1985; Mukhopadhyay et al, 1986; Pandey et al, 1986; Armstrong et al, 1994; Sanchez et al, 1996; Wiesner et al, 1998; Middendorff et al, 2000; Wayman et al, 2005). Guanylate cyclase-activating substances, which are secreted by several tissues in the genital tract, affect sperm motility, capacitation, and acrosomal reactivity; stimulation of sperm metabolism; and promotion of sperm ability to approach the oocyte, interact with it, and finally, to fertilize it (Garbers, 1989; Shapiro et al, 1990; Armstrong et al, 1994; reviewed in: Revelli et al, 2002; Kawase et al, 2004).

In the process of fertilization Ca<sup>2+</sup> plays a key role. The mechanisms controlling Ca<sup>2+</sup> entry into the sperm,

Ca<sup>2+</sup> conductance, and ultimately the spermatozoa movement toward the oocyte, remain unknown but are thought to involve various types of Ca<sup>2+</sup> channels (reviewed in: Benoff, 1998; Publicover and Barratt, 1999; Son et al, 2000; Jagannathan et al, 2002a,b; Castellano et al, 2003; Chiarella et al, 2004; Stamboulian et al, 2004; Trevino et al, 2004). The acrosomal activity of human spermatozoa is thought to be triggered by Ca<sup>2+</sup> influx mainly through  $\alpha$ 1H T-type Ca<sup>2+</sup> channels, expressed exclusively in spermatogenic cells (Son et al, 2000; Jagannathan et al, 2002a,b). A novel Ca<sup>2+</sup> channel, CatSper, present in the sperm has been cloned (Nikpoor et al, 2004). This gene encodes a unique and testes-specific Ca<sup>2+</sup> channel. A significant reduction in the level of CatSper gene expression has been observed among patients with lowered sperm motility (Nikpoor et al, 2004). Also a cyclic nucleotide-gated channel has been identified in the mammalian sperm (Weyand et al, 1994; Wiesner et al, 1998). The homooligomeric  $\alpha$  subunit cloned from bovine testes is roughly 200-fold more sensitive to cyclic GMP than to cyclic AMP, suggesting that the channel is specific and is a part of the cyclic GMP signaling pathway (Wiesner et al, 1998).

Cyclic GMP is synthesized by a group of enzymes termed guanylate cyclases. Depending on their cellular distribution, these enzymes have been classified into 2 families, membrane-bound and soluble. The soluble form is heterodimeric in structure and is stimulated by

Supported by The Polish Committee for Scientific Research award 3PO5E 11526 (AJ), NIH awards DC 005349 (RKS), and HL 070015 (TD).

Correspondence to: Anna Jankowska, ul. Swieckiego 6, 60-781 Poznan, Poland (e-mail: ajanko@amp.edu.pl).

Received for publication April 6, 2006; accepted for publication August 16, 2006.

DOI: 10.2164/jandrol.106.000182

nitric oxide and carbon dioxide (Murad, 1994; Middendorff et al, 1997; Pyriochou and Papapetropoulos, 2005). The membrane-bound form is a single transmembrane-spanning protein; it is monomeric, yet requires homodimerization for its activity (Chinkers and Wilson, 1992; Vaandrager et al, 1994; Wilson and Chinkers, 1995; Labrecque et al, 1999; Yu et al, 1999). On a biochemical basis, the membrane guanylate cyclase family is divided into 2 subfamilies, surface receptors and rod outer segment guanylate cyclase (ROS-GC). A central feature of the surface receptor subfamily is that all its members are receptors for peptide hormones or bacterial toxins (Fulle and Garbers, 1994; Leitman et al, 1994; Amin et al, 1996; Sharma and Duda, 1997; reviewed in: Sharma, 2002; Kuhn, 2003). These receptors, by binding their ligands, transduce the signal into production of the second messenger cyclic GMP.

The  $\text{Ca}^{2+}$ -modulated membrane guanylate cyclase subfamily has been named ROS-GC. ROS-GC subfamily is designed to transduce  $\text{Ca}^{2+}$  signals (reviewed in: Koch et al, 2002; Sharma, 2002; Sharma and Duda, 2006). ROS-GC machinery is a 2-component transduction system, the enzyme ROS-GC and the  $\text{Ca}^{2+}$  sensor. It is expressed specifically in the sensory and intermediary neurons of the retina (Hayashi and Yamazaki, 1991; Dizhoor et al, 1994; Goracznik et al, 1994; Lowe et al, 1995; Goracznik et al, 1997), pinealocytes (Venkataraman et al, 2000), the olfactory bulb (Duda et al, 2001a,b), and the anterior portion of the gustatory epithelium (Duda and Sharma, 2004). There are 3 members of this subfamily, ROS-GC1, ROS-GC2, and ONE-GC, expressed specifically in the olfactory neuroepithelium (Duda et al, 2001a,b). These cyclases, in contrast to the peptide hormone receptor guanylate cyclases, receive signals indirectly through  $\text{Ca}^{2+}$  sensor proteins, which belong to 2 classes, guanylyl cyclase activating proteins (GCAPs) and calcium dependent-GCAPs (CD-GCAPs). GCAPs receive the signals, undergo conformational change and inhibit ROS-GC. CD-GCAPs receive the signals, also undergo conformational change, and stimulate ROS-GC (reviewed in Koch et al, 2002; Sharma, 2002). GCAPs and CD-GCAPs sense  $\text{Ca}^{2+}$  signals through their EF-hand domains. There are 3 known functional GCAPs, 1, 2, and 3, and 2 CD-GCAPs, S100B and neurocalcin  $\delta$  (Palczewski et al, 1994; reviewed in Haeseleer et al, 1999; Sharma, 2002; Sharma and Duda, 2006).

The presence and functional activity of the soluble and particulate (ANF-RGC and CNP-RGC) guanylate cyclase has been shown in testes (Marala and Sharma, 1988; Middendorff et al, 1996; Middendorff et al, 1997; Middendorff et al, 2000; Muller et al, 2004). In the sperm of marine invertebrates, the presence of chemotactic peptide receptors guanylate cyclases has also been shown

(Ramarao and Garbers, 1985; Matsumoto et al, 2003), but there is no evidence of their presence in the mammalian sperm. Until now, the presence of  $\text{Ca}^{2+}$ -modulated ROS-GC transduction system has been thought to be the exclusive domain of sensory or sensory-linked neuronal cells. This study documents the biochemical, molecular, and functional identity of a  $\text{Ca}^{2+}$ -modulated membrane guanylate cyclase transduction machinery in bovine testes. The machinery is both inhibited and stimulated by free  $\text{Ca}^{2+}$  levels. The  $\text{Ca}^{2+}$ -sensor component of the inhibitory mode of the machinery is GCAP1 and for the stimulatory mode is S100B. The transduction component is the guanylate cyclase ROS-GC1.

## Materials and Methods

### Reagents

GCAP1 and GCAP2 were cloned, expressed, and purified as described previously (Venkataraman et al, 2000). S100B used was obtained commercially (Sigma Chemical Co, St Louis, Mo) or recombinant, prepared in our laboratory as described previously (Duda and Sharma, 2004).

### Antibodies

Characterization of highly specific antibodies raised against GCAP1, GCAP2, S100B, and ROS-GC1 has been described previously (Duda et al, 1998; Venkataraman et al, 2000; Duda et al, 2001a,b; Duda and Sharma, 2004). The antibodies were enriched by precipitating the immunoglobulin fraction using ammonium sulphate. ELISA and Western blotting were used to test the specificity and to determine the titer of purified antibodies.

### Reverse Transcription Polymerase Chain Reaction

Bovine testes were purchased from a local slaughterhouse. The tunica albuginea was removed, and total RNA was isolated from the fraction containing seminiferous tubules and interstitial tissue using TriPure isolation reagent (Roche Diagnostics, Mannheim, Germany) according to the manufacturer's protocols. The cDNA library was constructed using Advantage RT for PCR kit (BD-Biosciences, San Jose, Calif) and used for the amplification of the 548-bp fragment of ROS-GC1, 373 bp of ROS-GC2, 627 bp of GCAP1, 601 bp of GCAP2, and 279 bp of S100B coding region. The amplified fragments were purified on agarose gel and sequenced to confirm their identities. Additionally, as a control, the 238-bp specific fragment of a housekeeping gene, large ribosomal subunit (L30), was amplified from the cDNA library.

### Preparation of the Membrane Fraction of Testes

Membrane fraction of bovine testes was isolated according to the protocol described previously (Marala et al, 1991). The tissue fragments devoid of tunica albuginea, including seminiferous tubules and interstitial tissue, were homogenized in a buffer containing 250 mmol sucrose, 10 mmol Tris-HCl (pH 7.4), and

1 mmol phenylmethylsulfonyl fluoride and centrifuged at 150 *g* for 10 minutes to remove cell debris and nuclei. The post-mitochondrial supernatant (after centrifugation at 400 × *g* and 10 000 × *g*) was centrifuged at 40 000 × *g*. The pellet, designated as the membrane fraction, was suspended in the homogenization buffer and stored at −150°C until use.

### *Guanylate Cyclase Activity*

The membrane fraction (~0.1 μg protein/sample) was assayed for guanylate cyclase activity as described previously (Paul et al, 1987; Duda et al, 1998; Venkataraman et al, 2000; Duda and Sharma 2004). Briefly, membranes were preincubated on an ice-bath with or without regulatory proteins in the assay system containing 10 mmol theophylline, 15 mmol phosphocreatine, 20 μg creatine kinase, and 50 mmol Tris-HCl, pH 7.5, adjusted to appropriate free Ca<sup>2+</sup> concentrations with precalibrated Ca<sup>2+</sup>/EGTA solutions (Molecular Probes, Eugene, Ore). The total assay volume was 25 μL. The reaction was initiated by the addition of the substrate solution [4 mmol MgCl<sub>2</sub> and 1 mmol GTP (final concentrations)] and maintained by incubation at 37°C for 10 minutes. The reaction was terminated by the addition of 225 μL of 50 mmol sodium acetate buffer, pH 6.2, followed by heating in a boiling water bath for 3 minutes. The amount of cyclic GMP formed was determined by radioimmunoassay (Nambi et al, 1982). The RIA system detects 2 fmol of cyclic GMP/sample.

### *Western Blotting*

The procedure was carried out according to the previously published protocols (Venkataraman et al, 2000; Duda and Sharma, 2004). Briefly, after boiling in gel-loading buffer (62.5 mmol Tris-HCl [pH 7.5], 2% SDS, 5% glycerol, 1 mmol β-mercaptoethanol, and 0.005% bromophenol blue) ~150 μg of membrane protein was subjected to SDS-PAGE in a buffer containing 0.025 mmol Tris-HCl (pH 8.3), 0.192 mol glycine, and 0.1% SDS. The resolved proteins were transferred to nitrocellulose membranes, and the blot was incubated in Tris-buffered saline containing 0.05% Tween 20 (TBS-T), 5% powdered nonfat Carnation milk (blocking buffer) overnight at 4°C. The anti-ROS-GC1, GCAP1, GCAP2, or S100B rabbit polyclonal antibodies were added individually at dilution 1:1500, 1:1000, 1:1000, and 1:800, respectively. After 1 hour incubation the blot was rinsed with TBS-T and the incubation was continued with the secondary antibodies conjugated to horseradish peroxidase (1:10 000 dilution) in the blocking buffer for another hour. Finally the blot was treated with SuperSignal blaze chemiluminescent substrate (Pierce Biotechnology, Rockford, Ill) for 5 minutes according to the manufacturer's protocol. The immunoreactive band was detected by exposing the blot to Kodak X-ray film for 15 seconds. Images of the membranes with the immunoreactive bands were acquired by scanning and processed using Photoshop 6.0 software.

### *Immunohistochemistry*

Paraffin sections of bovine testes fixed in 4% paraformaldehyde were used for immunohistochemical detecting of ROS-

GC1, GCAP1, and S100B. Antigens were retrieved by microwave activation in citrate buffer (10 mmol, pH 6.0). After being blocked in PBS containing 3% bovine serum albumin and 0.1% Tween 20, sections were incubated with primary antibodies against ROS-GC1 diluted 1:200, GCAP1 diluted 1:100, and S100B diluted 1:50 in the same solution for 60 minutes at 37°C in a humidified chamber and washed for 60 minutes in PBS containing 0.1% Tween 20. AP-conjugated anti-rabbit IgG, diluted 1:200 (Sigma-Aldrich, Saint Louis, Mo) and NBT/BCIP as the substrate were used for detection. Incubation and washing conditions were as described for primary antibodies. Control included detection reactions carried out under identical conditions, except that the primary antibodies were replaced by nonimmune serum.

## **Results and Discussion**

With the background information that both cyclic GMP and Ca<sup>2+</sup> play an important role in the spermatogenesis and fertilization processes, the present study was aimed at determining if the activities of these 2 messengers are interlocked in the testes. Through molecular, biochemical, and functional approaches, this investigation shows that they are indeed linked and provides a mechanism for the interlocking process.

### *Testes Contain a Functional Ca<sup>2+</sup>-Dependent Membrane Guanylate Cyclase*

Membrane fraction of bovine testes was isolated and tested for guanylate cyclase activity. It was 3.5 pmol of cyclic GMP/mg protein/min. This is a cumulative value representing all putative guanylate cyclases present in the fraction. To assess whether the activity is Ca<sup>2+</sup>-modulated, the membrane fraction was exposed to increasing concentrations of free Ca<sup>2+</sup> and the guanylate cyclase activity was measured. The results (Figure 1A) show that the fraction contained a guanylate cyclase, which was either stimulated or inhibited depending on the free Ca<sup>2+</sup> concentration. A range of Ca<sup>2+</sup> concentration from 10 nmol to 0.8 μmol caused a dose-dependent decrease, and the higher range caused stimulation of the membrane guanylate cyclase. The inhibitory IC<sub>50</sub> value for Ca<sup>2+</sup> was ~300 nmol, and the stimulatory EC<sub>50</sub> value was ~5 μmol (Figure 1A). These results show that the testes contain membrane guanylate cyclase, which is both inhibited and stimulated by Ca<sup>2+</sup> signals.

### *Testes Express a Guanylate Cyclase With the Biochemical Attributes of ROS-GC1*

The 3 known Ca<sup>2+</sup>-dependent membrane guanylate cyclase forms, ROS-GC1, ROS-GC2, and ONE-GC, can be differentiated biochemically (Koch, 1991; Gor-




Figure 1. Ca<sup>2+</sup> regulation of guanylate cyclase activity in bovine testes. Membrane fraction of bovine testes was prepared and assayed for guanylate cyclase activity, as described in "Materials and Methods." (A) Response to Ca<sup>2+</sup>. Membranes were incubated in the presence of indicated concentration of Ca<sup>2+</sup>. (B) Response to GCAP1 and GCAP2. Membranes were incubated with increasing concentrations of GCAP1 or GCAP2 in the presence of 10 nmol free Ca<sup>2+</sup> and 4 μmol GCAP1 or 10 μmol GCAP2 in the presence of 100 μmol free Ca<sup>2+</sup>. (C) Response to S100B. Membranes were incubated with indicated concentrations of S100B in the presence of 100 μmol Ca<sup>2+</sup> and 8 μmol S100B in the presence of 10 nmol Ca<sup>2+</sup>. The molecular mass of 24 000 daltons was used to calculate the concentration of GCAP1 and GCAP2; 20 kd for S100B. Each

aczniak et al, 1998; Duda et al, 2001a,b). The ROS-GCs are stimulated in low nanomolar Ca<sup>2+</sup> concentrations and inhibited in the semimicro- to micromolar Ca<sup>2+</sup> concentrations by the GCAPs: ROS-GC1 by GCAP1 and GCAP2, and ROS-GC2 only by GCAP2 (reviewed in Palczewski et al, 1994; Haeseleer et al, 1999; Koch et al, 2002; Duda et al, 2005; Duda and Sharma, 2006). In the semimicro- to micro-molar Ca<sup>2+</sup> concentrations, both ROS-GCs are stimulated by the CD-GCAP, S100B and ROS-GC1 also by neurocalcin δ (reviewed in Koch et al, 2002; Sharma and Duda, 2006). ONE-GC is stimulated only by Ca<sup>2+</sup>-bound neurocalcin δ (Duda et al, 2001a,b; Sharma and Duda, 2006).

To determine which 1 (or more) of these cyclases is expressed in the testes, the membrane fraction of bovine testes was exposed to GCAP1, GCAP2, or S100B in the presence of 10 nmol and 100 μmol Ca<sup>2+</sup>. The stimulation by GCAP3 was not tested, since its expression was not detected in bovine tissue (Haeseleer et al, 1999). In 10 nmol Ca<sup>2+</sup> both GCAPs stimulated the cyclase activity in a dose-dependent manner, with an EC<sub>50</sub> of ~1 μmol for GCAP1 and of ~6 μmol for GCAP2 (Figure 1B). The maximal stimulation of the cyclase was ~2-fold above the basal value (Figure 1B). There was no stimulation by either GCAP in 100 μmol Ca<sup>2+</sup> (Figure 1B). The Ca<sup>2+</sup>-bound S100B also stimulated the cyclase activity in a dose-dependent fashion (Figure 1C). The half-maximal stimulation was achieved at 0.8 μmol S100B, and the maximal stimulation was ~2.5-fold. Ca<sup>2+</sup>-free S100B did not stimulate the cyclase activity (Figure 1C). All these stimulatory profiles are in accord with those established earlier for the reconstituted systems consisting exclusively of recombinant ROS-GC1 and individual Ca<sup>2+</sup> sensor proteins, GCAP1, GCAP2, or S100B (Duda et al, 1996a,b; Duda et al, 1998; Goraczniak et al, 1998; Krishnan et al, 1998). The ability of the cyclase to respond to both GCAPs (ROS-GC2 does not respond to GCAP1) and S100B indicates that the membrane fraction of bovine testes contains a guanylate cyclase that functionally mimics ROS-GC1 (Goraczniak et al, 1998; Haeseleer et al, 1999). It is noted that the ability of exogenous GCAPs or S100B (Figure 1B and C) to stimulate ROS-GC1 in the testes membrane fraction beyond the point achieved through only the removal or addition of Ca<sup>2+</sup> (Figure 1A) indicates that during the membrane preparation some Ca<sup>2+</sup> sensor proteins were lost; hence, their levels were lower than that necessary for maximal activation of the cyclase.

← experiment was done in triplicate and repeated 2 times. The results shown are from 1 representative experiment. Error bars are within the size of the symbols.

### The Transcript of ROS-GC1 Is Present in the Testes

To verify at molecular level the presence of ROS-GCs in bovine testes, total RNA was isolated from the tissue, reverse transcribed, and a 548-bp fragment corresponding to ROS-GC1 nucleotides 2042–2589 (GenBank accession number P55203) and a 373-bp fragment matching to ROS-GC2 nucleotides 3276–3649 (GenBank accession number U95958) were amplified. Amplification of a 238-bp fragment of the 30 kd ribosomal subunit (L30) served as a control. In the cases of ROS-GC1 and L30, the amplification yielded single bands of the predicted size (Figure 2A, lane 3, ROS-GC1; lane 4, L30). Sequencing of the amplified ROS-GC1 fragment confirmed its identity with bovine ROS-G1. The amplification of the ROS-GC2 fragment yielded no results (Figure 2B, lane 2); however, when cDNA from bovine retina was used as a control, the amplification yielded a single band of the predicted size of 373 bp (Figure 2B, lane 3). It is thus concluded that the ROS-GC1 but not ROS-GC2 transcript is present in the bovine testes.

### Biochemical Identification of ROS-GC1 in the Membranes of Bovine Testes

To determine the direct biochemical presence of ROS-GC1 in testes, Western blot analysis was carried out. The particulate fraction was electrophoresed, transferred onto nitrocellulose membrane, and probed with specific antibodies against ROS-GC1. Specificity of the antibody has been described previously (Venkataraman et al, 2000; Duda and Sharma, 2004). Membranes of COS cells expressing ROS-GC1 as well as membranes isolated from bovine retina were processed in parallel as a positive control. A single immunoreactive band of apparent mobility of ~116 kd was observed in control membranes (Figure 2C, lane 1 and 2). A band of identical mobility was detected in the membrane preparation from bovine testes (Figure 2C, lane 3). Identical results were obtained when antibodies against the catalytic domain of membrane guanylate cyclase corresponding to ROS-GC1 amino acid residues M872-S1016 (Fik-Rymarkiewicz et al, 2006) were used in Western blots (data not shown). Thus through 3 independent criteria, functional, molecular, and biochemical, it is concluded that the sole ROS-GC expressed in the membrane fraction of bovine testes is ROS-GC1.

### Expression of GCAP1 and S100B in Bovine Testes

To determine whether GCAP1 and/or GCAP2 are responsible for the cyclase stimulation in the absence of  $\text{Ca}^{2+}$  (Figure 1A; 10 nmol  $\text{Ca}^{2+}$ ) and if S100B is responsible for the  $\text{Ca}^{2+}$ -dependent stimulation (Figure 1A; 0.8–100  $\mu\text{mol Ca}^{2+}$ ), their presence was investigated at both the RNA and protein levels. Using




Figure 2. Expression of ROS-GC1 in bovine testes. **(A)** Detection of the ROS-GC1 transcript. RNA was isolated from the bovine testes and reverse transcribed. From the resulting cDNA a 548-bp fragment of ROS-GC1 (lane 3) and a 238-bp fragment of L30 (lane 4) were amplified. The predicted size fragment of ROS-GC1 corresponded to PCR product obtained from retinal cDNA (lane 2). **(B)** Detection of the ROS-GC2 transcript. ROS-GC2 was not detected in bovine testes (lane 2). When cDNA from bovine retina was used as a control, the amplification yielded a single band of the predicted size, 373 bp (lane 3). Molecular size markers are given in lane 1. **(C)** Detection of the ROS-GC1 protein. Membranes of COS cells expressing ROS-GC1 (100  $\mu\text{g}$  of total proteins; lane 1), retinal membranes (150  $\mu\text{g}$  of total proteins; lane 2) and the membranes of bovine testes (150  $\mu\text{g}$  of proteins; lane 3) were separated on a SDS-6% PAGE, transferred to nitrocellulose membrane, and probed with anti ROS-GC1 antibody, as described in "Materials and Methods." The immunoreactive bands were visualized by enhanced chemiluminescence (ECL). The ROS-GC1 immunoreactive band of the testes is indicated by an arrow. The positions of the molecular size markers are given alongside.

the specific primers designed based on the known sequences of S100B (GenBank accession number DQ195377), GCAP1 (GenBank accession number X95352), and GCAP2 (GenBank accession number U32856), RT-PCR was performed. A 279-bp fragment corresponding to the total coding region of S100B and a 627-bp fragment corresponding to the total coding region of GCAP1 were amplified (Figure 3A, lanes 2 and 3). Sequencing of the amplified fragments gave exact match to bovine GCAP1 and S100B cDNAs. There was, however, no amplification of a GCAP2

fragment (Figure 3A, lane 4). A single band of the predicted for GCAP2 size, 601 bp, was obtained in parallel control reaction with cDNA from bovine retina (Figure 3A, lane 5).

At the protein level, the identities of GCAPs and S100B in the membranes of bovine testes were established by Western blot analysis using specific antibodies against these proteins (Figure 3B). Anti-GCAP1 antibody detected a single band of apparent mobility of ~20 kd, identical to that of purified recombinant GCAP1 and retinal membranes (Figure 3B, panel "GCAP1&GCAP2," lanes 1, 2, and 3 for recombinant, retinal, and testes, respectively). In agreement with the cDNA amplification results, in testes membrane fraction there was no immunoreactivity with anti-GCAP2 antibody (Figure 3B, panel "GCAP1&GCAP2," lane 6). Anti-S100B antibody showed the presence of 2 immunoreactive bands (Figure 3B, panel "S100B," lane 3). Bands of identical mobility were observed for recombinant S100B and membranes isolated from retina (Figure 3B, panel "S100B," lanes 1 and 2). The mobility of these bands, ~10 and 20 kd, corresponds to monomeric and dimeric forms of S100B. It is therefore concluded that GCAP1, but not GCAP2, and S100B are expressed in bovine testes.

#### *ROS-GC1 Is Expressed in Germinal Cells of Bovine Testes*

To determine the localization of the components of the ROS-GC1 transduction system, immunohistochemical analyses were carried out with specific antibodies against ROS-GC1, GCAP1, and S100B. In all specimens, the labeling was randomly distributed in seminiferous tubules. The results are presented in Figure 4. Staining for ROS-GC1 was observed predominantly in spermatogenic cells, especially in primary spermatocytes and spermatids. However, single positively stained spermatogonia were observed as well (Figure 4A). The increase in staining was detected across the cells in different stages of the spermatogenesis process. The highest accumulation visible in long spermatids appears to correlate with the stage of the seminiferous cycle. The majority of spermatocytes and spermatids were stained for S100B (Figure 4B). A different pattern of immunostaining was observed with anti-GCAP1 antibody. Only a small population of spermatogonia was stained for GCAP1 (Figure 4C). It can be suggested that the expression of GCAP1 in these cells can be associated with their biological activity. No labeling was observed in the controls, where the primary antibodies were omitted (Figure 4D).

These results suggest that the ROS-GC1/GCAP1 and ROS-GC1/S100B transduction systems are mutually




Figure 3. Expression of ROS-GC1 modulators in the bovine testes. **(A)** Detection of GCAP1, GCAP2, and S100B transcripts. From a cDNA library constructed from bovine testes, a 279-bp coding region of S100B (lane 2) and a 627-bp fragment of GCAP1 (lane 3) were amplified. There was no amplification of GCAP2 fragment (lane 4). A single band of 601 bp characteristic for GCAP2 was obtained when cDNA from bovine retina was used as a control (lane 5). The products were resolved on agarose gel and visualized with ethidium bromide under UV light. Molecular size markers are in lane 1. **(B)** Detection of GCAP1, GCAP2, and S100B proteins. Membrane fraction from bovine testes was isolated and subjected to Western blot analyses with specific antibodies against GCAP1 (panel "GCAP1&GCAP2," lane 3), GCAP2 (panel "GCAP1&GCAP2," lane 6), and S100B (panel "S100B," lane 3), as described in "Materials and Methods." Recombinant GCAP1 (1  $\mu$ g) and GCAP2 (0.5  $\mu$ g) (panel "GCAP1&GCAP2," lanes 1 and 4, respectively) as well as commercial S100B (0.3  $\mu$ g) (panel "S100B," lane 1) were used as controls. Additionally, retinal membranes were probed with GCAP1, GCAP2, and S100B antibodies as a positive control of the study (panel "GCAP1&GCAP2," lane 2 for GCAP1 and 5 for GCAP2; panel "S100B," lane 2). The band corresponding to GCAP1 as well as monomeric and dimeric forms of S100B are indicated by arrows. Molecular size markers are given alongside.




Figure 4. Immunolocalization of ROS-GC1 in the bovine testes. **(A)** Immunohistochemistry was performed using specific antibodies against ROS-GC1 diluted 1:100 on paraffin section of bovine testes. The ROS-GC1 was localized in spermatogenic cells. The positive staining was observed in spermatogonia (Sg), primary spermatocytes (Sp), and spermatids (Sd), as indicated by the arrows. Bar = 100  $\mu$ m. **(B)** When the section were subjected to analysis with anti-S100B antibodies, the majority of positively stained spermatocytes and spermatids were observed. **(C)** Incubation of paraffin section of bovine testes with antibodies against GCAP1 resulted in staining of only a small population of spermatogonia. **(D)** Control. The procedure was exactly the same as for detection of ROS-GC1, except that nonimmune serum was used instead of ROS-GC1 antibodies. Lack of staining is characteristic for all cells of spermatogenic cycle.

exclusive and are functional in different phases of the spermatogenic cycle.

#### *The Ca<sup>2+</sup> Modulated GCAP1/S100B - ROS-GC1 Signal Transduction Model*

The present study, through functional, molecular, and biochemical approaches, shows the presence of ROS-GC1 in the testes. The enzyme coexists with its 2 Ca<sup>2+</sup>-dependent modulators, GCAP1 and S100B. In response to localized changes in free calcium concentration, these proteins bind to the specific domains, residing in the intracellular region of the cyclase, and inhibit or stimulate ROS-GC1. In this manner Ca<sup>2+</sup> pulses precisely regulate the levels of cyclic GMP generated in 1 (or

more) of the testes cells. Given the presence of a cyclic GMP-gated channel in these cells, the possibility exists that the levels control the hyper- or depolarized state of those cells. This prediction opens up a new venue of research. And the finding demonstrates that the presence of the ROS-GC transduction machinery is not unique to the neurosensory and neurosensory-linked systems; it also exists in the endocrinal system of the testes.

#### **References**

- Amin J, Carretero OA, Ito S. Mechanisms of action of atrial natriuretic factor and C-type natriuretic peptide. *Hypertension*. 1996;27:684-687.

- Armstrong VL, Clulow J, Murdoch RN, Jones RC. Intracellular signal transduction mechanisms of rat epididymal spermatozoa and their relationship to motility and metabolism. *Mol Reprod Dev.* 1994;38:77–84.
- Benoff S. Voltage dependent calcium channels in mammalian spermatozoa. *Front Biosci.* 1998;3:1220–1240.
- Castellano LE, Trevino CL, Rodriguez D, Serrano CJ, Pacheco J, Tsutsumi V, Felix R, Darszon A. Transient receptor potential (TRPC) channels in human sperm: expression, cellular localization and involvement in the regulation of flagellar motility. *FEBS Lett.* 2003;541:569.
- Chiarella P, Puglisi R, Sorrentino V, Boitani C, Stefanini M. Ryanodine receptors are expressed and functionally active in mouse spermatogenic cells and their inhibition interferes with spermatogonial differentiation. *J Cell Sci.* 2004;117:4127–4134.
- Chinkers M, Wilson EM. Ligand-independent oligomerization of natriuretic peptide receptors. Identification of heteromeric receptors and a dominant negative mutant. *J Biol Chem.* 1992;267:18589–18597.
- Dizhoor AM, Lowe DG, Olshevskaya EV, Laura RP, Hurley JB. The human photoreceptor membrane guanylyl cyclase, RetGC, is present in outer segments and is regulated by calcium and a soluble activator. *Neuron.* 1994;2:1345–1352.
- Duda T, Goraczniak R, Pozdnyakov N, Sitaramayya A, Sharma RK. Differential activation of rod outer segment membrane guanylate cyclases, ROS-GC1 and ROS-GC2, by CD-GCAP and identification of the signaling domain. *Biochem Biophys Res Commun.* 1998;242:118–122.
- Duda T, Goraczniak R, Sharma RK. Molecular characterisation of S100A1-S100B protein in retina and its activation mechanism of bovine photoreceptor guanylate cyclase. *Biochemistry.* 1996a;35:6263–6266.
- Duda T, Goraczniak R, Surgucheva I, Rudnicka-Nawrot M, Gorczyca WA, Palczewski K, Sitaramayya A, Baehr W, Sharma RK. Calcium modulation of bovine photoreceptor guanylate cyclase. *Biochemistry.* 1996b;35:8478–8482.
- Duda T, Jankowska A, Venkataraman V, Nagele R, Sharma RK. A novel calcium-regulated guanylate cyclase transduction system in olfactory neuroepithelium. *Biochemistry.* 2001b;40:12067–12077.
- Duda T, Sharma RK. S100B-modulated Ca(2+)-dependent ROS-GC1 transduction machinery in the gustatory epithelium: a new mechanism in gustatory transduction. *FEBS Lett.* 2004;577:393–398.
- Duda T, Venkataraman V, Krishnan A, Nagele RG, Sharma RK. Negatively calcium-modulated membrane guanylate cyclase signaling system in the rat olfactory bulb. *Biochemistry.* 2001a;40:4654–4662.
- Duda T, Venkataraman V, Ravichandran S, Sharma RK. ATP-regulated module (ARM) of the atrial natriuretic factor receptor guanylate cyclase. *Peptides.* 2005;26:969–984.
- Fik-Rymarkiewicz E, Duda T, Sharma RK. Novel frequenin-modulated Ca<sup>2+</sup>-signaling membrane guanylate cyclase (ROS-GC) transduction pathway in bovine hippocampus. *Mol Cell Biochem.* May 2006 [Epub ahead of print].
- Fulle HJ, Garbers DL. Guanylyl cyclases: a family of receptor-linked enzymes. *Cell Biochem Funct.* 1994;12:157–165.
- Garbers DL. Molecular basis of signaling in the spermatozoon. *J Androl.* 1989;10:99–107.
- Goraczniak RM, Duda T, Sharma RK. Calcium modulated signalling site in type 2 rod outer segment guanylate cyclase (ROS-GC2). *Biochem Biophys Res Commun.* 1998;245:447–453.
- Goraczniak RM, Duda T, Sharma RK. Structural and functional characterization of a second subfamily member of the calcium-modulated bovine rod outer segment membrane guanylate cyclase, ROSGC2. *Biochem Biophys Res Commun.* 1997;234:666–670.
- Goraczniak RM, Duda T, Sitaramayya A, Sharma RK. Structural and functional characterization of the rod outer segment membrane guanylate cyclase. *Biochem J.* 1994;302:455–461.
- Haeseleer F, Sokal I, Li N, Pettenati M, Rao N, Bronson D, Wechter R, Baehr W, Palczewski K. Molecular characterization of a third member of the guanylyl cyclase-activating protein subfamily. *J Biol Chem.* 1999;274:6526–6535.
- Hayashi F, Yamazaki A. Polymorphism in purified guanylate cyclase from vertebrate rod photoreceptors. *Proc Natl Acad Sci U S A.* 1991;88:4746–4750.
- Jagannathan S, Publicover SJ, Barratt CL. Voltage-operated calcium channels in male germ cells. *Reproduction.* 2002a;123:203–215.
- Jagannathan S, Punt EL, Gu Y, Arnoult C, Sakkas D, Barratt CL, Publicover SJ. Identification and localisation of T-type voltage-operated calcium channel subunits in human male germ cells. Expression of multiple isoforms. *J Biol Chem.* 2002b;8:8449–8456.
- Kawase O, Ueno S, Minakata H, Hoshi M, Matsumoto M. Guanylyl cyclase and cGMP-specific phosphodiesterase participate in the acrosome reaction of starfish sperm. *Zygote.* 2004;12:345–355.
- Koch KW. Purification and identification of photoreceptor guanylate cyclase. *J Biol Chem.* 1991;266:8634–8637.
- Koch K-W, Duda T, Sharma RK. Photoreceptor specific guanylate cyclases in vertebrate phototransduction. *Mol Cell Biochem.* 2002;230:97–106.
- Krishnan A, Goraczniak RM, Duda T, Sharma RK. Third calcium-modulated rod outer segment membrane guanylate cyclase transduction mechanism. *Mol Cell Biochem.* 1998;178:251–259.
- Kuhn M. Structure, regulation, and function of mammalian membrane guanylyl cyclase receptors, with a focus on guanylyl cyclase-A. *Circ Res.* 2003;93:700–709.
- Labrecque J, Mc Nicoll N, Marquis M, De Lean A. A disulfide-bridged mutant of natriuretic peptide receptor-A displays constitutive activity. Role of receptor dimerization in signal transduction. *J Biol Chem.* 1999;274:9752–9759.
- Leitman DC, Waldman SA, Murad F. Regulation of particulate guanylate cyclase by natriuretic peptides and Escherichia coli heat-stable enterotoxin. *Adv Pharmacol.* 1994;26:67–86.
- Lowe DG, Dizhoor AM, Liu K, Gu Q, Spencer M, Laura R, Lu L, Hurley JB. Cloning and expression of a second photoreceptor-specific membrane retina guanylyl cyclase (RetGC), RetGC-2. *Proc Natl Acad Sci U S A.* 1995;92:5535–5539.
- Marala RB, Sharma RK. Characterization of atrial-natriuretic-factor-receptor-coupled membrane guanylate cyclase from rat and mouse testes. *Biochem J.* 1988;251:301–304.
- Marala RB, Sitaramayya A, Sharma RK. Dual regulation of atrial natriuretic factor-dependent guanylate cyclase by ATP. *FEBS Lett.* 1991;281:73–76.
- Matsumoto M, Solzin J, Helbig A, Hagen V, Ueno S, Kawase O, Maruyama Y, Ogiso M, Godde M, Minakata H, Kaupp UB, Hoshi M, Weyand I. A sperm-activating peptide controls a cGMP-signaling pathway in starfish sperm. *Dev Biol.* 2003;260:314–324.
- Middendorff R, Davidoff MS, Behrends S, Mewe M, Miethens A, Muller D. Multiple roles of the messenger molecule cGMP in testicular function. *Andrologia.* 2000;32:55–59.
- Middendorff R, Muller D, Paust HJ, Davidoff MS, Mukhopadhyay AK. Natriuretic peptides in the human testis: evidence for a potential role of C-type natriuretic peptide in Leydig cells. *J Clin Endocrinol Metab.* 1996;81:4324–4328.
- Middendorff R, Muller D, Wichers S, Holstein AF, Davidoff MS. Evidence for production and functional activity of nitric oxide in



- seminiferous tubules and blood vessels of the human testes. *J Clin Endocrinol Metab.* 1997;82:4154–4161.
- Mukhopadhyay AK, Schumacher M, Leidenberger FA. Steroidogenic effect of atrial natriuretic factor in isolated mouse Leydig cells is mediated by cyclic GMP. *Biochem J.* 1986;239:463–467.
- Muller D, Mukhopadhyay AK, Speth RC, Guidone G, Potthast R, Potter LR, Middendorff R. Spatiotemporal regulation of the two atrial natriuretic peptide receptors in testes. *Endocrinol.* 2004;145:1392–1401.
- Murad F. Regulation of cytosolic guanylyl cyclase by nitric oxide: the NO-cyclic GMP signal transduction system. *Adv Pharmacol.* 1994;26:19–33.
- Nambi P, Aiyar NV, Sharma RK. Adrenocorticotropin-dependent particulate guanylate cyclase in rat adrenal and adrenocortical carcinoma: comparison of its properties with soluble guanylate cyclase and its relationship with ACTH-induced steroidogenesis. *Arch Biochem Biophys.* 1982;217:638–646.
- Nikpoor P, Mowla SJ, Movahedin M, Ziaee SA, Tiraihi T. CatSper gene expression in postnatal development of mouse testes and in subfertile men with deficient sperm motility. *Hum Reprod.* 2004;19:124–128.
- Ong SH, Whitley TH, Stowe NW, Steiner AL. Immunohistochemical localization of 3': 5'-cyclic AMP and 3': 5'-cyclic GMP in rat liver, intestine, and testis. *Proc Natl Acad Sci U S A.* 1975;72:2022–2026.
- Palczewski K, Subbaraya I, Gorczyca WA, Helekar BS, Ruiz CC, Ohguro H, Huang J, Zhao X, Crabb JW, Johnson RS, Walsh KA, Gray-Keller MP, Detwiler PB, Baehr W. Molecular characterization of retinal photoreceptor guanylyl cyclase activating protein. *Neuron.* 1994;13:395–404.
- Pandey KN, Pavlou SN, Kovacs WJ, Inagami T. Atrial natriuretic factor regulates steroidogenic responsiveness and cyclic nucleotide levels in mouse Leydig cells in vitro. *Biochem Biophys Res Commun.* 1986;38:399–404.
- Paul AK, Marala RB, Jaiswal RK, Sharma RK. Coexistence of guanylate cyclase and atrial natriuretic factor receptor in a 180-kD protein. *Science.* 1987;235:1224–1226.
- Publicover SJ, Barratt CL. Voltage-operated Ca<sup>2+</sup> channels and the acrosome reaction: which channels are present and what do they do? *Hum Reprod.* 1999;14:873–879.
- Pyriochou A, Papapetropoulos A. Soluble guanylyl cyclase: more secrets revealed. *Cell Signal.* 2005;17:407–413.
- Ramarao CS, Garbers DL. Receptor-mediated regulation of guanylate cyclase activity in spermatozoa. *J Biol Chem.* 1985;260:8390–8396.
- Revelli A, Ghigo D, Moffa F, Massobrio M, Tur-Kaspa I. Guanylate cyclase activity and sperm function. *Endocr Rev.* 2002;23:484–494.
- Rossi P, Pezzotti R, Conti M, Geremia R. Cyclic nucleotide phosphodiesterases in somatic and germ cells of mouse seminiferous tubules. *J Reprod Fertil.* 1985;74:317–327.
- Sanchez M, Menendez L, Garcia de Boto MJ, Hidalgo A. Role of cyclic nucleotides in contraction induced by oxytocin in the testicular capsule of the rat in vitro. *Pharmacology.* 1996;53:296–301.
- Shapiro BM, Cook S, Quest AF, Oberdorf J, Wothe D J. Molecular mechanisms of sea-urchin sperm activation before fertilization. *Reprod Fertil Suppl.* 1990;42:3–8.
- Sharma RK. Evolution of the membrane guanylate cyclase transduction system. *Mol Cell Biochem.* 2002;230:3–30.
- Sharma RK, Duda T. Calcium sensor neurocalcin  $\delta$ -modulated ROS-GC transduction machinery in the retinal and olfactory neurons. *Calcium Binding Proteins.* 2006;1:7–11.
- Sharma RK, Duda T. Plasma membrane guanylate cyclase. A multi-module transduction system. *Adv Expl Med Biol.* 1997;407:271–279.
- Son WY, Lee JH, Han CT. Acrosome reaction of human spermatozoa is mainly mediated by through  $\alpha$ 1H T-type calcium channels. *Mol Hum Reprod.* 2000;6:893–897.
- Stambouliau S, Kim D, Shin HS, Ronjat M, De Waard M, Arnoult C. Biophysical and pharmacological characterization of spermatogenic T-type calcium current in mice lacking the CaV3.1 (alpha1G) calcium channel: CaV3.2 (alpha1H) is the main functional calcium channel in wild-type spermatogenic cells. *J Cell Physiol.* 2004;200:116–124.
- Trevino CL, Felix R, Castellano LE, Gutierrez C, Rodriguez D, Pacheco J, Lopez-Gonzalez I, Gomora JC, Tsutsumi V, Hernandez-Cruz A, Fiordelisio T, Scaling AL, Darshon A. Expression and differential cell distribution of low-threshold Ca(2+) channels in mammalian male germ cells and sperm. *FEBS Lett.* 2004;563:87–92.
- Vaandrager AB, van der Wiel E, Hom ML, Luthjens LH, de Jonge HR. Heat-stable enterotoxin receptor/guanylyl cyclase C is an oligomer consisting of functionally distinct subunits, which are non-covalently linked in the intestine. *J Biol Chem.* 1994;269:16409–16415.
- Venkataraman V, Duda T, Nagele R, Sharma RK. Rod outer segment membrane guanylate cyclase type I-linked stimulatory and inhibitory calcium signaling systems in the pineal gland: biochemical, molecular, and immunohistochemical evidence. *Biochemistry.* 2000;39:6042–6052.
- Wayman C, Phillips S, Lunny C, Webb T, Fawcett L, Baxendale R, Burgess G. Phosphodiesterase 11 (PDE11) regulation of spermatozoa physiology. *Int J Impot Res.* 2005;17:216–223.
- Weyand I, Godde M, Frings S, Weiner J, Muller F, Altenhofen W, Hatt H, Kaupp UB. Cloning and functional expression of a cyclic-nucleotide-gated channel from mammalian sperm. *Nature.* 1994;368:859–863.
- Wiesner B, Weiner J, Middendorff R, Hagen V, Kaupp UB, Weyand I. Cyclic nucleotide-gated channels on the flagellum control Ca<sup>2+</sup> entry into sperm. *J Cell Biol.* 1998;42:473–484.
- Wilson EM, Chinkers M. Identification of sequences mediating guanylyl cyclase dimerization. *Biochemistry.* 1995;34:4696–4701.
- Yu H, Olshevskaya E, Duda T, Seno K, Hayashi F, Sharma RK, Dizhoor AM, Yamazaki A. Activation of retinal guanylyl cyclase-I by Ca<sup>2+</sup>-binding proteins involves its dimerization. *J Biol Chem.* 1999;274:15547–15555.