

What is (not) wrong with scalar gravity?

Domenico Giulini
University of Freiburg
Department of Physics
Hermann-Herder-Strasse 3
D-79104 Freiburg, Germany

Abstract

On his way to General Relativity (GR) Einstein gave several arguments as to why a special relativistic theory of gravity based on a massless scalar field could be ruled out merely on grounds of theoretical considerations. We re-investigate his two main arguments, which relate to energy conservation and some form of the principle of the universality of free fall. We find that such a theory-based *a priori* abandonment not to be justified. Rather, the theory seems formally perfectly viable, though in clear contradiction with (later) experiments. This may be of interest to those who teach GR and/or have an active interest in its history.

Contents

1	Introduction	2
2	Historical background	2
3	Scalar gravity	5
3.1	First guesses and a naive theory	5
3.2	A consistent model-theory for scalar gravity	7
4	Free-fall in static homogeneous fields	10
4.1	The scalar model-theory	10
4.2	The naive scalar theory	12
4.3	Vector theory	12
4.4	Discussion	14
5	Periapsis precession	15
5.1	Scalar model-theory	17
5.2	Naive scalar theory	18
5.3	Vector theory	19
6	Energy conservation	21
6.1	Einstein's argument	21
6.2	Energy conservation in the scalar model-theory	23
6.3	Discussion	24
7	Conclusion	24

1 Introduction

General Relativity (henceforth ‘GR’) differs markedly in many structural aspects from all other theories of fundamental interactions, which are all formulated as Poincaré invariant theories in the framework of Special Relativity. A common strategy to motivate the particular structure of GR to those already familiar with Special Relativity and Poincaré invariant field theories is to first carefully consider the obstructions that prevent gravity from also fitting into this framework. A natural way to proceed is then to consider fields according to mass and spin (the Casimir operators of the Poincaré group that label its irreducible representations), discuss their possible equations, the inner consistency of the mathematical schemes so obtained, and finally their experimental consequences. Since gravity is a classical and long-ranged field one usually assumes right at the beginning the spin to be integral and the mass parameter to be zero. The first thing to consider would therefore be a massless scalar field. What goes wrong with such a theory?

There can be no doubt that scalar gravity is ruled out. However, especially if one has to teach the subject, one should be careful to give the *right* reasons for its abandonment. In particular, two logically different types of arguments should be strictly kept apart:

1. The theory is internally inconsistent. In a trivial sense this may mean that it is mathematically contradictory, in which case this is the end of the story. On a more sophisticated level it might also mean that the theory violates accepted fundamental physical principles, like e.g. that of energy conservation, without being plainly mathematically contradictory
2. The theory is formally consistent and in accord with basic physical principles, but simply refuted by experiments.

Note that, generically, it does not make much sense to claim both shortcomings simultaneously, since ‘predictions’ of inconsistent theories should not be trusted. The question to be addressed here is whether scalar gravity falls under the first category, i.e. whether it can be refuted on the basis of formal arguments alone.

Many people think that it can, following A. Einstein who accused scalar theories to

3. violate some form of the principle of universality of free fall,
4. violate energy conservation.

We will see that the actual situation is not that easy. We will proceed by the standard (Lagrangian) methods of modern field theory and take what we perceive as the obvious route when working from first principles.

2 Historical background

The abandonment of scalar theories of gravity by Einstein is intimately linked with the birth of General Relativity, in particular with his conviction that general covariance must replace the principle of relativity as used in Special Relativity.

I will focus on two historical sources in which Einstein complains about scalar gravity not being adequate. One is his joint paper with Marcel Grossmann on the ‘Entwurf Theory’ ([7], Vol. 4, Doc. 13), of which Grossmann wrote the ‘mathematical part’ and Einstein the ‘physical part’. Einstein finished with § 7, whose title asks: “Can the gravitational field be reduced to a scalar ?” (“Kann das Gravitationsfeld auf einen Skalar zurückgeführt werden ?”). In this paragraph he presented a Gedankenexperiment-based argument which allegedly shows that any Poincaré-invariant scalar theory of gravity, in which the scalar gravitational field couples exclusively to the matter via the trace of its energy-momentum tensor, necessarily violates energy conservation and is hence physically inconsistent. This he presented as plausibility argument why gravity has to be described by a more complex quantity, like the $g_{\mu\nu}$ of the ‘Entwurf Paper’, where he and Grossmann considers ‘generally covariant’ equations for the first time. After having presented his argument, he ends § 7 (and his contribution) with the following sentences, expressing his conviction in the validity of the principle of general covariance:

Einstein Quote 1. *Ich muß freilich zugeben, daß für mich das wirksamste Argument dafür, daß eine derartige Theorie [eine skalare Gravitationstheorie] zu verwerfen sei, auf der Überzeugung beruht, daß die Relativität nicht nur orthogonalen linearen Substitutionen gegenüber besteht, sondern einer viel weitere Substitutionsgruppe gegenüber. Aber wir sind schon deshalb nicht berechtigt, dieses Argument geltend zu machen, weil wir nicht imstande waren, die (allgemeinste) Substitutionsgruppe ausfindig zu machen, welche zu unseren Gravitationsgleichungen gehört.*¹ ([7], Vol. 4, Doc. 13, p. 323)

The other source where Einstein reports in more detail on his earlier experiences with scalar gravity is his manuscript entitled “Einiges über die Entstehung der Allgemeinen Relativitätstheorie”, dated June 20th 1933 (reprinted in [1]). There he describes in words (no formulae are given) how the ‘obvious’ special-relativistic generalization of the Poisson equation,

$$\Delta\Phi = 4\pi G\rho, \quad (1a)$$

together with a (slightly less obvious) special-relativistic generalization of the equation of motion,

$$\frac{d^2\vec{x}(t)}{dt^2} = -\vec{\nabla}\Phi(\vec{x}(t)), \quad (1b)$$

lead to a theory in which the vertical acceleration of a test particle in a static homogeneous vertical gravitational field depends on its initial horizontal velocity and also on its internal energy content. In his own words:

Einstein Quote 2. *Solche Untersuchungen führten aber zu einem Ergebnis, das mich in hohem Maß mißtrauisch machte. Gemäß der klassischen Mechanik ist nämlich*

¹ *To be sure, I have to admit that in my opinion the most effective argument for why such a theory [a scalar theory of gravity] has to be abandoned rests on the conviction that relativity holds with respect to a much wider group of substitutions than just the linear-orthogonal ones. However, we are not justified to push this argument since we were not able to determine the (most general) group of substitutions which belongs to our gravitational equations.*

die Vertikalbeschleunigung eines Körpers im vertikalen Schwerfeld von der Horizontal Komponente der Geschwindigkeit unabhängig. Hiermit hängt es zusammen, daß die Vertikalbeschleunigung eines mechanischen Systems bzw. dessen Schwerpunktes in einem solchen Schwerfeld unabhängig herauskommt von dessen innerer kinetischer Energie. Nach der von mir versuchten Theorie war aber die Unabhängigkeit der Fallbeschleunigung von der Horizontalgeschwindigkeit bzw. der inneren Energie eines Systems nicht vorhanden. Dies paßte nicht zu der alten Erfahrung, daß die Körper alle dieselbe Beschleunigung in einem Gravitationsfeld erfahren. Dieser Satz, der auch als Satz über die Gleichheit der trägen und schweren Masse formuliert werden kann, leuchtete mir nun in seiner tiefen Bedeutung ein. Ich wunderte mich im höchsten Grade über sein Bestehen und vermutete, daß in ihm der Schlüssel für ein tieferes Verständnis der Trägheit und Gravitation liegen müsse. An seiner strengen Gültigkeit habe ich auch ohne Kenntnis des Resultates der schönen Versuche von Eötvös, die mir – wenn ich mich richtig erinnere – erst später bekannt wurden, nicht ernsthaft gezweifelt. Nun verwarf ich den Versuch der oben angedeuteten Behandlung des Gravitationsproblems im Rahmen der speziellen Relativitätstheorie als inadäquat. Er wurde offenbar gerade der fundamentalsten Eigenschaft der Gravitation nicht gerecht. [...] Wichtig war zunächst nur die Erkenntnis, daß eine vernünftige Theorie der Gravitation nur von einer Erweiterung des Relativitätsprinzips zu erwarten war.² ([1], pp. 135–136)

Einstein belief, that scalar theories of gravity are ruled out, placed him—in this respect—in opposition to most of his contemporary physicist who took part in the search for a (special-) relativistic theory of gravity (Nordström, Abraham, Mie, von Laue ..). (Concerning Nordström's theory and the Einstein-Nordström interaction, see e.g. the beautiful discussions in [5] and [6].) Some of them were not convinced, it seems, by Einstein's inconsistency argument. For example, even after General Relativity was completed, Max von Laue wrote a comprehensive review paper on Nordström's theory, thereby at least implicitly claiming inner consistency [4]. Remarkably, this paper of Laue's is not contained in his collected writings.

On the other hand, modern commentators seem to fully accept Einstein's claims and view them as important step in the development of General Relativity and possibly also as an important step towards the requirement of general covariance, whose important heuristic power is unquestioned. Historically speaking this may certainly be true. But are these arguments also physically correct, so as to be appropriately repeated

² *These investigations, however, led to a result which raised my strong suspicion. According to classical mechanics, the vertical acceleration of a body in a vertical gravitational field is independent of the horizontal component of its velocity. Hence in such a gravitational field the vertical acceleration of a mechanical system, or of its center of gravity, comes out independently of its internal kinetic energy. But in the theory I advanced, the acceleration of a falling body was not independent of its horizontal velocity or the internal energy of the system. This did not fit with the old experience that all bodies experience the same acceleration in a gravitational field. This statement, which can be formulated as theorem on the equality of inertial and gravitational mass, became clear to me in all its deeper meaning. I wondered to the highest degree as to why it should hold and conjectured that it be the key for a deeper understanding of inertia and gravitation. I did not question its rigorous validity, even without knowing about the beautiful experiments by Eötvös, of which—if I remember correctly—I became aware only later. I now abandoned my attempt as inadequate to address the problem of gravitation along the lines outlined above. It obviously could not account for the most fundamental property of gravitation. [...] The important insight at this stage was that a reasonable theory of gravitation could only be expected from an extension of the principle of relativity.*

in, say, a modern course on GR? Unfortunately Einstein’s recollections do not give us sufficient insight into the precise mathematical formulations he had in mind when making the statements just quoted. But what one can do is writing down a plausible scalar theory and check whether its shortcomings are of the type Einstein mentions.

3 Scalar gravity

We wish to construct a Poincaré-invariant theory of a scalar gravitational field, Φ , coupled to matter. Before we will do so in a systematic manner, using Lagrangian methods, we will mention the obvious first and naive guesses for a Poincaré invariant generalization of formulae (1). Our conventions for the Minkowski metric are ‘mostly minus’, that is, $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$.

3.1 First guesses and a naive theory

There is an obvious way to generalize the left hand side of (1a), namely to replace the Laplace operator by minus (due to our ‘mostly minus’ convention) the d’Alembert operator:

$$\begin{aligned} \Delta &:= \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \\ \mapsto -\square &:= \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} = -\eta^{\mu\nu} \frac{\partial^2}{\partial x^\mu \partial x^\nu}. \end{aligned} \quad (2)$$

This is precisely what Einstein reported:

Einstein Quote 3. *Das einfachste war natürlich, das Laplacesche skalare Potential der Gravitation beizubehalten und die Poisson Gleichung durch ein nach der Zeit differenziertes Glied in naheliegender Weise so zu ergänzen, daß der speziellen Relativitätstheorie Genüge geleistet wurde.*³ ([1], p. 135)

At this stage Einstein gives no information (in his 1933 notes) how to replace the right hand side of (1a). Since the mass density ρ is not a scalar, it is clear that something must be done about it. Note that the rest-mass density is not a scalar either, since albeit the rest-mass is a scalar, the volume—as measured by the different inertial observers—is not. In Special Relativity the energy density is the 00 component of the energy-momentum tensor $T^{\mu\nu}$, which corresponds to a mass density T^{00}/c^2 . Hence a sensible generalization for the right hand side of (1a) is:

$$\rho \mapsto T/c^2 := \eta^{\mu\nu} T_{\mu\nu}/c^2, \quad (3)$$

so that (1a) translates to

$$\square\Phi = -\kappa T, \quad \text{where } \kappa := 4\pi G/c^2. \quad (4)$$

The next step is to generalize (1b). With respect to this problem Einstein remarks:

³ *The most simple thing to do was to retain the Laplacian scalar potential and to amend the Poisson equation by a term with time derivative, so as to comply with special relativity.*

Einstein Quote 4. *Auch mußte das Bewegungsgesetz des Massenpunktes im Gravitationsfeld der speziellen Relativitätstheorie angepaßt werden. Der Weg hierfür war weniger eindeutig vorgeschrieben, weil ja die Träge Masse eines Körpers vom Gravitationspotential abhängen konnte. Dies war sogar wegen des Satzes von der Trägheit der Energie zu erwarten.*⁴ ([1], p. 135)

It should be clear that the structurally obvious choice,⁵

$$\ddot{x}^\mu(\tau) = \eta^{\mu\nu} \nabla_\nu \Phi(x(\tau)), \quad (5)$$

cannot work. Four velocities are normed,

$$\eta(\dot{x}, \dot{x}) = \dot{x}_\mu \dot{x}^\mu = c^2, \quad (6)$$

so that

$$\eta(\dot{x}, \ddot{x}) = \dot{x}_\mu \ddot{x}^\mu = 0. \quad (7)$$

hence (5) implies the integrability condition $\dot{x}^\mu(\tau) \nabla_\mu \Phi(x(\tau)) = d\Phi(x(\tau))/d\tau = 0$. Hence (5) implies that Φ must stay constant along the worldline of the particle, with renders it physically totally useless. The reason for this failure lies in the fact that we replaced the three independent equations (1b) by four equations, which led to an overdetermination since the four-velocity still represents only three independent functions due to the kinematical constraint (6). More specifically, it is the component parallel to the four-velocity \dot{x} of the four-vector equation (5) that leads to the unwanted restriction. The obvious way out it to just retain the part of (5) perpendicular to \dot{x} :

$$\ddot{x}^\mu(\tau) = P^{\mu\nu}(\tau) \nabla_\nu \Phi(x(\tau)), \quad (8a)$$

where $P^{\mu\nu}(\tau) = \eta^{\nu\lambda} P_\lambda^\mu(\tau)$ and

$$P_\nu^\mu(\tau) := \delta_\nu^\mu - \dot{x}^\mu(\tau) \dot{x}_\nu(\tau) / c^2 \quad (8b)$$

is the one-parameter family of projectors orthogonal to the four-velocity $\dot{x}(\tau)$, for each point $x(\tau)$ of the particle's worldline. Hence, by construction, this modified equation of motion avoids the difficulty just mentioned. We will call the theory based on (4) and (8) the *naive theory*. We also note that (8) is equivalent to

$$\frac{d}{d\tau} (m(x(\tau)) \dot{x}^\mu(\tau)) = m(x(\tau)) \eta^{\mu\nu} \nabla_\nu \Phi(x(\tau)), \quad (9)$$

where m is a spacetime dependent mass, given by

$$m = m_0 \exp((\Phi - \Phi_0)/c^2). \quad (10)$$

Here m_0 is a constant, corresponding to the mass at gravitational potential Φ_0 (e.g. $\Phi_0 = 0$).

⁴ Also, the law of motion of a mass point in a gravitational field had to be adjusted to special relativity. Here the route was less uniquely mapped out, since the inertial mass of a body could depend on the gravitational potential. Indeed, this had to be expected on grounds of the law of inertia of energy.

⁵ Throughout we write ∇_μ for $\partial/\partial x^\mu$.

We could now work out consequences of this theory. However, before doing this, we would rather put the reasoning employed so far on a more systematic basis as provided by variational principles. This also allows us to discuss general matter couplings and check whether the matter coupling that the field equation (4) expresses is consistent with the coupling to the point particle, represented by the equation of motion (8). This has to be asked for if we wish to implement the equivalence principle in the following form:

Requirement 1 (Principle of universal coupling). All forms of matter (including test particles) couple to the gravitational field in a universal fashion.

We will see that in this respect the naive theory is not quite correct.

3.2 A consistent model-theory for scalar gravity

Let us now employ standard Lagrangian techniques to construct a Poincaré-invariant theory of a scalar gravitational field, Φ , coupled to matter. We take seriously the field equation (4) and seek an action which makes it the Euler-Lagrange equation. It is easy to guess⁶:

$$S_{\text{field}} + S_{\text{int}} = \frac{1}{\kappa c^3} \int d^4x \left(\frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi - \kappa \Phi T \right), \quad (11)$$

where S_{field} , given by the first term, is the action for the gravitational field and S_{int} , given by the second term, accounts for the interaction with matter.

To this we have to add the action S_{matter} for the matter, which we only specify insofar as we assume that the matter consists of a point particle of rest-mass m_0 and a ‘rest’ of matter that needs not be specified further for our purposes here. Hence $S_{\text{matter}} = S_{\text{particle}} + S_{\text{rom}}$ (rom = rest of matter) where

$$S_{\text{particle}} = -m_0 c^2 \int d\tau. \quad (12)$$

The quantity $d\tau = \frac{1}{c} \sqrt{\eta_{\mu\nu} dx^\mu dx^\nu}$ is the proper time along the worldline of the particle. We now invoke the principle of universal coupling to find the particle’s interaction with the gravitational field. It must be of the form ΦT_p , where T_p is the trace of the particle’s energy momentum tensor. The latter is given by

$$T_p^{\mu\nu}(x) = m_0 c \int \dot{x}^\mu(\tau) \dot{x}^\nu(\tau) \delta^{(4)}(x - x(\tau)) d\tau, \quad (13)$$

so that the particle’s contribution to the interaction term in (11) is

$$S_{\text{int-particle}} = -m_0 \int \Phi(x(\tau)) d\tau. \quad (14)$$

⁶ Note that Φ has the physical dimension of a squared velocity, κ that of length-over-mass. The prefactor $1/\kappa c^3$ gives the right hand side of (11) the physical dimension of an action. The overall signs are chosen according to the general scheme for Lagrangians: kinetic minus potential energy.

Hence the total action can be written in the following form:

$$\begin{aligned}
S_{\text{tot}} = & -m_0 c^2 \int (1 + \Phi(x(\tau))/c^2) d\tau \\
& + \frac{1}{\kappa c^3} \int d^4x \left(\frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi - \kappa \Phi T_{\text{rom}} \right) \\
& + S_{\text{rom}} .
\end{aligned} \tag{15}$$

By construction the field equation that follow from this action is (4), where the energy momentum-tensor refers to the matter without the test particle (the self-gravitational field of a *test* particle is always neglected). The equations of motion for the test particle are then given by

$$\ddot{x}^\mu(\tau) = P^{\mu\nu}(\tau) \partial_\nu \phi(x(\tau)) , \tag{16a}$$

$$\text{where } P^{\mu\nu}(\tau) = \eta^{\mu\nu} - \dot{x}^\mu(\tau) \dot{x}^\nu(\tau) / c^2 \tag{16b}$$

$$\text{and } \phi := c^2 \ln(1 + \Phi/c^2) . \tag{16c}$$

Three things are worth remarking at this point:

- The projector $P^{\mu\nu}$ now appears naturally.
- The difference between (8) and (16) is that in the latter it is ϕ rather than Φ that drives the four acceleration. This (only) difference to the naive theory was imposed upon us by the principle of universal coupling, which, as we have just seen, determined the motion of the test particle. This difference is small for small Φ/c^2 , since, according to (16c), $\phi \approx \Phi(1 + \Phi/c^2 + \dots)$. But it becomes essential if Φ gets close to $-c^2$, where ϕ diverges and the equations of motion become singular. We will see below that the existence of the critical value $\Phi = -c^2$ is more of a virtue than a deficiency and that it is the naive theory which displays an unexpected singular behavior.
- The universal coupling of the gravitational field to matter only involves the trace of energy-momentum tensor of the latter. As a consequence of the tracelessness of the pure electromagnetic energy-momentum tensor, there is no coupling of gravity to the *free* electromagnetic field, like e.g. a light wave in otherwise empty space. A travelling electromagnetic wave will not be influenced by gravitational fields. Hence this theory predicts no bending of light-rays that pass the neighborhoods of stars of other massive objects, in disagreement with experimental observations. Note however that the interaction of electromagnetic fields with other matter will change the trace of the energy-momentum tensor of the latter. For example, electromagnetic waves trapped in a material box with mirrored walls will induce additional stresses in the box's walls due to radiation pressure. This will increase the weight of the box corresponding to an additional mass $\Delta m = E_{\text{rad}}/c^2$, where E_{rad} is the energy of the radiation field. In this sense *bounded* electromagnetic fields *do* carry weight.

Let us now focus on the equations of motion specialized to static situations. That is, we assume that there exists some inertial coordinate system x^μ with respect to which Φ and hence ϕ are static, i.e. $\nabla_0 \Phi = \nabla_0 \phi = 0$. We have

Proposition 1. For static potentials (16) is equivalent to

$$\vec{x}''(t) = -(1 - \beta^2(t)) \vec{\nabla} \phi(\vec{x}(t)), \quad (17)$$

where here and below we write a prime for d/dt , and use the standard shorthands $\vec{v} = \vec{x}'$, $\vec{\beta} = \vec{v}/c$, $\beta = \|\vec{\beta}\|$, and $\gamma = 1/\sqrt{1 - \beta^2}$.

Proof. We write in the usual four-vector component notation: $\dot{x} = c\gamma(1, \vec{\beta})$. Using $d/d\tau = \gamma d/dt$ and $d\gamma/dt = \gamma^3(\vec{\alpha} \cdot \vec{v}/c^2)$, we have on one side

$$\ddot{x}^\mu = \gamma^4 (\vec{\alpha} \cdot \vec{\beta}, \vec{\alpha}_\parallel + \gamma^{-2} \vec{\alpha}_\perp), \quad (18a)$$

with $\vec{\alpha} := d\vec{v}/dt$ and where $\vec{\alpha}_\parallel := \beta^{-2} \vec{\beta} (\vec{\beta} \cdot \vec{\alpha})$ and $\vec{\alpha}_\perp := \vec{\alpha} - \vec{\alpha}_\parallel$ are respectively the spatial projections of $\vec{\alpha}$ parallel and perpendicular to the velocity \vec{v} . On the other hand we have

$$-\dot{x}^\mu \dot{x}^\nu \nabla_\nu \phi / c^2 = -\gamma^2 (\vec{\beta} \cdot \vec{\nabla} \phi) (1, \vec{\beta}) \quad (18b)$$

$$\eta^{\mu\nu} \nabla_\nu \phi = (0, -\vec{\nabla} \phi), \quad (18c)$$

so that

$$(\eta^{\mu\nu} - \dot{x}^\mu \dot{x}^\nu / c^2) \nabla_\nu \phi = -\gamma^2 (\beta \cdot \nabla \phi, \vec{\nabla}_\parallel \phi + \gamma^{-2} \vec{\nabla}_\perp \phi), \quad (18d)$$

where $\vec{\nabla}_\parallel := \beta^{-2} \vec{\beta} (\vec{\beta} \cdot \vec{\nabla})$ and $\vec{\nabla}_\perp := \vec{\nabla} - \vec{\nabla}_\parallel$ are the projections of the gradient parallel and perpendicular to \vec{v} respectively. Equating (18a) and (18d) results in

$$\vec{\alpha} \cdot \vec{\beta} = -\gamma^{-2} \vec{\beta} \cdot \vec{\nabla} \phi, \quad (18e)$$

$$\vec{\alpha} = -\gamma^{-2} \vec{\nabla} \phi. \quad (18f)$$

Since (18e) is trivially implied by (18f), (18f) alone is equivalent to (16) in the static case, as was to be shown. \square

Einstein's second quote suggests that he also arrived at an equation like (17), which clearly displays the dependence of the acceleration in the direction of the gravitational field on the transversal velocity. We will come back to this in the discussion section.

We can still reformulate (17) so as to look perfectly Newtonian:

Proposition 2. Let m be the rest-mass of the point particle. Then (17) implies

$$m\vec{\alpha} = -\vec{\nabla} \tilde{\phi}(\vec{x}(t)) \quad \text{with} \quad \tilde{\phi} := (mc^2/2) \gamma_0^{-2} \exp(2\phi/c^2), \quad (19)$$

where γ_0 is an integration constant.

Proof. Scalar multiplication of (17) with \vec{v} leads to

$$(\ln \gamma + \phi/c^2)' = 0, \quad (20)$$

which integrates to

$$\gamma = \gamma_0 \exp(-\phi/c^2), \quad (21)$$

where γ_0 is a constant. Using this equation to eliminate the γ^{-2} on the right hand side of (17) the latter assumes the form (19). \square

4 Free-fall in static homogeneous fields

4.1 The scalar model-theory

Suppose that with respect to some inertial reference frame with coordinates (ct, x, y, z) the gravitational potential ϕ just depends on z . Let a body be released at $x = y = z = 0$ with proper velocity $\dot{y}_0 = \dot{z}_0 = 0$, $\dot{x}_0 = c\beta\gamma$, and $\dot{t}_0 = \gamma$, where $c\beta := v := \dot{x}_0/\dot{t}_0$ is the ordinary velocity and $\gamma := 1/\sqrt{1 - \beta^2}$. Here an overdot denotes the derivative with respect to proper time and we used the fact that $c^2\dot{t}^2 - \dot{x}^2 - \dot{y}^2 - \dot{z}^2 = c^2$. We take the gravitational field to point into the negative z direction so that ϕ is a function of z with positive derivative ϕ' . Note that $\dot{z}(\phi' \circ z) = d(\phi \circ z)/d\tau$ for which we simply write $\dot{\phi}$ with the usual abuse of notion (i.e. taking ϕ to mean $\phi \circ z$). Finally, we normalize ϕ such that $\phi(z = 0) = 0$.

The equations of motion (16a) now simply read

$$\ddot{t} = -\dot{t}\dot{\phi}/c^2, \quad (22a)$$

$$\ddot{x} = -\dot{x}\dot{\phi}/c^2, \quad (22b)$$

$$\ddot{y} = -\dot{y}\dot{\phi}/c^2, \quad (22c)$$

$$\ddot{z} = -(1 + \dot{z}^2/c^2)\phi'. \quad (22d)$$

The first integrals of the first three equations, keeping in mind the initial conditions, are

$$(\dot{t}(\tau), \dot{x}(\tau), \dot{y}(\tau)) = (1, c\beta, 0) \gamma \exp(-\phi(z(\tau))/c^2). \quad (23)$$

Further integration requires the knowledge of $z(\tau)$, that is, the horizontal motion couples to the vertical one if expressed in proper time.⁷ Fortunately, the vertical motion does *not* likewise couple to the horizontal one, that is, the right hand side of (22d) just depends on $z(\tau)$. Writing it in the form

$$\frac{\ddot{z}/c^2}{1 + \dot{z}^2/c^2} = -\dot{\phi}/c^2 \quad (24)$$

its integral for $\dot{z}(\tau = 0) = 0$ and $\phi(z = h) = 0$ (so that $\phi(z < h) < 0$) is

$$\dot{z} = -c\sqrt{\exp(-2\phi/c^2) - 1}. \quad (25)$$

From this the eigentime τ_h for dropping from $z = 0$ to $z = -h$ with $h > 0$ follows by one further integration, showing already at this point its independence of the initial horizontal velocity.

Here we wish to be more explicit and solve the equations of motion for the one-parameter family of solutions to (4) for $T = 0$ and a Φ that just depends on z , namely $\Phi = gz$, for some constant g that has the physical dimension of an acceleration. As already announced we normalize Φ such that $\Phi(z = 0) = 0$. These solutions correspond to what one would call ‘homogeneous gravitational field’. But note that these solutions are *not* globally regular since $\phi = c^2 \ln(1 + \Phi/c^2) = c^2 \ln(1 + gz/c^2)$ exists only for $z > -c^2/g$ and it is the quantity ϕ rather than Φ that corresponds to the Newtonian potential (i.e. whose negative gradient gives the local acceleration).

⁷ In terms of coordinate time the horizontal motion decouples: $dx/dt = \dot{x}/\dot{t} = c\beta \Rightarrow x(t) = c\beta t$.

Upon insertion of $\Phi = gz$, (25) can be integrated to give $z(\tau)$. Likewise, from (25) and (23) we can form $dz/dt = \dot{z}/\dot{t}$ and $dz/dx = \dot{z}/\dot{x}$ which integrate to $z(t)$ and $z(x)$ respectively. The results are

$$z(\tau) = -\frac{c^2}{g} \left\{ 1 - \sqrt{1 - (\tau g/c)^2} \right\}, \quad (26a)$$

$$z(t) = -\frac{2c^2}{g} \sin^2(gt/2\gamma c), \quad (26b)$$

$$z(x) = -\frac{2c^2}{g} \sin^2(gx/2\beta\gamma c^2). \quad (26c)$$

For completeness we mention that direct integration of (23) gives for the other component functions, taking into account the initial conditions $t(0) = x(0) = y(0) = 0$:

$$(t(\tau), x(\tau), y(\tau)) = (1, c\beta, 0) (\gamma c/g) \sin^{-1}(g\tau/c). \quad (27)$$

The relation between τ and t is

$$\tau = (c/g) \sin(gt/\gamma c). \quad (28)$$

Inversion of (26a) and (26a) leads to the proper time, τ_h , and coordinate time, t_h , that it takes the body to drop from $z = 0$ to $z = -h$:

$$\tau_h = \frac{c}{g} \sqrt{1 - (1 - gh/c^2)^2} \approx \sqrt{2h/g}, \quad (29a)$$

$$t_h = \gamma \frac{2c}{g} \sin^{-1}\left(\sqrt{gh/2c^2}\right) \approx \gamma \sqrt{2h/g}. \quad (29b)$$

The approximations indicated by \approx refer to the leading order contributions for small values of gh/c^2 (and any value of γ). The appearance of γ in (29b) signifies the quadratic dependence on the initial horizontal velocity: the greater the inertial horizontal velocity, the longer the span in inertial time for dropping from $z = 0$ to $z = -h$. This seems to be Einstein's point (cf. Quote 2). In contrast, there is no such dependence in (29a), showing the independence of the span in *eigentime* from the initial horizontal velocity.

The eigentime for dropping into the singularity at $z = -h = -c^2/g$ is $\tau_* = c/g$. In particular, it is finite, so that a freely falling observer experiences the singularity of the gravitational field $-\vec{\nabla}\phi$ in finite proper time. We note that this singularity is also present in the static spherically symmetric vacuum solution $\Phi(r) = -Gm/r$ to (4), for which $\phi(r) = c^2 \ln(1 + \Phi/c^2)$ exists only for $\Phi > c^2$, i.e. $r > Gm/c^2$. The Newtonian acceleration diverges as r approaches this value from above, which means that stars of radius smaller than that critical value cannot exist because no internal pressure can support the infinite inward pointing gravitational pull.

Knowing General Relativity, this type of behavior does not seem too surprising after all. Note that we are here dealing with a non-linear theory, since the field equations (4) become non-linear if expressed in terms of ϕ according to (16c).

4.2 The naive scalar theory

Let us for the moment return to the naive theory, given by (4) and (8). Its equations of motion in a static and homogeneous vertical field are obtained from (22) by setting $\phi = gz$. Insertion into (25) leads to $z(\tau)$. The expressions $z(t)$ and $z(x)$ are best determined directly by integrating $dz/dt = \dot{z}/\dot{t}$ using (25) and (27). One obtains

$$z(\tau) = \frac{c^2}{g} \ln\left(\cos(g\tau/c)\right), \quad (30a)$$

$$z(t) = -\frac{c^2}{g} \ln\left(\cosh(gt/\gamma c)\right), \quad (30b)$$

$$z(x) = -\frac{c^2}{g} \ln\left(\cosh(gx/\beta\gamma c^2)\right). \quad (30c)$$

The proper time and coordinate time for dropping from $z = 0$ to $z = -h$ are therefore given by

$$\tau_h = \frac{c}{g} \cos^{-1}\left(\exp(-hg/c^2)\right) \approx \sqrt{2h/g}, \quad (31a)$$

$$t_h = \frac{c}{g} \gamma \cosh^{-1}\left(\exp(hg/c^2)\right) \approx \gamma \sqrt{2h/g}, \quad (31b)$$

where \approx gives again the leading order contributions for small gh/c^2 .⁸ The general relation between τ and t is obtained by inserting (30a) into the expression (23) for \dot{t} and integration:

$$\tau = \frac{2c}{g} \left\{ \tan^{-1}\left(\exp(gt/\gamma c)\right) - \pi/4 \right\}. \quad (32)$$

The surprising feature of (31a) is that τ_h stays finite for $h \rightarrow \infty$. In fact, $\tau_\infty = c\pi/2g$. So even though the solution $\phi(z)$ is globally regular, the solution to the equations of motion is in a certain sense not, since the particle reaches the ‘end of spacetime’ in finite proper time. This should presumably be seen as a worse singular behavior than that discussed before, since it is not associated with any singular behavior of the field itself. Except perhaps for the fact that the very notion of an infinitely extended homogeneous field is itself regarded as unphysical.

4.3 Vector theory

For comparison it is instructive to look at the corresponding problem in a vector (spin 1) theory, which we here do not wish to discuss in detail. It is essentially given by Maxwell’s equations with appropriate sign changes to account for the attractivity of like ‘charges’ (here masses). This causes problems, like that of runaway solutions, due to the possibility to radiate away negative energy. But the problem of free fall in a homogeneous gravitoelectric field can be addressed, which is formally identical to that of free fall of a charge e and mass m in a static and homogeneous electric field $\vec{E} = -E\vec{e}_z$. So let us first look at the electrodynamical problem.

The equations of motion (the Lorentz force law) are

$$m\dot{z}^\mu = e\eta^{\mu\nu}F_{\nu\lambda}\dot{z}^\lambda, \quad (33)$$

⁸ To see this here use e.g. the identity $\cos^{-1}(x) = \tan^{-1}(\sqrt{x^2 - 1})$.

where $F_{03} = -F_{30} = -E/c$ and all other components vanish. Hence, writing

$$\mathcal{E} := eE/mc, \quad (34)$$

we have

$$c\ddot{t} = -\mathcal{E}z, \quad (35a)$$

$$\ddot{x} = 0, \quad (35b)$$

$$\ddot{y} = 0, \quad (35c)$$

$$\ddot{z} = -\mathcal{E}c\dot{t}. \quad (35d)$$

With the same initial conditions as in the scalar case we immediately have

$$x(\tau) = c\beta\gamma\tau, \quad y(\tau) = 0. \quad (36)$$

(35a) and (35d) are equivalent to

$$(ct \pm z)'' = \mp\mathcal{E}(ct \pm z)', \quad (37)$$

which twice integrated lead to

$$ct(\tau) \pm z(\tau) = A_{\pm} \exp(\mp\mathcal{E}\tau) + B_{\pm}, \quad (38)$$

where A_+ , A_- , B_+ , and B_- are four constants of integration. They are determined by $z(0) = \dot{z}(0) = t(\tau) = 0$ and $c\dot{t}^2 - \dot{x}^2 - \dot{y}^2 - \dot{z}^2 = c^2$, leading to

$$t(\tau) = (\gamma/\mathcal{E}) \sinh(\mathcal{E}\tau) \quad (39)$$

and also

$$z(\tau) = - (2c\gamma/\mathcal{E}) \sinh^2(\mathcal{E}\tau/2). \quad (40a)$$

Using (39) and (36) to eliminate τ in favour of t or x respectively in (40a) gives

$$z(t) = -\frac{\gamma c}{\mathcal{E}} \left(\sqrt{1 + (t\mathcal{E}/\gamma)^2} - 1 \right), \quad (40b)$$

$$z(x) = -\frac{2\gamma c}{\mathcal{E}} \sinh^2(x\mathcal{E}/2\beta\gamma c). \quad (40c)$$

Inverting (40a) and (40b) gives the expressions for the spans of eigentime and inertial time respectively, that it takes for the body to drop from $z = 0$ to $z = -h$:

$$\tau_h = (2/\mathcal{E}) \sinh^{-1} \left(\sqrt{\mathcal{E}h/2\gamma c} \right) \approx \gamma^{-1/2} \sqrt{2h/\mathcal{E}c}, \quad (41a)$$

$$t_h = (\gamma/\mathcal{E}) \sqrt{(1 + \mathcal{E}h/\gamma c)^2 - 1} \approx \gamma^{+1/2} \sqrt{2h/\mathcal{E}c}. \quad (41b)$$

This is the full solution to our problem in electrodynamics, of which we basically just used the Lorentz force law. It is literally the same in a vector theory of gravity, we just have to keep in mind that the ‘charge’ e is now interpreted as gravitational mass, which is to be set equal to the inertial mass m , so that $e/m = 1$. Then $\mathcal{E}c$ becomes equal to the ‘gravitoelectric’ field strength E , which directly corresponds to the strength g of the scalar gravitational field. Having said this, we can directly compare (41) with (29). For small field strength we see that in both cases t_h is larger by a factor of γ than τ_h , which just reflects ordinary time dilation. However, unlike in the scalar case, the eigentime span τ_h also depends on γ in the vector case. The independence of τ_h on the initial horizontal velocity is therefore a special feature of the scalar theory.

4.4 Discussion

Let us reconsider Einstein’s statements in Quote 2, in which he dismisses scalar gravity for it predicting an unwanted dependence on the vertical acceleration on the initial horizontal velocity. As already noted, we do not know exactly in which formal context Einstein derived this result (i.e. what the “von mir versuchten Theorie” mentioned in Quote 2 actually was), but it seems most likely that he arrived at an equation like (17), which clearly displays the alleged behavior. In any case, the diminishing effect of horizontal velocity onto vertical acceleration is at most of *quadratic* order in v/c .

Remark 1. How could Einstein be so convinced that such an effect did not exist? Certainly there were no experiments at the time to support this. And yet he asserted that such a prediction “did not fit with the *old experience* [my italics] that all bodies experience the same acceleration in a gravitational field” (cf. Quote 2). What was it based on?

One way to rephrase/interpret Einstein’s requirement is this: the time it takes for a body in free fall to drop from a height h to the ground should be independent of its initial horizontal velocity. More precisely, if you drop two otherwise identical bodies in a static homogeneous vertical gravitational field at the same time from the same location, one body with vanishing initial velocity, the other with purely horizontal initial velocity, they should hit the ground simultaneously.

But that is clearly impossible to fulfill in *any* special relativistic theory of gravity, independent of whether it is based on a scalar (or vector) field. The reason is this: suppose $-\nabla^\mu\phi = (0, 0, 0, -g)$ is the gravitational field in one inertial frame. Then it takes exactly the same form in any other inertial frame which differs from the first one by 1) spacetime translations, 2) rotations about the z axis, 3) boosts in any direction within the xy -plane. So consider a situation where with respect to an inertial frame F body 1 and body 2 are released simultaneously at $z = 0$ with initial velocities $\vec{v}_1 = (0, 0, 0)$ and $\vec{v}_2 = (v, 0, 0)$. One is interested whether the bodies hit the ‘ground’ simultaneously. The ‘ground’ is represented in spacetime by the hyperplane $z = -h$ and ‘hitting the ground’ is taken to mean that the worldline of the particle in question intersects this hyperplane. Let another inertial frame, F' , move with respect to F at speed v along the x axis. With respect to F' both bodies are released simultaneously at $z' = 0$ with initial velocities $\vec{v}'_1 = (-v, 0, 0)$ and $\vec{v}'_2 = (0, 0, 0)$. The field is still static, homogeneous, and vertical with respect to F' .⁹ In F' the ‘ground’ is defined by $z' = -h$, which defines the *same* hyperplane in spacetime as $z = -h$. This is true since F and F' differ by a boost in x -direction, so that the z and z' coordinates coincide. Hence ‘hitting the ground’ has an invariant meaning in the class of inertial systems considered here. However, if ‘hitting the ground’ are simultaneous events in F they cannot be simultaneous in F' and vice versa, since these events differ in their x coordinates. This leads us to the following

⁹ This is true for gravitational fields that derive from a scalar potential as well as vector potentials. In the scalar case even the strength, $\|\vec{\nabla}\phi\|$, of the field is the same in F and F' , whereas in the vector case the strength in F' is enhanced by a factor $\gamma = 1/\sqrt{1 - v^2/c^2}$. For our argument to work we just need that the field is again static, homogeneous, and vertical. It therefore applies to the scalar as well as the vector case.

Remark 2. Due to the usual relativity of simultaneity, the requirement of ‘hitting the ground simultaneously’ cannot be fulfilled in any Poincaré invariant theory of gravity.

But there is an obvious reinterpretation of ‘hitting the ground simultaneously’, which makes perfect invariant sense in Special Relativity, namely the condition of ‘hitting the ground after the same lapse of eigentime’. As we have discussed in detail above, the scalar theory does indeed fulfill this requirement (independence of (29a) from γ) whereas the vector theory does not (dependence of (41a) on γ).

Remark 3. The scalar theory is distinguished by its property that the eigentime for free fall from a given altitude does *not* depend on the initial horizontal velocity.

In general, with regard to this requirement, the following should be mentioned:

Remark 4. Einstein’s requirement is (for good reasons) not implied by any of the modern formulations of the (weak) equivalence principle, according to which the world-line of a freely falling test-body (without higher mass-multipole-moments and without charge and spin) is determined by its initial spacetime point and four velocity, i.e. independent of the further constitution of the test body. In contrast, Einstein’s requirement relates two motions with *different* initial velocities.

Finally we remark on Einstein’s additional claim in Quote 2, that there is also a similar dependence on the vertical acceleration on the internal energy. This claim, too, does not survive closer scrutiny. Indeed, one might e.g. think at first that (17) also predicts that, for example, the gravitational acceleration of a box filled with a gas decreases as temperature increases, due to the increasing velocities of the gas molecules. But this arguments incorrectly neglects the walls of the box which gain in stress due to the rising gas pressure. According to (4) more stress means more weight. In fact, a general argument due to von Laue [3] shows that these effects precisely cancel. This has been lucidly discussed by Norton [6] and need not be repeated here.

5 Periapsis precession

The Newtonian laws of motion predict that the line of apsides remains fixed relative to absolute space for the motion of a body in a potential with $1/r$ -falloff. Any deviation from the latter causes a rotation of the line of apsides within the orbital plane. This may also be referred to as precession of the periapsis, the orbital point of closest approach to the center of force.

A convenient way to calculate the periapsis precession in perturbed $1/r$ -potentials is provided by the following proposition (taken from an exercise in the textbook on mechanics by Landau & Lifshitz):

Proposition 3. *Consider the Newtonian equations of motion for a test particle of mass m in a perturbed Newtonian potential*

$$U(r) = -\frac{\alpha}{r} + \Delta U(r), \quad (42)$$

where $\alpha > 0$ and $\Delta U(r)$ is the perturbation. The potential is normalized so that it tends to zero at infinity, i.e. $\Delta U(r \rightarrow \infty) \rightarrow 0$. Let $2\pi + \Delta\varphi$ denote the increase of the

polar angle between two successive occurrences of periapsis. Hence $\Delta\varphi$ represents the excess over a full turn, also called the ‘periapsis shift per revolution’. Then the first-order contribution of ΔU to $\Delta\varphi$ is given by

$$\Delta\varphi = \frac{\partial}{\partial L} \left\{ \frac{2m}{L} \int_0^\pi r_*^2(\varphi; L, E) \Delta U(r_*(\varphi; L, E)) d\varphi \right\}. \quad (43)$$

Here $\varphi \mapsto r_*(\varphi; L, E)$ is the solution of the unperturbed problem (Kepler orbit) with angular momentum L and energy E . (As we are interested in bound orbits, we have $E < 0$.) It is given by

$$r_*(\varphi; L, E) = \frac{p}{1 + \varepsilon \cos \varphi}, \quad (44a)$$

where

$$p := \frac{L^2}{m\alpha}, \quad (44b)$$

$$\varepsilon := \sqrt{1 + \frac{2EL^2}{m\alpha^2}}. \quad (44c)$$

Note that the expression in curly brackets on the right hand side of (43) is understood as function of L and E , and the partial differentiation is to be taken at constant E .

Proof. In the Newtonian setting, the conserved quantities of energy and angular momentum for the motion in a plane coordinatized by polar coordinates, are given by

$$E = \frac{1}{2}m(\dot{r}^2 + r^2\dot{\varphi}^2) + U(r), \quad (45)$$

$$L = mr^2\dot{\varphi}', \quad (46)$$

where a prime represents a t -derivative. Eliminating $\dot{\varphi}'$ in (45) via (46) and also using (46) to re-express t -derivatives in terms of φ -derivatives, we have

$$\frac{L^2}{m^2r^4} \left((dr/d\varphi)^2 + r^2 \right) = 2 \frac{E - U}{m}. \quad (47)$$

This can also be write in differential form,

$$d\varphi = \frac{\pm dr L/r^2}{\sqrt{2m(E - U(r)) - L^2/r^2}}, \quad (48)$$

whose integral is just given by (44).

Now, the angular change between two successive occurrences of periapsis is twice the angular change between periapsis (i.e. r_{\min}) and apoapsis (i.e. r_{\max}):

$$\begin{aligned} \Delta\varphi + 2\pi &= 2 \int_{r_{\min}}^{r_{\max}} \frac{dr L/r^2}{\sqrt{2m(E - U(r)) - L^2/r^2}} \\ &= -2 \frac{\partial}{\partial L} \left\{ \int_{r_{\min}}^{r_{\max}} dr \sqrt{2m(E - U(r)) - L^2/r^2} \right\}, \end{aligned} \quad (49)$$

where the term in curly brackets is considered as function of L and E and the partial derivative is for constant E .

Formula (49) is exact. Its sought after approximation is obtained by writing $U(r) = -\alpha/r + \Delta U(r)$ and expanding the integrand up to linear order in ΔU . Taking into account that the zeroth order term just cancels the 2π on the left hand side, we get:

$$\begin{aligned} \Delta\varphi &\approx \frac{\partial}{\partial L} \left\{ 2m \int_{r_{\min}}^{r_{\max}} \frac{\Delta U(r) dr}{\sqrt{2m(E + \alpha/r) - L^2/r^2}} \right\} \\ &\approx \frac{\partial}{\partial L} \left\{ \frac{2m}{L} \int_0^\pi r_*^2(\varphi; L, E) \Delta U(r_*(\varphi; L, E)) d\varphi \right\}. \end{aligned} \quad (50)$$

In the second step we converted the r -integration into an integration over the azimuthal angle φ . This we achieved by making use of the identity that one obtains from (48) with $U(r) = -\alpha/r$ and r set equal to the Keplerian solution curve $r_*(\varphi; L, M)$ for the given parameters L and E . Accordingly, we replaced the integral limits r_{\min} and r_{\max} by the corresponding angles $\varphi = 0$ and $\varphi = \pi + \Delta\varphi/2$ respectively. Since the integrand is already of order ΔU , we were allowed to replace the upper limit by $\varphi = \pi$, so that the integral limits now correspond to the angles for the minimal and maximal radius of the unperturbed Kepler orbit $r_*(\varphi; L, E)$ given by (44a). \square

Let us apply this proposition to the general class of cases where $\Delta U = \Delta_2 U + \Delta_3 U$ with

$$\Delta_2 U(r) = \delta_2/r^2, \quad (51a)$$

$$\Delta_3 U(r) = \delta_3/r^3. \quad (51b)$$

In the present linear approximation in ΔU the effects of both perturbations to $\Delta\varphi$ simply add, so that $\Delta\varphi = \Delta_2\varphi + \Delta_3\varphi$. The contributions $\Delta_2\varphi$ and $\Delta_3\varphi$ are very easy to calculate from (43). The integrals are trivial and give $\pi\delta_2$ and $\pi\delta_3/p$ respectively. Using (44b) in the second case to express p as function of L , then doing the L differentiation and finally eliminating L again in favour of p using (44b), we get

$$\Delta_2\varphi = -2\pi \left[\frac{\delta_2/\alpha}{p} \right] = -2\pi \left[\frac{\delta_2/\alpha}{\alpha(1-\varepsilon^2)} \right], \quad (52a)$$

$$\Delta_3\varphi = -6\pi \left[\frac{\delta_3/\alpha}{p^2} \right] = -6\pi \left[\frac{\delta_3/\alpha}{\alpha^2(1-\varepsilon^2)^2} \right], \quad (52b)$$

where we also expressed p in terms of the semi-major axis a and the eccentricity ε via $p = a(1 - \varepsilon^2)$, as it is usually done. Clearly this method allows to calculate in a straightforward manner the periastris shifts for general perturbations $\Delta_n U = \delta_n/r^n$. For example, the case $n = 3$ is related to the contribution from the quadrupole moment of the central body.

5.1 Scalar model-theory

All this applies directly to the scalar theory if its equation of motion is written in the Newtonian form (19). The static, rotationally symmetric, source-free solution to (4) is

$\Phi(r) = -GM/r$ and hence

$$\tilde{\Phi}(r) = (m_0 c^2/2) \gamma_0^{-2} \left(1 - \frac{GM}{rc^2}\right)^2. \quad (53)$$

In order to normalize the potential so that it assumes the value zero at spatial infinity we just need to drop the constant term. This leads to to

$$\alpha = \gamma_0^{-2} GMm, \quad (54a)$$

$$\delta_2 = \alpha \frac{GM}{2c^2}, \quad (54b)$$

so that

$$\Delta\varphi = \Delta_2\varphi = -\pi \left[\frac{GM/c^2}{a(1-\varepsilon^2)} \right] = -\frac{1}{6} \Delta_{\text{GR}}\varphi, \quad (55)$$

where $\Delta_{\text{GR}}\varphi$ is the value predicted by General Relativity. Hence scalar gravity leads to a *retrograde* periastris precession

5.2 Naive scalar theory

In the naive scalar theory we have $\phi(r) = -GM/r$ in (19) and therefore

$$\begin{aligned} \tilde{\Phi}(r) &= (m_0 c^2/2) \gamma_0^{-2} \exp(-2GM/c^2 r) \\ &= (m_0 c^2/2) \gamma_0^{-2} \left\{ 1 - 2 \left(\frac{GM}{c^2 r} \right) + 2 \left(\frac{GM}{c^2 r} \right)^2 - \frac{4}{3} \left(\frac{GM}{c^2 r} \right)^3 + \dots \right\}. \end{aligned} \quad (56)$$

Again we subtract the constant term to normalize the potential so as to assume the value zero at infinity. Then we simply read off the coefficients α , δ_2 , and δ_3 :

$$\alpha = 2(GM/c^2) (m_0 c^2/2) \gamma_0^{-2}, \quad (57a)$$

$$\delta_2 = 2(GM/c^2)^2 (m_0 c^2/2) \gamma_0^{-2}, \quad (57b)$$

$$\delta_3 = -\frac{4}{3}(GM/c^2)^3 (m_0 c^2/2) \gamma_0^{-2}. \quad (57c)$$

Hence we have

$$\Delta\varphi = \Delta_2\varphi + \Delta_3\varphi, \quad (58a)$$

where

$$\Delta_2\varphi = -2\pi \left[\frac{GM/c^2}{a(1-\varepsilon^2)} \right], \quad (58b)$$

$$\Delta_3\varphi = +4\pi \left[\frac{GM/c^2}{a(1-\varepsilon^2)} \right]^2. \quad (58c)$$

Recall that (52) neglects quadratic and higher order terms in ΔU . If we expand ΔU in powers of $GM/c^2 r$, as done in (56), it would be inconsistent to go further than to third order because ΔU starts with the quadratic term so that the neglected corrections of order $(\Delta U)^2$ start with fourth powers in $GM/c^2 r$. Hence (58) gives the optimal accuracy obtainable with (43). For solar-system applications $GM/c^2 a$ is of the order of 10^{-8} so that the quadratic term (58c) can be safely neglected. Comparison of (58b) with (55) shows that the naive scalar theory gives a value twice as large as that of the consistent model theory, i.e. $-1/3$ times the correct value (predicted by GR).

5.3 Vector theory

We start from the following

Proposition 4. *The equations of motion (33) for a purely ‘electric’ field, where $F_{0i} = -F_{i0} = E_i/c$ and all other components of $F_{\mu\nu}$ vanish, is equivalent to*

$$(\gamma(t)\vec{x}'(t))' = \vec{\mathcal{E}}(\vec{x}(t)), \quad (59)$$

where $'$ denotes d/dt , $\gamma(t) := 1/\sqrt{1 - \|\vec{x}'(t)\|^2/c^2}$, and $\vec{\mathcal{E}} := e\vec{E}/mc$.

Proof. We have $d/ds = \gamma d/dt$, $d\gamma/dt = \gamma^3(\vec{\beta} \cdot \vec{\beta}')$. Now,

$$\ddot{z}^\mu = c\gamma(\gamma', (\gamma\vec{\beta})'), \quad \text{and} \quad (e/m)F_{\nu}^{\mu}\dot{z}^\nu = c\gamma(\vec{\mathcal{E}} \cdot \vec{\beta}, \vec{\mathcal{E}}), \quad (60)$$

so that (33) is equivalent to

$$\vec{\mathcal{E}} \cdot \vec{\beta} = \gamma' = \gamma^3(\vec{\beta} \cdot \vec{\beta}'), \quad (61a)$$

$$\vec{\mathcal{E}} = (\gamma\vec{\beta})' = \gamma^3\vec{\beta}'_{\parallel} + \gamma\vec{\beta}'_{\perp}. \quad (61b)$$

where \parallel and \perp refer to the projections parallel and perpendicular to $\vec{\beta}$ respectively. Since (61b) implies (61a), (33) is equivalent to the former. \square

We apply this to a spherically symmetric field, where $\vec{\mathcal{E}} = -\vec{\nabla}\phi$ with $\phi = \phi(r) = -GM/r$. This implies conservation of angular momentum, the modulus of which is now given by

$$L = \gamma m r^2 \varphi' \quad (62)$$

Note the explicit appearance of γ , which e.g. is not present in the scalar case, as one immediately infers from (17). This fact makes Proposition 3 not immediately applicable. We proceed as follows: scalar multiplication of (59) with $\vec{v} = \vec{x}'$ and m leads to the following expression for the conserved energy:

$$E = mc^2(\gamma - 1) + U, \quad (63)$$

where $U = m\phi$. This we write in the form

$$\gamma^2 = \left(1 + \frac{E - U}{mc^2}\right)^2. \quad (64a)$$

On the other hand, we have

$$\gamma^2 \equiv 1 + (\beta\gamma)^2 = 1 + (\gamma/c)^2(r'^2 + r^2\varphi'^2) = 1 + \frac{L^2}{m^2c^2r^4} \left((dr/d\varphi)^2 + r^2 \right), \quad (64b)$$

where we used (62) to eliminate φ' and convert r' into $dr/d\varphi$, which also led to a cancellation of the factors of γ . Equating (64a) and (64b), we get

$$\frac{L^2}{m^2r^4} \left((dr/d\varphi)^2 + r^2 \right) = 2 \frac{\tilde{E} - \tilde{U}}{m} \quad (65)$$

where

$$\tilde{E} := E(1 + E/2mc^2), \quad (66a)$$

$$\tilde{U} := U(1 + E/mc^2) - U^2/2mc^2. \quad (66b)$$

Equation (65) is just of the form (47) with \tilde{E} and \tilde{U} replacing E and U . In particular we have for $U = m\phi = -GMm/r$:

$$\tilde{U}(r) = -\frac{\alpha}{r} + \frac{\delta_2}{r^2} \quad (67)$$

with

$$\alpha = GMm(1 + E/mc^2), \quad (68a)$$

$$\delta_2 = -\frac{G^2M^2m}{2c^2}. \quad (68b)$$

In leading approximation for small E/mc^2 we have $\delta_2/\alpha = -GMm/2c^2$. The advance of the periapsis per revolution can now be simply read off (52a):

$$\Delta\varphi = \pi \left[\frac{GM/c^2}{a(1 - \varepsilon^2)} \right] = \frac{1}{6}\Delta_{GR}\varphi. \quad (69)$$

This is the same amount as in the scalar model-theory (compare (55)) but of opposite sign, corresponding to a *prograde* periapsis precession of 1/6 the value predicted by General Relativity.

6 Energy conservation

From a modern viewpoint, Einstein’s claim as to the violation of energy conservation seems to fly in the face of the very concept of a Poincaré invariant theory. After all, time translations are among the symmetries of the Poincaré group, thus giving rise to a corresponding conserved Noether charge. Its conservation is a theorem and cannot be questioned. The only thing that seems logically questionable is whether this quantity does indeed represent physical energy. So how does Einstein arrive at his conclusion?

6.1 Einstein’s argument

Einstein first pointed out that the source for the gravitational field must be a scalar built from the matter quantities alone, and that the only such scalar is the trace T_{μ}^{μ} of the energy-momentum tensor (as pointed out to Einstein by von Laue, as Einstein acknowledges, calling T_{μ}^{μ} the “Laue Scalar”). Moreover, for *closed stationary systems* the so-called Laue-Theorem (first proven in [3] for static systems and later slightly generalized to stationary ones) states that the integral over space of $T^{\mu\nu}$ must vanish, except for $\mu = 0 = \nu$; hence the space integral of T_{μ}^{μ} equals that of T^{00} , which means that the total (active and passive) gravitational mass of a closed stationary system equals its inertial mass. However, if the system is not closed, the weight depends on the stresses (the spatial components T^{ij}).

His argument proper is then as follows (compare Fig. 1): consider a box B filled with electromagnetic radiation of total energy E . We idealize the walls of the box to be inwardly perfectly mirrored and of infinite stiffness, i.e. they can support normal stresses (pressure) without any deformation. The box has an additional vertical strut in the middle connecting top and bottom walls, which supports all the vertical material stresses that counterbalance the radiation pressure, so that the side walls merely sustain normal and no tangential stresses. The box can slide without friction along a vertical shaft whose cross section corresponds exactly to that of the box. The walls of the shaft are likewise idealized to be inwardly perfectly mirrored and of infinite stiffness. The whole system of shaft and box is finally placed in a homogeneous static gravitational field, \vec{g} , which points vertically downward. Now we perform the following process. We start with the box being placed in the shaft in the upper position. Then we slide it down to the lower position; see Fig. 2. There we remove the side walls of the box—without any radiation leaking out—such that the sideways pressures are now provided by the shaft walls. The strut in the middle is left in position to further take all the vertical stresses, as before. Then the box together with the detached side walls are pulled up to their original positions; see Fig. 3. Finally the system is reassembled so that it assumes its initial state. Einstein’s claim is now that in a very general class of imaginable scalar theories the process of pulling up the parts needs less work than what is gained in energy in letting the box (with side walls attached) down. Hence he concluded that such theories necessarily violate




Figure 1: Sliding box filled with radiation in a gravitational field \vec{g} .




Figure 2: Lowering the box in the gravitational field with side walls attached.




Figure 3: Raising the box in the gravitational field with side walls taken off.

energy conservation.

Indeed, radiation-plus-box is a closed stationary system in von Laue's sense. Hence the weight of the total system is proportional to its total energy E , which we may pretend to be given by the radiation energy alone since the contributions from the rest masses of the walls will cancel in the final energy balance, so that we may formally set them to zero at this point. Lowering this box by an amount h in a static homogeneous gravitational field of strength g results in an energy gain of $\Delta E = hgE/c^2$. So despite the fact that radiation has a traceless energy-momentum tensor, *trapped* radiation has a weight given by E/c^2 . This is due to the radiation pressure which puts the walls of the trapping box under tension. Tension makes an independent contribution to weight, independent of the material that supports it. For each parallel pair of side-walls the tension is just the radiation pressure, which is one-third of the energy density. So each pair of side-walls contribute $E/3c^2$ to the (passive) gravitational mass (over and above their rest mass, which we set to zero) in the lowering process when stressed, and zero in the raising process when unstressed. Hence, Einstein concluded, there is a net gain in energy of $2E/3c^3$ (there are two pairs of side walls).

But it seems to me that Einstein neglects a crucial contribution to the energy balance. In contrast to the lowering process, the state of the shaft S is changed during the lifting process, and it is this additional contribution which just renders Einstein's argument inconclusive. Indeed, when the side walls are first removed in the lower position, the walls of the shaft necessarily come under stress because they now need to provide the horizontal balancing pressures. In the raising process that stress distribution of the shaft is translated upwards. But that *does* cost energy in the theory discussed here, even though it is not associated with any proper transport of the material the shaft is made from. As already pointed out, stresses make their own contribution to weight, independent of the nature of the material that supports them. In particular, a redistribution of stresses in a material immersed in a gravitational field generally makes a non-vanishing contribution to the energy balance, even if the material does not move.

This will be seen explicitly below.

6.2 Energy conservation in the scalar model-theory

Corresponding to Poincaré-invariance there are 10 conserved currents. In particular, the total energy E relative to an inertial system is conserved. For a particle coupled to gravity it is easily calculated and consists of three contributions corresponding to the gravitational field, the particle, and the interaction-energy of particle and field:

$$E_{\text{gravity}} = \frac{1}{2\kappa c^2} \int d^3x \left((\partial_{\text{ct}}\Phi)^2 + (\vec{\nabla}\Phi)^2 \right), \quad (70a)$$

$$E_{\text{particle}} = m_0 c^2 \gamma(v), \quad (70b)$$

$$E_{\text{interaction}} = m_0 \gamma(v) \Phi(\vec{z}(t), t), \quad (70c)$$

Let's return to general matter models and let $T_{\text{total}}^{\mu\nu}$ be the total stress-energy tensor of the gravity-matter-system. It is the sum of three contributions:

$$T_{\text{total}}^{\mu\nu} = T_{\text{gravity}}^{\mu\nu} + T_{\text{matter}}^{\mu\nu} + T_{\text{interaction}}^{\mu\nu}, \quad (71)$$

where¹⁰

$$T_{\text{gravity}}^{\mu\nu} = \frac{1}{\kappa c^2} \left(\partial^\mu \Phi \partial^\nu \Phi - \frac{1}{2} \eta^{\mu\nu} \partial_\lambda \Phi \partial^\lambda \Phi \right), \quad (72a)$$

$$T_{\text{matter}}^{\mu\nu} = \text{depending on matter model}, \quad (72b)$$

$$T_{\text{interaction}}^{\mu\nu} = \eta^{\mu\nu} (\Phi/c^2) T_{\text{matter}}. \quad (72c)$$

Energy-momentum-conservation is expressed by

$$\partial_\mu T_{\text{total}}^{\mu\nu} = F_{\text{external}}^\nu, \quad (73)$$

where F_{external}^ν is the four-force of a possible *external* agent. The 0-component of it (i.e. energy conservation) can be rewritten in the form

$$\text{external power supplied} = \frac{d}{dt} \int_D d^3x T_{\text{total}}^{00} + \int_{\partial D} T_{\text{total}}^{0k} n_k d\Omega. \quad (74)$$

If the matter system is of finite spatial extent, meaning that outside some bounded spatial region, D , we have that $T_{\text{matter}}^{\mu\nu}$ vanishes identically, and if we further assume that no gravitational radiation escapes to infinity, the surface integral in (74) vanishes identically. Integrating (74) over time we then get

$$\text{external energy supplied} = \Delta E_{\text{gravity}} + \Delta E_{\text{matter}} + \Delta E_{\text{interaction}}, \quad (75)$$

with

$$E_{\text{interaction}} = \int_D d^3x (\Phi/c^2) T_{\text{matter}}, \quad (76)$$

and where $\Delta(\text{something})$ denotes the difference between the initial and final value of 'something'. If we apply this to a process that leaves the *internal* energies of the

¹⁰ We simply use the standard expression for the canonical energy-momentum tensor, which is good enough in the present case. If $S = \int L dt d^3x$, it is given by $T_\nu^\mu := (\partial L / \partial \Phi_{,\mu}) \Phi_{,\nu} - \delta_\nu^\mu L$.

gravitational field and the matter system unchanged, for example a processes where the matter system, or at least the relevant parts of it, are *rigidly* moved in the gravitational field, like in Einstein’s Gedankenexperiment of the ‘radiation-shaft-system’, we get

$$\text{external energy supplied} = \Delta \left\{ \int_{\text{D}} d^3x (\Phi/c^2) T_{\text{matter}} \right\}. \quad (77)$$

Now, my understanding of what a valid claim of energy non-conservation in the present context would be is to show that *this* equation can be violated. But this is *not* what Einstein did (compare Conclusions).

If the matter system stretches out to infinity and conducts energy and momentum to infinity, than the surface term that was neglected above gives a non-zero contribution that must be included in (77). Then a proof of violation of energy conservation must disprove this modified equation. (Energy conduction to infinity as such is not in any disagreement with energy conservation; you have to prove that they do not balance in the form predicted by the theory.)

6.3 Discussion

For the discussion of Einstein’s Gedankenexperiment the term (76) is the relevant one. It accounts for the *weight of stress*. Pulling up a radiation-filled box inside a shaft also moves up the stresses in the shaft walls that must act sideways to balance the radiation pressure. This lifting of stresses to higher gravitational potential costs energy, according to the theory presented here. This energy was neglected by Einstein, apparently because it is not associated with a transport of matter. He included it in the lowering phase, where the side-walls of the box are attached to the box and move with it, but neglected them in the raising phase, where the side walls are replaced by the shaft, which does not move. But as far as the ‘weight of stresses’ is concerned, this difference is irrelevant. What (76) tells us is that raising stresses in an ambient gravitational potential costs energy, irrespectively of whether it is associated with an actual transport of the stressed matter or not. This would be just the same for the transport of heat in a heat-conducting material. Raising the heat distribution against the gravitational field costs energy, even if the material itself does not move.

7 Conclusion

From the foregoing I conclude that, taken on face value, neither of Einstein’s reasonings that led him to dismiss of a scalar theories of gravity prior to being checked against experiments are convincing. For example, I would not use them in lectures on General Relativity.

First, energy—as defined by Noether’s theorem—is conserved in our model theory. Note also that the energy of the free gravitational field is positive definite in this theory. Secondly, the *eigentime* for free fall in a homogeneous static gravitational field *is* independent of the initial horizontal velocity. Hence our model-theory serves as an example of an internally consistent theory which, however, is experimentally ruled out. As we have seen, it predicts $-1/6$ times the right perihelion advance of Mercury and also no light deflection (not to mention Shapiro time-delay and various other accurately measured effects which are correctly described by GR).

The situation is slightly different in a special-relativistic vector theory of gravity (Spin 1, mass 0). Here the energy is clearly still conserved (as in any Poincaré invariant theory), but the energy of the radiation field is negative definite due to a sign change in Maxwell's equations which is necessary to make like charges (i.e. masses) attract rather than repel each other. Hence there exist runaway solutions in which a massive particle self-accelerates unboundedly by radiating negative gravitational radiation. Also, the free-fall eigentime now does depend on the horizontal velocity. Hence, concerning these theoretical aspects, scalar gravity is much better behaved.

Finally I wish to mention another general aspect that is relevant to the present discussion.¹¹ Consider the dynamical problem of an electromagnetically bound system, like an atom, where (classically speaking) an electron orbits a charged nucleus (both modelled as point masses). Place this system in a gravitational field that varies negligibly over the spatial extent of the atom and over the time of observation. The electromagnetic field produced by the charges will be unaffected by the gravitational field (due to its traceless energy momentum tensor). In contrast, (15) tells us that the dynamics of the particle is influenced by the gravitational field, even for strictly constant potentials. The effect can be conveniently summarized by saying that the masses of point particles scale by a factor of $1 + \Phi/c^2 = \exp(\Phi/c^2)$ when placed in the potential Φ . This carries over to Quantum Mechanics so that atomic length scales, like the Bohr radius (in MKSA units)

$$a_0 := \frac{\varepsilon_0 \hbar^2}{m \pi e^2} \quad (78)$$

and time scales, like the Rydberg period (inverse Rydberg frequency)

$$T_R := \frac{8\varepsilon_0^2 \hbar^3}{m e^4} \quad (79)$$

change by a factor $\exp(-\Phi/c^2)$ due to their inverse proportionality to the electron mass m (\hbar is Planck's constant, e the electron charge, and ε_0 the vacuum permittivity). This means that, relative to the units on which the Minkowski metric is based, atomic units of length and time vary in a way depending on the potential. Transporting the atom to a spacetime position in which the gravitational potential differs by an amount $\Delta\Phi$ results in a diminishment (if $\Delta\Phi > 0$) or enlargement (if $\Delta\Phi < 0$) of its size and period *relative to Minkowskian units*. This effect is universal for all atoms.

The question then arises as to the operational significance of the latter. Should we not rather *define* spacetime lengths by what is measured using atoms? After all, as Einstein repeatedly remarked, physical notions of spatial lengths and times should be based on physically constructed rods and clocks which are consistent with our dynamical equations. The Minkowski metric would then merely turn into a redundant structure without direct observational content.¹² From that perspective one may indeed criticize special-relativistic scalar gravity for making essential use of dispensable absolute structures, which eventually should be eliminated, just like in the 'flat-spacetime-approach' to GR (see e.g. [8] and Sect. 5.2 in [2] for more references). In view of

¹¹ I thank John Norton for asking a question that led to these remarks.

¹² Note that the argument presented here, which is merely based on the Lagrangian for point-particles (which is also the relevant one in Quantum-Mechanics) does not show that in full generality. 'Clocks' and 'rods' not based on atomic frequencies and lengths scales are clearly conceivable.

Quote 1 one might conjecture that this more sophisticated point was behind Einstein's criticism. If so, it is well taken. But physically it should be clearly separated from the other accusations which we discussed here.

References

- [1] Albert Einstein. *Mein Weltbild*. Ullstein Verlag, Berlin, 1977.
- [2] Domenico Giulini and Norbert Straumann. Einstein's impact on the physics of the twentieth century. *Studies in the History and Philosophy of Modern Physics*, 37:115–173, 2006. ArXiv physics/0507107.
- [3] Max von Laue. Zur Dynamik der Relativitätstheorie. *Annalen der Physik*, 35:524–542, 1911. *Gesammelte Schriften und Vorträge* (Friedrich Vieweg & Sohn, Braunschweig, 1961), Vol. I, p 524-542.
- [4] Max von Laue. Die Nordströmsche Gravitationstheorie. *Jahrbuch der Radioaktivität und Elektronik*, 14(3):263–313, 1917.
- [5] John Norton. Einstein, Nordström and the early demise of scalar Lorentz-covariant theories of gravitation. *Archive for the History of Exact Sciences*, 45:17–94, 1992.
- [6] John Norton. Einstein and Nordström: Some lesser known thought experiments in gravitation. In John Earman, Michel Janssen, and John Norton, editors, *The Attraction of Gravitation: New Studies in History of General Relativity*, pages 3–29, Boston, MA, 1993. Birkäuser Verlag.
- [7] John Stachel et al., editors. *The Collected Papers of Albert Einstein, Vols. 1-9*. Princeton University Press, Princeton, New Jersey, 1987-2005.
- [8] Walter E. Thirring. An alternative approach to the theory of gravitation. *Annals of Physics*, 16:96–117, 1961.