

The Opportunities of Using Mushroom Compost Waste in Strawberry Growing

Ahsen Işık ÖZGÜVEN

Department of Horticulture, Faculty of Agriculture, University of Çukurova, 01330 Adana-TURKEY

Received: 13.01.1997

Abstract: The aim of this study is to see whether Mushroom compost waste (MCW) can be used as an alternative fertilizers to Farm yard manure (FYM) because the latter is very expensive in Strawberry growing.

Experiment was carried out with frigo plants of Douglas cv. in summer planting system. In this experiment, both MCW (test material) and FYM (as control) were used. The obtained MCW and FYM were used 10, 20 and 40 tons per hectare. Experiment was designed as Randomized Blocks with 3 replicates. Applications in question during the growing season are observed the earliness, the yield per plant and fruit quality.

According to the results of this two-year-experiment, it was determined that MCW can be used as an alternative fertilizer to FYM in strawberry growing.

Çilek Yetiştiriciliğinde Mantar Kompostu Atıklarının Kullanılma Olanakları

Özet: Bu çalışmanın amacı yanmış çiftlik gübresinin çok pahalı olmasından dolayı çilek yetiştiriciliğinde mantar kompostu atığının çiftlik gübresine alternatif olarak kullanılıp kullanılamayacağını araştırmaktır.

Deneme Douglas çilek çeşidiyle, yaz dikim sisteminde frigo bitkiler ile kurulmuş ve hem mantar kompostu atığı (test materyali) hem de yanmış çiftlik gübresi (kontrol) kullanılmıştır. Mantar kompost atığı ve yanmış çiftlik gübresi hektara 10, 20 ve 40 ton dozlarında kullanılmıştır. Deneme 3 yinelemeli Tesadüf Bloklarına göre yapılmıştır. Yetiştirme periyodu boyunca erkencilik, bitki başına verim ve meyve kaliteleri incelenmiştir.

İki yıllık denemenin sonuçlarına göre çilek yetiştiriciliğinde mantar kompostu atığının yanmış çiftlik gübresine alternatif olarak kullanılabileceği saptanmıştır.

Introduction

In Turkey's agriculture, generally farm yard manure (FYM) is used commonly to enrich the soil by organic matter. But to take a good result from FYM, it must have been matured. Otherwise some plants sometimes die because of nonmaturation of the FYM. Nowadays maturing the FY is done in ordinary conditions. However there are several studies on maturing it (1-7). As a result, some important criteria were established. It is stated that it can prevent the effect of harmful microorganism and stimulate the growth and development of the plant in question.

Because matured FYM is very expensive, and the germination of different weed seed problem increases the cost, people try to find other ways that will help to increase the yield and quality (8, 9).

In this study, it is hypothesized that we can use waste of mushroom compost (MCW) as the organic material. The waste mushroom compost has the characteristics of

plant growing medium both physically and chemically. Some research conducted for this purpose has shown that MCW can be a good soilless cultur substrate when it is mixed with soil as a farmyard fertilizer or used alone in order to recover the physical conditions of the soil (10-14).

In Turkey, mushroom growing has developed in the last twenty years. The result of some studies have revealed that mushroom growing is approximately 7000-10000 tonnes per year. Since such an increasing amount of waste, has been obtained this waste can be used in strawberry growing.

Materials and Methods

The obtained mushroom compost waste (MCW) administered to the raised beds in the rates of 10, 20 and 40 tons per hectare (ha), and were mixed on the soil. As control, farm yard manure (FYM) was used also in the same rates.

	N(%)	P(%)	K(%)	Fe (ppm)	Zn (ppm)	Mn (ppm)	Cu (ppm)
FYM	1.1795	0.1673	0.9	2989.9	77.5	384.8	44.0
MC	1.6275	0.1238	1.0	606.2	44.8	158.9	157.8

Media	MC doses			FYM doses		
	Yield (g/plant)					
Months	10 ton/ha	20 ton/ha	40 ton/ha	10 ton/ha	20 ton/ha	40 ton/ha
March	24.4	36.2	50.1	32.9	27.1	38.9
April	266.9	260.6	239.3	233.6	262.0	241.2
May	171.9	141.5	130.1	122.2	143.8	130.5
June	158.7	128.5	128.4	119.5	140.1	185.4
Total	621.9	566.8	547.9	508.2	573.0	596.0
Mean	578.9			559.1		

Table 1. Levels of macro and micro elements in the FYM and MC.

Table 2. The effects of MC and FYM on the total and monthly yield (g/plant) during 1993-94 period.

D%5 (Doses)=N.S. D%5 (Media)=N.S. D%5 (Doses*Media)=N.S.

Experiment was carried out with frigo plants of Douglas cv. as a summer planting. The experiment was designed as a Randomized Blocks with 3 replicates. Each replicate which is a parcel contains 30 plants. Statistical analyses were resolved by COSTAT pocket program.

The experiment continued for two years. In the first year, plants were grown under high tunnel during winter period. In the second year, plants were grown in the open conditions within the same period of time. In both trials, the effects of MCW and FYM on the earliness, yield per plant, and berry quality were observed (15).

Result and Discussion

Mineral Nutrient Contents of MCW and FYM

Macro and micro nutrient contents of the MCW and FYM are given in Table 1. In terms of macro elements, the amount of P and K (except N) were almost a like in both media. The amount of N in MCW was observed to be higher (1.6275%) than FYM. In terms of micro elements, Fe, Zn and Mn (except Cu) were higher in FYM than in MCW.

The yield and its monthly distribution

During the whole growing period of 1993-94, the highest yield (621.9 g/plant) was obtained from the 10 tons/ha dose of MCW (Table 2, Figure 1). It was followed by 40 tons/ha of the FYM (596.0 g/plant).

During the 1993-94 season in all the treatments the harvests started in March. The highest yield in March was obtained from the 40 tons/ha dose of MCW (50.1 g/plant).

During the whole growing period of 1994-95, the highest yield (490.5 g/plant) was obtained from the 10 tons/ha dose of MCW as it was the same last year (Table 3, Figure 2). It was followed by 10 tons/ha of the FYM (472.2 g/plant).

As earliness yield, during the 1994-95 season in March the highest yield was obtained from the 40 tons/ha of the FYM and 10 tons/ha of the MCW (22.0 g, 19.3 g/plant, respectively) (Table 3).

When Table 2 and 3 are studied together, it can be seen that the yield is higher in 1993-94 than 1994-95 season. The reason of this low yield may be that plants were grown in walk-in high tunnel in the first year (16). The same results were observed in precocity. In the trial of 1993-94, the yield of March was more than the yield of March 1994-95. This may be due to the fact that the plants were grown in high tunnel in which they grow, develop, and give flowers earlier than plants in the field (16).

Average Berry Weights

Average berry weights in both experimental year are given Table 4 and 5. The largest berries were obtained in March of 1993-94 and 1994-95 seasons (Figure 3, 4).




Figure 1. The effects of MCW and FYM on the yield during 1993-94 period (M10: 10 tons/ha; M20: 20 tons/ha; M40: 40 tons/ha of MCW and F10: 10 tons/ha; F20: 20 tons/ha; F40: 40 tons/ha of FYM).




Figure 2. The effects of MCW and FYM on the yield during 1994-95 period (M10: 10 tons/ha; M20: 20 tons/ha; M40: 40 tons/ha of MCW and F10: 10 tons/ha; F20: 20 tons/ha; F40: 40 tons/ha of FYM).

Media	MCW doses			FYM doses		
	Yield (g/plant)					
Months	10 ton/ha	20 ton/ha	40 ton/ha	10 ton/ha	20 ton/ha	40 ton/ha
March	19.3	13.1	6.7	16.6	5.1	22.0
April	322.0	234.3	119.5	277.4	186.3	280.2
May	149.2	118.7	57.3	178.2	111.2	149.1
Total	490.5	366.1	183.5	472.2	302.5	451.2
Mean	346.7			408.6		

Table 3. The effects of MCW and FYM on the total and monthly yield (g/plant) during 1994-95 period.

D%5 (Doses)=N.S. D%5 (Media)=N.S. D%5 (Doses*Media)=N.S.

In 1993-94, the average weights of berries were 24.4 g with 40 tons/ha dose of MCW and 22.9 g with 10 tons/ha and 20 tons/ha of FYM.

In 1994-95, the largest berries were obtained from

10 tons/ha of MCW with 24.7 g/plant and 40 tons/ha of FYM with 22.4 g/plant. In all trials, as the season progressed the berry weight decreased. This decrease is due to the characteristic of strawberry because the earlier flowers in strawberry give larger fruits (16).

Media	MCW doses			FYM doses		
	10 ton/ha	20 ton/ha	40 ton/ha	10 ton/ha	20 ton/ha	40 ton/ha
Berry Weight (g)						
Months	10 ton/ha	20 ton/ha	40 ton/ha	10 ton/ha	20 ton/ha	40 ton/ha
March	20.6	20.7	24.4	22.9	22.9	18.0
April	9.4	10.0	10.0	9.7	10.6	9.3
May	8.8	8.4	9.1	8.7	8.7	8.5
June	7.1	5.8	7.0	6.2	6.2	6.2
Average	11.5	11.2	12.6	11.9	12.1	10.5
	11.8			11.5		

Table 4. The effects of MCW and FYM on the average berry weights (g) in different months of 1993-94 period.

D%5 (Doses)=N.S. D%5 (Media)=N.S. D%5 (Doses*Media)=N.S.




Figure 3. The effects of MCW and FYM on the average berry weights (g) in different months of 1993-94 period (M10: 10 tons/ha; M20: 20 tons/ha; M40: 40 tons/ha of MCW and F20: 10 tons/ha; F20: 20 tons/ha; F40: 40 tons/ha of FYM).




Figure 4. The effects of MCW and FYM on the average berry weights (g) in different months of 1994-95 period (M10: 10 tons/ha; M20: 20 tons/ha; M40: 40 tons/ha of MCW and F10: 10 tons/ha; F20: 20 tons/ha; F40: 40 tons/ha of FYM).

Total Soluble Solid Contents of Berries

TSS contents of the berries generally increased as the season progressed; that means is warmer the days sweeter the berries (Table 6 and 7, Figure 5 and 6). In all treatments there is no statistical significant difference in terms of TSS.

In the first year of the experiment the seasonal mean

TSS values were changed between 7.6% and 8.6% (Table 6).

In the second year, the highest content of TSS was obtained from 20 tons/ha dose of FYM with 8.8% and this was almost the same with the other doses.

These results support the findings of other studies (10-14) that when MCW is mixed with soil just as FYM or

Media	MCW doses			FYM doses		
	Berry Weight (g)					
Months	10 ton/ha	20 ton/ha	40 ton/ha	10 ton/ha	20 ton/ha	40 ton/ha
March	24.7	21.0	18.8	21.2	20.2	22.4
April	13.8	13.2	11.4	15.5	14.4	14.1
May	8.8	8.9	8.1	8.6	8.1	9.0
Average	15.8	14.4	12.8	15.1	14.2	15.1
		14.3			14.8	

Table 5. The effects of MCW and FYM on the average berry weights (g) in different months of 1994-95 period.

D%5 (Doses)=N.S. D%5 (Media)=N.S. D%5 (Doses*Media)=N.S.

Media	MCW doses			FYM doses		
	TSS (%)					
Months	10 ton/ha	20 ton/ha	40 ton/ha	10 ton/ha	20 ton/ha	40 ton/ha
March	7.6	8.0	6.0	7.7	7.0	7.8
April	8.1	8.6	7.7	7.7	7.4	8.2
May	8.4	8.6	8.9	8.3	8.9	9.6
June	8.5	8.8	8.3	8.3	8.2	8.6
Mean	8.2	8.5	7.6	7.9	7.9	8.6

Table 6. The effects of MCW and FYM doses on the TSS contents (%) of the berries during 1993-94.

D%5 (Doses)=N.S. D%5 (Media)=N.S. D%5 (Doses*Media)=N.S.

Media	MCW doses			FYM doses		
	TSS (%)					
Months	10 ton/ha	20 ton/ha	40 ton/ha	10 ton/ha	20 ton/ha	40 ton/ha
March	8.2	7.6	8.1	7.9	7.8	8.1
April	7.9	8.1	8.0	7.7	8.2	7.8
May	9.9	9.6	9.9	8.9	10.4	10.1
Mean	8.7	8.4	8.7	8.2	8.8	8.7

Table 7. The effects of MCW and FYM doses on the TSS contents (%) of the berries during 1994-95.

D%5 (Doses)=N.S. D%5 (Media)=N.S. D%5 (Doses*Media)=N.S.

when it is used without soil, it recovers the physical conditions of media.

According to the results, it can be said that the MCW has almost the same positive effects with FYM on the

yield and quality of the Douglas strawberry variety. In other words MCW can be considered as an alternative to FYM.




Figure 5. The effects of MCW and FYM on the total soluble solid contents (%) of berries during 1993-94 period (M10: 10 tons/ha; M20: 20 tons/ha; M40: 40 tons/ha of MCW and F20: 10 tons/ha; F20: 20 tons/ha; F40: 40 tons/ha of FYM).




Figure 6. The effects of MCW and FYM on the total soluble solid contents (%) of berries during 1994-95 period (M10: 10 tons/ha; M20: 20 tons/ha; M40: 40 tons/ha of MCW and F10: 10 tons/ha; F20: 20 tons/ha; F40: 40 tons/ha of FYM).

References

1. Hadar, Y., Inbar, Y., Chen, Y., Effect of compost maturity on tomato seedling growth. *Scientia Horticulturae*, 27: 199-208. Elsevier Science Publishers B.V., Amsterdam-Netherlands, 1985.
2. Hoitink, H.A.J. Fahy, P.C. Basis for the control of soilborne plant pathogens with composts. *Ann. Rev. Phytopathol.* 24: Phytopathol. 1986.
3. Özgüven, M., Kaya, Z. Tütün atıklarının tarımda gübre olarak kullanılma olanakları üzerinde bir araştırma. *Ulusal Çevre Simpozyumu Tebliğ Metinleri*, Adana. TÜBİTAK Yayınları. No:12. 240-245. 1984.
4. Inbar, Y., Chen, Y., Hadar, Y., Composting of agricultural waste for their use as container media. Simulation of the composting process biological waste 26: 247-259. Elsevier Science Publishers Ltd. England. 1988.
5. Inbar, Y., Chen, Y., Hadar, Y. Hoitink, H.A.J. Approaches to determining compost maturity. *The biocycle guide to the art and science of composting*. The JG Press, Inc. Emmaus, Pennsylvania 183-187. 1990.
6. Inbar, Y., Chen, Y., Hadar, Y. Carbon-13 CPMAS NMR and FTIR spectroscopic analysis of organic matter transformations during composting of solid wastes from wineries. *Soil Science Vol. 152* No.41 p:272-282. 1991.
7. Türemiş, N., Kaşka, N., Kaya, Z., Özgüven, A.I., Bazı tarımsal atıkların kompost yapmak suretiyle tekrar kullanılma olanakları. *Tarım-Çevre İlişkileri Sempozyumu*. 13-15 Mayıs 1996 Mersin (In press).
8. Özgüven, A.I., Kaşka, N., Türemiş, N. The opportunities of using tobacco compost in strawberry growing. 3rd. Inter. Strawberry symposium Holland. *Acta Horticulturae* (in press). 1996.

9. Özgüven, A.I. Kaşka, N., Türemiş, N. The effects of gross clipping wastes on the precocity, yield and quality of the strawberry. 3rd. Inter. Strawberry Symposium. Holland. Acta Horticulturae (in press), 1996.
10. Abak, K., Yanmaz, R., Koçyiğit, A.E. Mantar kompost atığının sebze yetiştiriciliğinde değerlendirilmesi. Tarım ve Köyüşleri Bah. Dergisi 25, 16-18, 1986.
11. Abak, K., Yanmaz, R., Ilbay, M.E. Kullanılmış mantar kompostunun serada biber yetiştiriciliğinde kullanımı. Türkiye I. Ulusal Bahçe Bitkileri Kongresi Tebliğleri, Cilt II, Ekim 1992, İzmir p:367-370. 1992.
12. Çelikel, G. Organik ve inorganik kökenli bazı ortamların serada topraksız yetiştiricilikte kullanılabilirliği ile domates, biber ve patlıcanda bitki gelişmesi, verim, erkencilik ve kalite üzerine etkileri. Çukurova Univ. Fen Bilimleri Enstitüsü (Doktora Tezi), Adana, 1994.
13. Paksoy, M. Domateste topraksız yetiştiricilikte değişik substrat karışımlarının ve bitki kök bölgesi ısıtmasının bitki gelişimi, verim, erkencilik ve ürün kalitesine etkileri. Çuk. Üniv. Fen Bilim. Enstitüsü (Doktora Tezi), Adana 1995.
14. Abak, K., Çelikel, G. Mantar kompostunun seralarda topraksız yetiştiricilikte kullanımı. Türkiye 5. Yemeklik mantar kongresi (Bildiri ve Poster Özetleri) 5-7 Kasım 1996. Atatürk Bahçe Kültürleri Merkez Araştırma Enstitüsü Yalova p:28 1996.
15. Özgüven, A.I., Kaşka, N., Effects of slow release and chemical fertilizers on precocity, yield and fruit quality of strawberries J. Agric. Fac. Ç.Ü. 7(3) 177-190. 1992.
16. Kaşka, N., Özgüven, A.I., Paydaş, S., Biçici, M., Türemiş, N., Küden, A. Türkiye için yeni bazı çilek çeşitlerinin Adana'da yaz ve kış dikim sistemiyle örtü altında yetiştiriciliğinin verim, kalite ve erkencilik üzerine etkileri. Doğa Bilim Dergisi (ser.D2) 10(1): 84-104. 1986.