

# Development of the nose and soft tissue profile

By Jeffrey S. Genecov, DDS, MSD; Peter M. Sinclair, DDS, MSD; and Paul C. Dechow, PhD

As orthodontic diagnosis and treatment planning have become more sophisticated and scientific, much attention has been paid to techniques for determining the skeletal pattern, the amount and direction of facial growth, and the position of the dentition.<sup>1-7</sup> Far less attention has been directed towards providing information which would aid the clinician in producing a well-proportioned, balanced and harmonious soft tissue profile at the end of treatment.<sup>8-11</sup>

Facial esthetics have interested orthodontists for many years and although opinions as to what constitutes an attractive face have come from many sources,<sup>3,12-14</sup> there is still a considerable lack of information as to the longitudinal development of the nose, lips and soft tissue chin. This deficiency effectively precludes the development of any scientifically based diagnostic scheme to aid in the objective integration

of soft tissue factors into an overall treatment plan.

Burstone stated that "the facial objective of the orthodontist might be considered the achievement of the optimal in facial harmony consistent with the maximum in functional occlusion within the limits of therapy."<sup>15</sup> Early studies suggested there was a constancy of the pattern of skeletal growth and assumed that the development of the soft tissue profile was coincident with the underlying hard tissues.<sup>16,17</sup> However, a considerable body of evidence now exists which suggests that different components of the soft tissue profile have differing rates and timing of growth and that all parts of the soft tissue profile do not grow in direct proportion to their skeletal bases.<sup>18-22</sup> Increasing amounts of soft tissue convexity with age have been demonstrated, as has considerable sexual dimorphism with regard to soft tissue thickness, as well as



## Abstract

Cephalometric radiographs from a sample of 64 untreated persons (32 Class I and 32 Class II) were evaluated to determine the amount, direction and timing of facial soft tissue development. Twenty-five parameters were evaluated in the mixed dentition (7 to 9 years), the early permanent dentition (11 to 13 years), and early adulthood (16 to 18 years). Results showed that anteroposterior growth and subsequent increased anterior projection of the nose continued in both males and females after skeletal growth had subsided. However, females had concluded a large proportion of their soft tissue development by age 12 while in males continued growth was noted until age 17 resulting in their having greater soft tissue dimensions for many of the parameters evaluated. During the developmental period, the angular shapes and positional relationships of the nose, lips and chin remained relatively constant for both sexes and was relatively independent of the underlying hard tissues. Treatment planning implications may be drawn from the amounts and timing of the soft tissue development found in this study.

This manuscript was submitted September, 1989.

## Key Words

Soft tissue development • Sexual dimorphism • Treatment planning



**Figure 1a**  
Anatomical points digitized

**Figure 1b**  
Constructed points digitized

the timing, amount, and direction of soft tissue growth in the post-pubertal period.<sup>23-25</sup> Several studies have also suggested that differences develop in the size, thickness and position of the lips and chin between males and females as they approach maturity.<sup>18,25,26</sup>

Studies specifically devoted to nasal growth and development are infrequently found in the literature.<sup>27</sup> Those studies which have focused on the nose have suggested that nasal growth proceeds at a relatively constant rate into adulthood and that the nose increases in relative prominence as maturity approaches.<sup>23,24,28-31</sup> Other researchers approached the problem from a different perspective, trying to determine the effect the underlying hard tissue had on the soft tissue profile of the nose. They have suggested that patients who begin adolescence with a Class II skeletal profile often continue to lack facial harmony as they mature. This lack of harmony is manifested by a greater tendency to develop disproportionate nasal size and an increased prominence of the nasal dorsum.<sup>26,32</sup>

To date, however, there is a lack of quantitative data on the size, shape and relative position of the nose and lips that would enable to the clinician to make rational decisions such as whether or not to extract teeth or plan to retract maxillary incisors in order to obtain the optimum in facial esthetics and harmony. The purpose of this study therefore was four-fold.

1. To determine the course of nasal development.
2. To evaluate the relative position of the nose

in relation to the rest of the soft tissue profile during maturation.


3. To determine the size, shape and position of the lips during maturation.
4. To search for associations between the patterns of hard and soft tissue development.

**Materials and methods**

The sample used in this study consisted of 64 untreated cases from the records of the Bolton study at Case Western Reserve University in Cleveland, Ohio. This sample was comprised of 32 Class I cases (ANB: 2-4 degrees) and 32 Class II cases (ANB: > 5 degrees) (16 males, 16 females in each group). In order to be included in the study, each case had to be of Caucasian origin, with no history of orthodontic treatment, allergies or airway problems. Lateral cephalometric radiographs of each case were evaluated at three time periods: (1) in the early mixed dentition when yearly orthodontic treatment might be considered ( $T_1$ ,  $\bar{x}$ : 7 years, 6 months), so as to provide a baseline for the soft tissue profile; (2) in the early permanent dentition when full orthodontic treatment might be considered ( $T_2$ ,  $\bar{x}$ : 12 years, 5 months), in order to evaluate the changes from  $T_1$  and provide a baseline for comparison to adults; (3) in adulthood ( $T_3$ ,  $\bar{x}$ : 17 years, 2 months), representing a mature soft tissue profile. Each radiograph was traced and then digitized with the assistance of a backlit digitizing table\* interfaced with a minicompu-

\* Hi-Pad Digitizing Tablet, Houston Instruments, Inc., Austin, TX 78753

**Figure 2**  
Antero-posterior nasal measurements and data  
(\* =  $p < 0.05$ , \*\* =  $p < 0.01$ ,  
\*\*\* =  $p < 0.001$ , from the  
previous time period)



ter.\*\* A custom program\*\*\* collated the data and translated the x-y coordinates of each landmark into the desired distances and angles. A total of 46 skeletal, soft tissue and constructed cephalometric points were digitized (Figures 1a and 1b) and a total of 25 cephalometric parameters were evaluated, comprising 21 linear and four angular measurements. Both S-N and Frankfort horizontal were used as reference planes with the large ear rods in use at the Bolton study necessitating the use of a machine porion (Figure 1a, point #5). The anteroposterior skeletal classification was conducted using ANB rather than the "Wits" analysis as the landmarks were more clearly identifiable and were less affected by occlusal changes during development. Ten randomly selected tracings were redigitized to evaluate measurement error which was determined to be insignificant at the  $p > 0.05$  level.

Statistical analysis was conducted on a micro-computer\*\*\*\* using a standard statistical analysis program\*\*\*\*\*. Means, standard deviations and ranges were calculated for each parameter at each time period for each of the four subgroups. Paired Student's *t*-tests using a signifi-

cance level of  $p < 0.05$  were performed to assess the significance of the changes for each subgroup from T<sub>1</sub> to T<sub>2</sub> and from T<sub>2</sub> to T<sub>3</sub>. A three-way analysis of variance was used to search for significant changes between time periods and between subgroups, using the specific factors of sex, Angle class, and time. Tukey's test for multiple comparisons at a significance level of  $p < 0.05$  was then used to identify the source of any significant differences.

## Results

### Antero-posterior growth of the nose (Figure 2)

When measured relative to hard and soft tissue nasion between four and six millimeters of forward growth of the nose was found to occur from age 7 to age 12 years ( $p < 0.01$ ). Males tended to be at the low end of this range while females tended to show the greater amounts of growth. From 12 to 17 years males continued to show four to five millimeters of antero-posterior nasal growth (i.e. one millimeter/year) ( $p < 0.01$ ), while for the females a total of only one to two millimeters of growth was noted ( $p = n.s.$ ). As a result, at age 17 slightly greater amounts of nasal projection were noted in males, with only the Class II males showing significantly greater ( $p < 0.05$ ) nasal projection than Class II females.

When evaluated to measurements reflecting the position of the nose relative to the upper lip, from age 7 to age 12 years all groups demonstrated about two to three millimeters of increased nasal projection. However, from age 12


\*\* Analog Digital Systems, 10820 Comanche N.E., Suite F, Albuquerque, NM 87111

\*\*\* Data General, 4400 Computer Drive, Westboro MA 01581

\*\*\*\* Heath-Zenith 151, Heath Company, Benton Harbor, MI 49022

\*\*\*\*\* SYSTAT. Systat, Inc., 2902 Central, Evanston, IL 60201

**Figure 3**  
**Vertical facial measurements and data**  
 (\*= $p < 0.05$ , \*\*= $p < 0.01$ ,  
 \*\*\* =  $p < 0.001$ , from the  
 previous time period)



to 17 years, while the females continued to grow at a similar rate the males demonstrated a considerable increase in their rate of nasal projection (i.e. four to five millimeters), resulting in a tendency for increased nasal projection in males at age 17.

**Vertical facial development (Figure 3)**

At age 7, males and females had similar total hard tissue total facial heights (N-Me) in the range of 101-103 millimeters. By age 17, the males had outgrown the females by seven millimeters resulting in a greater ( $p < 0.05$ ) male total facial height at 124 millimeters compared to the 117 millimeters total facial height seen in females. However, from T<sub>1</sub> to T<sub>2</sub> both sexes had similar amounts of growth (males +9.3 millimeters, females +10.5 millimeters) while from T<sub>2</sub> to T<sub>3</sub> the males continued to grow rapidly (+11.5 millimeters) even as the females' growth slowed down (+4.5 millimeters). Similar growth patterns were seen in the midface (N - ANS) and the lower face (ANS - Me). Two thirds of the growth seen in total facial height occurred in the lower face which was also responsible for the majority of the greater male facial height seen at T<sub>3</sub>.

When evaluating the comparable soft tissue parameters a much more diverse set of results was seen. For both sexes, from T<sub>1</sub> to T<sub>2</sub>, the soft tissue nasion to subnasale distance demonstrated increases in the five to seven millimeter range which were similar to those seen in the hard tissue N-ANS distance. However, from T<sub>2</sub> to T<sub>3</sub> the males continued to demonstrate similar in-


creases in growth in their soft tissue (+ six millimeters) while the females only grew very slightly (+ one millimeter). As a result, the males soft tissue midface was three to four millimeters longer than that of the females at T<sub>3</sub> (57 millimeters vs 53 millimeters). Similarly the stomion to soft tissue menton distance in males was considerably larger at T<sub>3</sub> (55 millimeters vs 51 millimeters) as a result primarily of greater T<sub>2</sub> to T<sub>3</sub> growth. In contrast were the changes seen in upper lip length (Sn - Sto). Here only a two millimeter overall T<sub>1</sub>-T<sub>3</sub> increase was seen in males (21 to 23 millimeters) and only a one millimeter increase (20 to 21 millimeters) in females, with most of the change occurring primarily from T<sub>1</sub> to T<sub>2</sub>.

**Tissue thickness (Figure 4)**

At age 7 there was a tendency for the soft tissue thickness in the upper lip area (ANS/ SLS and Supradentale/UL) to be one to two millimeters thinner in females than in males. By age 17 this trend had become very clear with females demonstrating an upper lip thickness two to three millimeters less than that of the males. As in other areas, the males and females demonstrated similar growth rates from age 7 to 12 years (males + 2.1 millimeters, females + 2.4 millimeters), but while males continued to grow from 12 to 17 years (+ 2.2 millimeters), females showed only small changes (+ 0.5 millimeters).

The lower lip in females started off at age 7 being 1.6 millimeters thinner than the males (12.8 vs 14.4 millimeters) and ended up at age 17 with a similar deficiency (16.0 vs 17.4 millime-

**Figure 4**  
Soft tissue thickness measurements and data (\* =  $p < 0.05$ , \*\* =  $p < 0.01$ , \*\*\* =  $p < 0.001$ , from the previous time period)



ters). However, the majority of the females' increase of 3.2 millimeters in lower lip thickness occurred from 7 to 12 years while in the males a similar increase (3.0 millimeters) was almost evenly distributed over the two time periods.

In contrast, soft tissue chin thickness in females at T<sub>1</sub> was greater than in males (11.7 vs 10.8 millimeters) but only demonstrated a 1.6 millimeter increase up to age 17, while the males demonstrated 2.4 millimeters of increase in tissue thickness over this period. As a result both sexes had similar (i.e. 13.3 millimeters) soft tissue chin thicknesses at age 17 years.

#### Angular nasal changes (Figure 5)

The nasal bone tended to demonstrate an increase in its anterior projection from age 7 to 17 years as measured to the S-N line (measurement #1). In males a 10 degree increase ( $p < 0.001$ ) was noted, while in females the increase was seven degrees ( $p = n.s.$ ), with both groups showing the changes evenly distributed throughout the evaluation period.

The nasal dorsum (measurement #2) showed a minor tendency to become slightly flatter (one to two degrees,  $p = n.s.$ ) while the columella (measurement #3) became more vertical by three to five degrees ( $p = n.s.$ ) in both sexes from age 7 to 17 when years when measured to Frankfort.

The nasolabial angle showed a tendency to decrease about three to four degrees in both sexes ( $p = n.s.$ ) from age 7 to age 17 with no differences being noted between the Class I and Class II samples.

#### Facial profile to subnasal vertical line (Figure 6)

All five parameters evaluated in this area demonstrated no significant changes during maturation, with no significant differences being seen between the different sexes, or between different skeletal and dental classifications.


#### Discussion

The overall impression gained from the data collected in this study was for both sexes and Angle classes to show quite similar amounts of soft tissue growth from age seven to age 12 years, but for a considerable dichotomy to occur between the sexes from age 12 to age 17 years.

This is clearly evident when examining the absolute amount of antero-posterior growth of the nose and its resultant projection relative to the rest of the profile. In several of the parameters evaluated (Figure 2) the females demonstrated equal, if not slightly greater, amounts of nasal projection at age 7 and experienced slightly more growth up to age 12 than the males (+ five to six millimeters, vs + three to four millimeters) at which age they showed a consistent trend for greater nasal projection. By age 17, while the females' nasal projection had remained virtually constant since age 12, the males had demonstrated continued, and in many cases increased nasal growth (+ four to five millimeters), resulting in a slightly greater degree of nasal prominence in males at age 17 in many of the parameters evaluated.

These soft tissue findings are thus in general agreement with previously published theories of skeletal growth patterns which have sug-

**Figure 5**  
Angular nasal change measurements and data (\* =  $p < 0.05$ , \*\* =  $p < 0.01$ , \*\*\* =  $p < 0.001$ , from the previous time period)



gested that there is a strong tendency for growth in males to continue for several years beyond that seen in females.<sup>18,20-22,26</sup> Where they differ is in suggesting that soft tissue development, particularly in males does tend to lag behind skeletal development and that this finding should be taken into account in the orthodontic treatment planning process. For instance, a clinician evaluating a Class II female at age 12 years could thus reasonably expect that on average only minimal increases in nasal projection will occur over the next two years. However, in a male of similar age any procedure that resulted in upper lip retraction might, in combination with the several millimeters of expected anterior nasal growth, produce a less than optimal final relationship between the lips and nose.

In contrast to the complex sexually dimorphic changes in nasal tip antero-posterior projection, the parameters reflecting the nasal complex's angular characteristics remained virtually unchanged over the 10 year period evaluated in this study (Figure 5). These findings are in agreement with the reports of Sorrell<sup>33</sup> and Farkas, et al.,<sup>34</sup> and would suggest to the clinician that despite the differences in nasal growth, the overall contours of the nose, and in particular the clinically significant naso-labial angle, are not likely to change by more than three to four degrees as a child matures. The only exception to these findings was in the angulation of the nasal bone which showed a strong tendency to become more horizontally inclined with time. This tendency has been reported previously with the T<sub>3</sub>


projection being up to 10 degrees more horizontal than at T<sub>1</sub>.<sup>28,32</sup> It has been suggested that greater elevation and projection of the nasal bone is seen in Class II cases resulting in a dorsal nasal hump occurring more frequently in Class II cases.<sup>24,28</sup>

The data from this study agrees with these findings showing both a more anterior position of the nasal dorsum and a more anteriorly angulated nasal bone in the Class II cases.

The tendency for females to have smaller soft tissue dimensions than males was clearly demonstrated in the relative thicknesses of the soft tissue of the lips and chin (Figure 4). Females demonstrated thinner lips at age seven and while growing similar amounts to males up to age 12, the final lip thicknesses in females were often less by two millimeters or more at age 17 due to almost no increase occurring in females from age 12 to 17 years. This data tends to confirm the findings of Subtelny,<sup>18</sup> Mauchamp and Sassouni,<sup>25</sup> as well as Riolo, et al.<sup>26</sup> Thus, a clinician retracting upper incisors in a 12 year female might expect little compensatory lip growth, while in a male a less detrimental facial effect might be expected if the normal two millimeter increase in upper lip thickness occurred from age 12 to age 17 years.

When evaluated relative to the subnasale vertical reference line, the components of the soft tissue profile below the nose (e.g.: superior labial sulcus, upper lip, and soft tissue chin) showed remarkably little change over the seven to 17 years age period (Figure 6). This data seems to

**Figure 6**  
Subnasale vertical line measurements and data  
(\* =  $p < 0.05$ , \*\* =  $p < 0.01$ ,  
\*\*\* =  $p < 0.001$ , from the  
previous time period)



confirm the overall impression gained that the components of soft tissue profile mature relatively similarly, although the timing and amount of change is very different in males and females.<sup>34</sup> The data from the subnasale vertical reference line would tend to suggest to the clinician that the relative positions of the upper lip and soft tissue chin are likely to remain constant with maturation of the child and that this should be taken into account in treatment planning. The absolute values seen at age 17 of the Class I and Class II cases were very similar to those seen in Spradley's study<sup>35</sup> of ideal faces with the exception of the more retrusive soft tissue chin seen in the Class II cases in this study.

As this study was concluded at age 17 the potential for future soft tissue development should not be ruled out. However, as Sarnas and Solow have demonstrated, any future changes are likely to be small, averaging less than 0.5 millimeters or 0.5 degrees.<sup>36</sup>

### Conclusions

On the basis of cephalometric records from a sample of 64 untreated persons examined in the mixed dentition, early permanent dentition and early adulthood, the following conclusions were reached:

1. Antero-posterior growth and increased anterior projection of the nose continued in both

males and females after skeletal growth had subsided.

2. Females had concluded a large proportion of their soft tissue development by age 12 while in males growth continued until age 17 resulting in greater soft tissue dimensions in many parameters.
3. The angular shapes and positional relationships of the nose, lips and chin remained relatively constant throughout the developmental period for both sexes.
4. No relationships were found between the amount of nasal development and skeletal class or sex of the subjects. The growth observed was relatively independent of the underlying skeletal hard tissue.
5. The sexual dimorphism in the amount and timing of soft tissue development may have treatment planning implications.

### Acknowledgments

The authors would like to express their gratitude to Dr. B. Holly Broadbent of the Bolton Study for allowing them access to the patient records; to Dr. Jim Economides, Paul Economides and Dr. Martin Wagner for their assistance in the statistical and computer portion of this project; to Dr. Wick Alexander and Dr. Joe Jacobs for their advice and guidance; and to Mrs. Martha Black for typing this manuscript.

**Author address**

Peter M. Sinclair  
Orthodontic Department  
School of Dentistry  
University of North Carolina  
Chapel Hill, North Carolina 27599-7450

*J.S. Genecov is an orthodontist in private practice in Dallas, Texas. This paper is based on a thesis by Dr. Genecov, submitted*

*to the Department of Orthodontics, Baylor College of Dentistry, Baylor University, in partial fulfillment of the requirements for the degree of Master of Science in Dentistry.*

*P.M. Sinclair is an associate professor in the Orthodontic Department at the University of North Carolina and formerly was an Associate Professor at Baylor College of Dentistry.*

*P.C. Dechow is an assistant professor in the Department of Anatomy at Baylor College of Dentistry.*

**References**

1. Ricketts, R.M., Bench, R.W., Hilgers, J.J., Schult-hof, R.: An overview of computerized cephalometrics. *Am. J. Orthod.*, 61:1-28, 1972.
2. Steiner, C.: Cephalometrics in clinical practice. *Angle Orthod.*, 29:8-29, 1959.
3. Tweed, C.H.: Indications for the extraction of teeth in orthodontic procedures. *J. Orthod. Oral Surg.*, 30:401-60, 1944.
4. Björk, A.: Prediction of mandibular growth rotation. *Am. J. Orthod.*, 55:585-94, 1969.
5. Downs, W.B.: The role of cephalometrics in orthodontic case analysis and diagnosis. *Am. J. Orthod.*, 38:162-82, 1952.
6. Harvold, E.: Morphogenetic response to activator treatment. *Am. J. Orthod.*, 60:478-90, 1971.
7. Jacobson, A.: The Wits appraisal of jaw disharmony. *Am. J. Orthod.*, 67:125-38, 1975.
8. Reidel, R.A.: An analysis of dentofacial relationships. *Am. J. Orthod.*, 43:103-19, 1957.
9. Holdaway, R.A.: A soft tissue cephalometric analysis and its use in orthodontic treatment planning. Part I. *Am. J. Orthod.*, 84:1-28, 1984.
10. Holdaway, R.A.: A soft-tissue cephalometric analysis and its use in orthodontic treatment planning. Part II. *Am. J. Orthod.*, 85:279-93, 1984.
11. Merrifield, L.: Profile line as an aid in critically evaluating facial esthetics. *Am. J. Orthod.*, 52:804-21, 1966.
12. Angle, E.H.: Treatment of malocclusion of the teeth and fractures of the maxilla. 6th ed. S.S. White Dental Mfg. Co., Philadelphia, 1900.
13. Peck, H., Peck, S.: A concept of facial esthetics. *Angle Orthod.*, 40:284-317, 1970.
14. Shaw, W.C.: The influence of children's dentofacial appearance on their social attractiveness as judged by peers and lay adults. *Am. J. Orthod.*, 79:299-314, 1981.
15. Burstone, C.J.: The integumental profile. *Am. J. Orthod.*, 44:1-25, 1958.
16. Brodie, A.G., Downs, W.B., Goldstein, A., Myer, E.: Cephalometric appraisal of orthodontic results: a preliminary report. *Angle Orthod.*, 8:261-65, 1938.
17. Fishman, L.S.: A longitudinal cephalometric study of the normal cranio-facial profile, utilizing a proportional analysis of skeletal, soft-tissue and dental structures. *Int. Dent. J.*, 19:352-79, 1969.
18. Subtelny, J.D.: A longitudinal study of soft-tissue facial structures and their profile characteristics defined in relation to underlying skeletal structure. *Am. J. Orthod.*, 45:281-507, 1959.
19. Björk, A., Skieller, V.: Facial development and tooth eruption. *Am. J. Orthod.*, 62:339-82, 1972.
20. Bishara, S.E.: Longitudinal cephalometric standards from five years to adulthood. *Am. J. Orthod.*, 79:35-44, 1981.
21. Bishara, S.E., Hession, T.J., Peterson, L.C.: Longitudinal soft-tissue profile changes: a study of three analyses. *Am. J. Orthod.*, 88:209-23, 1985.
22. Bishara, S.C., Peterson, L.C., Bishara, E.C.: Changes in facial dimensions and relationships between the ages of five and 25 years. *Am. J. Orthod.*, 85:238-52, 1984.
23. Posen, J.M.: A longitudinal study of the growth of the nose. *Am. J. Orthod.*, 53:746-56, 1967.
24. Chaconas, S.J.: A statistical evaluation of nasal growth. *Am. J. Orthod.*, 56:403-14, 1969.
25. Mauchamp, O., Sassouni, V.: Growth and prediction of the skeletal and soft tissue profiles. *Am. J. Orthod.*, 64:83-94, 1973.
26. Riolo, M.L., Moyers, R.E., TenHove, T.R., Mayers, C.A.: Facial soft tissue changes during adolescence. In: *Craniofacial Growth During Adolescence*. Edited by David S. Carlson and Katherine A. Ribbens. University of Michigan Center for Human Growth and Development, Ann Arbor, pp 117-33, 1987.
27. Meredith, H.V., Knott, V.B., Hixon, E.H.: Relation of the nasal and subnasale components of facial height in childhood. *Am. J. Orthod.*, 44:285-94, 1958.
28. Clements, B.S.: Nasal imbalance and the orthodontic patient. *Am. J. Orthod.*, 55:244-64, 1969(a).
29. Clements, B.S.: Nasal imbalance and the orthodontic patient. *Am. J. Orthod.*, 55:329-52, 1969(b).
30. Clements, B.S.: Nasal imbalance and the orthodontic patient. *Am. J. Orthod.*, 55:477-98, 1969(c).
31. Neger, M.: A quantitative method for the evaluation of the soft-tissue facial profile. *Am. J. Orthod.*, 45:738-51, 1959.
32. Robinson, J.M., Rinchuse, D.J., Zullo, T.G.: Relationship of skeletal pattern and nasal form. *Am. J. Orthod.*, 89:499-506, 1986.
33. Sorrell, M.: A study of the heritability of the soft tissue nasal profile. Master's Thesis, University of Pittsburgh, 1982.
34. Farkas, L.G., Sohm, P., Kolar, J.C., Munro, I.R.: Inclinations of the facial profile: art versus reality. *Plast. Reconstr. Sur.*, 75:509-19, 1985.
35. Spradley, F.L., Jacobs, J.D., Crowe, D.P.: Assessment of the antero-posterior soft-tissue contour of the lower facial third in the ideal young adult. *Am. J. Orthod.*, 79:316-25, 1981.
36. Sarnas, K.V., Solow, B.: Early adult changes in the skeletal and soft-tissue profile. *Eur. J. Orthod.*, 2:1-12, 1980.