

Diabetes in old age, a review

Abbaszadeh Ahranjani S, Tabatabaei Malazy O, Pajouhi M*

Endocrinology and Metabolism Research Center, Tehran University of Medical Sciences, Tehran, Iran

Abstract

The elderly population is usually defined as those at least 60 or 65 years of age. In tandem with the rest of the world, Iran has seen a dramatic increase in this age group in recent decades. It is estimated that the percentage of men and women over 65 years old in 2020 will be 9.4% and 9.1%, respectively. The old are liable to suffer from coincidental diseases; decisions as to the most appropriate treatment modality should, therefore, be made on an individual basis in harmony with the physical and mental status of the patients as well as possible drug interactions. Diabetes, together with its complications such as hypoglycemia, is one of the most prevalent chronic diseases and its prevalence rises considerably by aging. Educating patients and their families empowers them to maintain their independence in the management and treatment of their medical condition. In addition, regular ophthalmologic, cardiovascular, and other laboratory examinations can reduce treatment costs and burdens and confers a better overall quality of life. Given the increase in the aged population around the globe and in Iran and the relative paucity of data on diabetes in this age group, it seems advisable that more attention be paid to raising awareness of the problems of diabetes in this age group.

Keywords: Diabetes, Elderly, Education

Introduction

The elderly are the symbols and a treasure trove of a country's customs and traditions. The cultural-religious context of our country, Iran, demands special reverence for the aged, not least during debilitating diseases.

Diabetes deteriorates lifespan by imposing damages on multiple vital organs [1]. In 1997, it was estimated that 124 million people were afflicted with this metabolic disorder worldwide, the majority (97%) of them had type2 diabetes [2]. Unfortunately, it estimates that the number of diabetics in the adult population (aged ≥ 20 years old) to have risen to 300 million by the year 2025 [3]. In the Eastern Mediterranean Region (EMRO), which includes such countries as Saudi Arabia, Islamic Republics of Iran, Kuwait and Iraq, the prevalence of diabetes amongst the adult population was 14.5% in the year 2003 [4]; simultaneously, the prevalence of the diabetic population in Iran stayed at 5% [5]. Currently, it is estimated that there are 22 million diabetic patients out of total 290 million adult populations in the EMRO; the feature seems to elevate to 300 million by the year 2025 [4].

The elderly comprise a portion of the general population that are at least 60 or 65 years of age according to different definitions [6]. In the next fifty years, the elderly may account for 2 billion people in the world [7]. The rapid population growth in the developing world by comparison with developed countries means that the former has much less time to devise suitable programs for the prevention of diabetes [7].

Diabetes and its consequent complications place a very heavy burden on health care systems. Abolhasani et al. revealed that the burden of diabetes and its complications in term of the Disability-Adjusted of Life Years (DALYs) was 306,440 years in Iran in 2001 [8].

In short, diabetes mellitus is a common disorder in the elderly population and

imposes a great burden on individuals and health care systems. The present study sought primarily to probe into diabetes in old age using such keywords as type2 diabetes mellitus, elderly, diabetes complications, and treatment of diabetes.

Epidemiology

Age is one of the variables influencing the prevalence of diabetes and Impaired Glucose Tolerance (IGT) [1]. For instance, in those 45 – 55 years of age in the USA, diabetes is four times more common than in the ones in the 20 – 44 aged. [1] Currently, at least 20% of people aged over 65 years old are estimated to be suffering from diabetes [9], and the figure amounts to 50% in some races such as Hispanics [10]. On the other hand, the world's elderly population is on the increase [11]. Different epidemiological studies have reported varying diabetes prevalence in the elderly, with the reasons cited being different in terms of not only the dates of the studies but also the diagnostic criteria and the racial groups recruited into the studies [12]. A study, conducted between 1993 and 2001 on old people (≥ 67 years), evaluated the prevalence and incidence of diabetes mellitus in four racial groups and reported that Asians had the highest prevalence. In that study, the prevalence of diabetes raised alongside aging and reached its peak amongst those in the 75-79 aged before it started to drop [13]. It should be noted, however, that the highest prevalence of diabetes mellitus tends to occur in the oldest age groups in the majority of the world's populations [2].

Until 1979, fasting blood glucose ≥ 140 mg/dl was the standard diagnosis of diabetes. In 1997, this standard was lowered to 126 mg/dl, which accounts for the lower epidemics having been reported in published epidemiological data since then. In the Japan 2002 Survey, a total of 7.4 million people were suspected to suffer from diabetes, 2.26 million of them being in the age bracket of 60 and 2.51 million being

at least 70 years old [14]. The annual rate of diabetes appearance amongst individuals over 65 years old in the 1994-1995 time period in the USA was about 23% before it soared to 62% in 2003-2004 [15, 16]. The total prevalence of diabetes in adult groups (≥ 20 years) in Iran in 2001 was reported at 4.67% (1.6 million patients). In the same year, the rate of diabetes in those aged between 60 and 69 years and those aged over 70 years was 15.49% and 14.07%, respectively. The average rate of diabetes in the elderly population (≥ 65 years old) was 14%, with the prevalence in women being higher than that in men in all age groups [5]. In the Tehran Lipid and Glucose Study (TLGS) between 1999 and 2001, 1799 persons from a total number of 15005 participants were 60 years old or higher. In this population based study, the prevalence of diabetes and IGT was 26.2% and 20.8% in the women and men, respectively [17]. In a study conducted on 1694 elderly persons over 60 years of age in the Iranian city of Isfahan in the year 2004, 17.4% suffered from diabetes and there was a significant difference in terms of diet between the diabetics and non-diabetics [18].

The time period between the years 1980 and 2003 life expectancy in Iran increase from 55 years to 68 years in men and to 70 years in women; this change is a testament to the further aging of the population and also the increase in non-communicable diseases such as diabetes mellitus in Iran [19]. A population based survey in the year 1986 showed that old people accounted for 3.03% of the entire Iranian population. This figure rose to 4.32% in 1996 and 5.18% in 2006 [20]. In the Shieber et al. study in 1995, men and women over 65 years of age accounted for 6% and 5.7% of the total study population, respectively. The authors predicted that by 2020, those percentages would have risen to 9.4 and 9.1% in men and women, respectively [21].

Of the factors contributing to a higher prevalence of diabetes in old age are the presence of other chronic diseases, higher

"hope for life", [22] lower activity level, higher prevalence of obesity due to development of technology and westernized diet, availability of healthy and therapeutic facilities, and more efficient laboratory evaluations of high blood sugar. In both genders, the prevalence of obesity in middle age makes the person susceptible to diabetes [14, 22].

A major health issue, diabetes has adverse effects on the quality of life and life span [4]. The fact that diabetes worsens with advancing age means that it creates even more problems in elderly people [23].

Senile diabetes pathophysiology and etiology

Insulin resistance and IGT are more prevalent in the aged than in young people, and the former are more susceptible to have type2 diabetes. Some reasons for lower sensitivity to insulin in the elderly are: adiposity, changing in fat distribution, decrease in free fat mass, abnormal muscle content, bad food habits, and lack of physical activity. Insulin resistance and defects in insulin secretion in response to glucose are the main contributing factors for IGT or diabetes in old obese and old slim persons, respectively. Also, autoimmune changes like antibody against islet cells as well as insulin deficiency may also be responsible [24, 25].

Type2 diabetes is divided into two categories according to the time of appearance: one that starts from middle age and has the most prevalence (55.28%) and another that commences after 65 years of age with a 44.72% prevalence rate. In rare cases, a continuation of juvenile diabetes (type I) is also seen in old age [26]. The etiopathology of diabetes in the elderly is complex and multi-factorial and it can happen due to a lower function of the islets of Langerhans because of partial vein damage to the pancreas or ramification of general atherosclerosis process [27].

Diagnosis of diabetes in the elderly

Natural differences in the physiology of the body are the corollaries to aging and exert a direct bearing upon the signs and symptoms of diabetes and are likely to cause problems in the diagnosis or treatment of diabetes. As a consequence, diabetes does not tend to present with gross hyperglycemia in elderly individuals. Moreover, at least 50% of elderly persons are oblivious to their diabetes [11]. The reasons, albeit not very clear, may be attributable to the unawareness of health care providers about the criteria for diagnosis of diabetes, absence of classic signs of diabetes, or the increase in the urinary extraction limit of glucose secondary to aging [11]. For instance, polydipsia in the elderly caused by a decline in thirst perception is less common than that in other age groups, rendering dehydration more prevalent. Hyperglycemia in these people can be accompanied with such symptoms as dizziness and incontinence. In old people, autonomic alerting signs set off by a decrease in glucagon and epinephrine, moreover the secretion of growth hormones in response to hypoglycemia decreases. Conversely, sometimes the epinephrine response increases and may induce severe hypoglycemia [25]. It is, therefore, advisable that blood tests be included in routine medical assessments of the aged.

Standard laboratory criteria for the diagnosis of diabetes in old people are similar to those in other age groups: blood glucose <100 mg/dl is normal and >126 mg/dl is indicative of diabetes. Blood glucose of 100-126 mg/dl is not regarded as diabetes, but it is called IGT and an individual with that amount of blood glucose is usually susceptible to some diabetic complications, especially, cardiovascular diseases. After the age of 50; for each decade that passes, fasting blood sugar and 2-hour postprandial blood sugar in the GTT (Glucose Tolerance Test) are liable to increase by 1.08 and 9 mg/dl, respectively [24].

Diabetes complications in the elderly

Geriatric syndromes

Cognitive disorders, depression, chronic pains, polypharmacy, injuries from falling, and difficulties in urination are some of the geriatric symptoms [28]. The risk of these syndromes in diabetics is higher than those in non-diabetics [29]. Depression and psychosocial stressors are more common in diabetics than in non-diabetic groups [30, 31]. A cross-sectional study recruited 2522 old men and women between 70 and 79 years old for an investigation into the relationship between diabetes and depression and found that the probability of depression symptoms was 23.5% and 19% in the diabetics and non-diabetics, respectively, and the rates of relapsing were 8.8% against 4.3%. The highest relation between diabetes and relapse of depressive symptoms was for the most part observed in the patients with weak blood sugar control [32]. Depression is one of the salient contributors to movement disability [33]. On account of the fact that diabetics are at higher risk of suffering from major depression, necessitates screening measures at the beginning of the diagnosis (i.e. first 3 months) or whenever it appeared a change in the patients' clinical conditions [28]. Cognitive problems and dementia are more prevalent in elderly diabetics than non-diabetics [34]. In an examination of cognitive disorders, in a case-control study in Tehran, with 160 type 2 diabetics and non-diabetics recruitment, the average values in Mini Mental Status Examination (MMSE) in the case group were less than those of the control group and a significant reverse correlation between these values, age, and duration of diabetes was reported. This report showed that old diabetics had some degrees of cognitive problems, which worsened with advancing age [35]. Functional disability is defined as difficulty in performing activities of daily living and social activities [33]. Diabetes can aggravate the disability and delay the

patients' recovery, as a consequence, leads to a higher prevalence of disability. This relation magnifies with the appearance of diabetes complications in the eye and diabetic foot, related conditions like obesity and depression or high blood pressure, and such social issues as lower education levels and lower income [29]. Not only will all these factors bring about higher rates of hospitalization and stays at nursing homes, but they will also increase need for care and the burden of diabetes in this age-group [29, 36].

Cardiovascular diseases

Both diabetes and aging tend to exacerbate cardiovascular diseases with symptoms ranging from completely silent and without any sign to obviously overt diseases [37]. Without considering the influence of age on diabetes, the most common outcome of type2 diabetes is emerging fatal and non-fatal vascular complications such as cardiovascular diseases and peripheral vascular or cerebrovascular conditions [38, 39]. In a study conducted over an eleven-year period, death due to diabetes in the old people with recent diagnosis was 9.2% higher than that in the non-diabetics. Also, 91.8% of the diabetic patients had more micro and macro vascular complications compared to 72% of the non-diabetics, and the diabetics were at higher risk of lower limb complications [40].

The metabolic syndrome is a disorder that is normally accompanied with central obesity, hypertension, and abnormal glucose tolerance test and lipid profile test results. This syndrome is a precursor to such serious diseases as cardiovascular diseases and diabetes. In one study carried out on 12600 persons in Iran, the incidence of the metabolic syndrome in the urban and rural populations was 25.4% and 21.7%, respectively. Additionally, the highest incidence (68.9%) was found amongst those over 60 years old in the rural areas of Isfahan, Iran [41].

The rate of heart failure in diabetic patients with respect to age as a major risk factor has been reported to be twice as the rate in non-diabetics [42]. The rate of heart failure in diabetics is also believed to be related to the quality of blood sugar control in as much as for each percentage increase in HbA1c¹, the risk of heart failure will increase by 15% [43]. According to the ACCORD² study, the mortality rate in the diabetics with HbA1c>6.5% tends to increase, especially in the elderly with prior history of cardiovascular diseases [44].

In a prospective study aimed at examining random cases with heart failure, the rates of having heart failure in a four-year period for men and women (≥ 65 years old) were 15% and 18%, respectively. Twenty years later and with the same people and time period, the respective percentages increased to 22% and 24% [45].

The treatment of hypertension and dyslipidemia in elderly groups can reduce the morbidity and mortality of cardiovascular diseases [46]. The goal is to control blood pressure at $\leq 130/80$ mmHg, LDL³ at <100 mg/dl, and total cholesterol / HDL⁴ ratio at < 4 in elderly groups [50]. In individuals with a history of cardiovascular diseases, it is usually recommended that LDL be reduced to levels lower than 70mg/dl [47].

Hypoglycemia

Hypoglycemia is a severe and prevalent consequence in the treatment of diabetes in old people and is defined as the Whipple triad, i.e. presence of clinical symptoms resulting from a drop in the blood sugar level followed by a decrease in the blood sugar level (less than 50 mg/dl) and symptoms fading after the blood sugar level returns to the normal mode [48]. Warning symptoms are dizziness or changing level of consciousness, pallor, sweating,

1 -Glycosylated Hemoglobin A1C

2 -Action to Control Cardiovascular Risk in Diabetes

3 -Low Density Lipoprotein

4 -High Density Lipoprotein

difficulty concentration, irritability, and paresthesia around the lips, low body temperature, nightmares, and insomnia. Ramifications of hypoglycemia in old people are more severe due to weaker physical strength and macrovascular diseases. On the other hand, after hypoglycemia, patients are at higher risk for macrovascular incidents like myocardial infarction and brain stroke. The occurrence of several and sudden hypoglycemia attacks in old people triggers dizziness, weakness, temporary unconsciousness and fall, undermining the patient's confidence and sense of independence. Accordingly, a desirable blood sugar level in old people is set at higher levels than that in the young [49].

Unawareness of hypoglycemia is widespread. Approximately, one fourth of insulin-administered patients may experience hypoglycemia without any alerting sign. Hypoglycemia is likely to be followed by changes in the brain blood circulation; that may not distinctly manifest itself in old patients because of hard vessels and nervous system disorders. An increase in the duration of diabetes concurs with rise in the probability of unawareness of emerging hypoglycemia [49].

Diabetic foot

According to a reported study in Iran in the year 2001 [8], the incidence diabetic foot in those over 20 years old was 3%. Another study looked into the rate of lower limb amputations and hospital stays due to diabetic foot in 281 patients hospitalized in Imam Khomeini Hospital and Dr Shariati Hospital, two educational hospitals in Tehran, between 1979 and 2001 and following findings reported: the amputation rate dropped from 40% in 1995 to 14% in 2001 and the average hospital stay was 4 weeks [50].

Like young people, old patients are not spared from diabetic ulcers. Some of the principal causes of this symptom are sensory neuropathy [51], foot deformities, peripheral arterial diseases, sight weakness, low activity, or presence of other

diseases. The risk of other major amputations (proximal amputations to the foot heel)[52] due to aging and increase in visual loss may intensify as well. Moreover, sight weakness leads to not daily foot checking by old patients [53]. For prevention of diabetic foot ulcers, an identification of high risk people, avoiding wearing unsuitable shoes, and not walking with barefoot are usually suggested. Peripheral vascular disease is one of the important leading factors for amputations which can be prevented by carefully long-term blood sugar and dyslipidemia control, treatment of hypertension, and smoking cessation [54-56]. It is recommended that diabetic foot be managed by a multidisciplinary team approach encompassing general practitioners, nurses, endocrinologists, vascular surgeons, orthopedists, and physiatrists [57-59].

Visual problems

Visual impairments tend to increase amongst the old to the extent that one out of three elderly persons suffers from some form of impairment [60]. The most severe visual impairments are age-related macular degeneration, cataract, glaucoma, and diabetic retinopathy [61]. After age-dependent degeneration in the macula, cataract, and glaucoma, diabetes is the fourth common cause of blindness in the elderly and the first in the West and in people under 60 years of age [62]. Diabetic retinopathy is distinct with microvascular injuries in the retina [63], with its prevalence in Asia reported from 10.5% in India to 35% in Taiwan [64].

There are only a few epidemiological studies in the existing literature on diabetic retinopathy in the aged. In Afkhami et al. study of 590 patients with type 2 diabetes referring to the Yazd Diabetes Clinic, a city in centre of Iran, 39.3% of the study population suffered from diabetic retinopathy and age, body mass index, glucose level, and HbA1c were amongst the majority of

risk factors for diabetes-related visual impairments [65].

Diabetic retinopathy is evident in 15% of diabetics over 70 years old at the time of diagnosis [66]. The prevalence of diabetic retinopathy is highly dependent upon the duration of diabetes. In a study of people over 65 years of age with an average diabetes background of 15 years, while the prevalence of diabetic retinopathy was 43%, the prevalence observed in 23% of people in the same group but with a shorter history [67].

Increase in age, diabetes duration, and the intensity of retinopathy (less important) are the main risk factors for developing cataract. In diabetics, the clinical appearances of cataract are not specific to diabetics [62]. Although aging is the main factor for appearance of glaucoma, no relation between diabetes and glaucoma has been mentioned in numerous studies [62, 68]. Old diabetics who have recently been diagnosed, must be underwent ophthalmologic examination [28, 62]. Old diabetics who are at higher risk of suffering eye disease (e.g. having eye symptoms, retinopathy, glaucoma and cataract in the first examination or during the last two years, HbA1C $\geq 8\%$, type 1 diabetes, and blood pressure $\geq 140/80$ mmHg) must be examined annually by an ophthalmologist [28, 62]. For low-risk people (e.g. recent diabetes diagnosis, with no diabetic retinopathy, controlled blood sugar, and no diabetes adverse effects) examining at least once every two years is suggested [28, 62]. Regular visual assessments can assist elderly persons in maintaining their life activities and self-sufficiency [61].

Like younger diabetics, regularly control of plasma sugar and blood pressure in order to decrease in incidence and progress to diabetic retinopathy is crucial and HbA1c must be kept at an acceptable level. Needless to say, the probability of hypoglycemia and blood pressure drop must not be neglected in the aged, hence

the advisability to treat everyone individually [62].

Diabetic nephropathy

This complication is characterized as microalbuminuria ≥ 30 mg/day. If albuminuria amounts to ≥ 300 mg/24h, it is referred to as clinical nephropathy or obvious albuminuria [69]. Diabetic nephropathy has no prominent symptoms until advanced stages of the disease. This condition is aggravated when accompanied with such problems as anemia or cardiac failure. The detection of proteinuria and diabetic nephropathy necessitates routine screening [70-72].

For the majority of elderly persons, the presence of macrovascular diseases, apart from nephropathy, is a great problem. The clinician needs to take heed of the patients' age and serum creatinine when evaluating the patients' albuminuria. The general approach to diabetic nephropathy in the elderly is similar to that in young diabetics and includes controlling blood pressure, blood sugar, and lipid profiles, administering angiotensin-converting enzyme inhibitor drugs, and periodically assessing serum creatinine and urine albumin. It is recommended that precautions be taken when treating patients with high levels of serum creatinine. For example, the use of metformin is generally contraindicated and the use of glibenclamide should be closely monitored because of the tendency of the drug to cause prolonged hypoglycemia [70]. In one study in Iran, 123 diabetics with respect to macroalbuminuria were evaluated, accordingly, 20.3% of the patients had microalbuminuria, 61.1% normoalbuminuria, and 10.6% macroalbuminuria. There was a significant statistical difference between the mean age of those with microalbuminuria (58.8 years) and those with normoalbuminuria (50.3 years), and the renal function in the patients with type 2 diabetes progressively deteriorated with

advancing age and the duration of diabetes [73].

Oral and teeth diseases

Oral and teeth diseases are regarded as one of the frequent consequences of diabetes. These diseases are more severe and prevalent in diabetics than in the healthy population. Diabetics are more susceptible to some clinical manifestations of oral diseases like candidiasis, which weakened the control of blood sugar. Appearance of xerostomia due to an increase in glucose levels in oral secretions and dysfunction of local immune mechanisms, is usually observed [74]. In an examination of diabetes-related causes of oral symptoms in 116 diabetics, the most common symptoms were gingivitis (96%), periodontitis (90%), and mouth dryness (67%). Although there was no statistically significant relationship between diabetes-related causes of mouth symptoms and blood glucose control, the patients with good blood glucose control had a lower incidence of the foregoing diseases [75].

Sexual dysfunction

Sexual dysfunction is seen in both types of diabetes (types 1 and 2). In elderly women, diabetes alongside with menopause, can cause vaginal dryness, low resistance to normal vaginal flora infections, and high susceptibility for fungal infections [76, 77]. Erectile dysfunction (impotency) is the most common sexual dysfunction in diabetic men regardless of age, and its prevalence increases with advancing age [78]. It has been reported that the prevalence of erectile dysfunction in male diabetics is 35-75%, which rises to 55-95% after 60 years old. Etiology of erectile dysfunction is multifactorial, and neuropathy, vascular insufficiency, and failure to control of blood sugar are some of frequent risk factors [76].

Mofid et al. study of 700 diabetic men between 20 and 69 years of age in Tehran

[79] reported that 246 (35%) persons suffered from erectile dysfunction according to the International Index of Erectile Function (IIEF) and this figure rose to 43% in those over 60 years old. Erectile dysfunction is more frequently observed with advancing the duration of diabetes and unfavorable blood glucose control.

Treatment of diabetes in the aged

The principles of the treatment of diabetes in the elderly are similar to those in the young; the difference lies in the changes taking place in insulin pharmacokinetics, also, its absorption, metabolism, and clearance are liable to changes. Furthermore, alongside diseases, onset of diabetes, presence or absence of microvascular and macrovascular complications, and varying physical and emotional problems of old people are all reasons justifying to make the best decision on drug regimens on an individual basis [25, 80].

Most of the diabetic patients are suffering from accompanied diseases and up to 70% have more than two comorbid conditions [81, 82]. Moreover, regular evaluation, management, and follow-up of complications such as hypertension, obesity, dyslipidemia, and cardiovascular diseases are similar as those in youth [83]. Nonetheless, clinicians must be cognizant of the dominant features of the disease for choosing appropriate treatment modalities.

Self-care is one of the most important aspects of diabetes management. Patients should adopt a healthy life style as regards diet and physical activities and have sufficient knowledge about the disease and its management; thus, the significance of education as an important factor in blood glucose control is unfolded [84]. Diabetic patients and their families or caregivers should be equipped with the knowledge on the symptoms, diagnosis and management of hyper and hypoglycemia. Education in this regard includes knowing the causing

factors, early warning signs and symptoms, prevention, diagnosis and control of the symptoms, treatment modalities, use of the glucometer, and appropriate scheduling for seeking professional help [28]. Usually, patients who have been diagnosed at middle-age are more familiar with appropriate management of diabetes than those of the elderly who have recently been diagnosed.

The management of diabetes in old age based on three fundamentals: nutritional changes, physical activity, and medications if the desired blood sugar level can not be achieved.

Eating three regular meals in addition to at least two snacks between meals reduces the risk of a drop in the blood glucose level. It is, therefore, important that patients be educated for healthy lifestyle and regular physical activities alongside with routine medication [27]. The elderly diabetics should regularly receive consultations about dieting and weight reduction [28, 85]. The consumption of vitamin C, vitamin E, and minerals (zinc, magnesium, and complements) may improve blood glucose control in selected groups of elderly diabetics [86-89]. Several investigations have demonstrated that physical activity can significantly promote control of blood glucose as well as lipids in elderly diabetics [90-92]. Needless to say, the intensity of physical activity should correspond to the health and physical condition of the patient [93]. Vigorous exercise is not usually recommended for elderly diabetics because of the possibility of cardiovascular complications [94].

Primary or secondary failure of taking oral hypoglycemic agents is conceivable in elderly diabetics who have diagnosed during middle-ages. In addition, oral agents are not markedly efficacious in senile diabetes when administering early in the course of treatment; moreover, their efficacy tend to decrease over the time. Indeed, no desirable response can be observed with oral hypoglycemic

agents in most elderly diabetics and majority end up using insulin as the last available therapeutic choice. Different treatment modalities have various effects on the quality of life. Occasionally, insulin prescription may improve as quality of life [27]. In one study, the patients under treatment with insulin had less satisfaction in life and worse general condition than did those receiving medical-nutritional treatments. This difference, however, was not significant when taking into account the duration of diabetes [94]. In some patients, insulin treatment hastens need for institutionalization [29]. Another study, however, reported more satisfaction and improvement in the general condition of diabetics between 39 and 81 years old after 7 months of insulin therapy [95].

The optimal levels of plasma glucose vary in the elderly because of possible accompanied diseases, drug interactions, and occurrence of some unpredictable conditions. Accordingly, the goal in the management of diabetes in the elderly is to maintain an appropriate level of blood sugar without increase in the risk of hypoglycemia. For effectively management of diabetes in the elderly, a multidisciplinary approach is warranted; moreover, endeavors must target efficiently lifestyle changes. Several studies have shown that the foregoing approach can markedly improve blood glucose control, enhance effectiveness of treatments, and improve the quality of life in elderly patients [96, 97]. For monitoring the long-term control of blood sugar in old patients, HbA1c assessment at least every six months is recommended [98]; the goal is to attain HbA1c of 6.5 to 7.5% in the elderly with no coincident diseases and 7.5 to 8.5% in ones who suffering from accompanied conditions. A desirable fasting blood sugar level is 90-126 mg/dl [27].

It is worthy of note, however, that the costs of hospitalization for elder diabetics estimated 60% higher than those for non-

diabetics [99]. Table 1 presents a list of oral hypoglycemic agents for the treatment of type2 diabetes [63, 100], and Table 2 introduces various products of insulin which are frequently used in Iran [100]. Given the significantly increase in the aged population around the world and in Iran and the importance of maintaining

the dignity of this age group, it is judicious to separately notice their problems.

Taken together, it seems advisable that more attention be paid to raising awareness of the problems of diabetes in this age group.

Table 1. Oral hypoglycemic agents for type 2 diabetes mellitus

	Sulfonylureas (generation2: Glybenclamide*)	Meglitinides (Repaglinide*)	Biguanides (Metformin*)	Thiazolidinediones (Pioglitazone*)	Alpha-glucosidase inhibitors (Acarbose*)
Mechanism	Stimulate insulin release from beta cells of pancreas	Stimulate insulin release from beta cells of pancreas	↓hepatic neoglucogenesis, ↑ insulin uptake in peripheral muscles	Enhance sensitivity to insulin in liver & muscle	Reversible inhibition of glucosidase in intestinal tract
Decrease in HbA1C (%)	1-2	1-2	1-2	1-2	0.5-1
Decrease in FBS (mg/dl)	50-60	60	40-80	60-70	30
Intake time	Before meal	Before meal	By meal or after meal	Before meal	Before meal
Daily recommended dosage (mg) *	5-20	1-16	500-3000	15-45	25-300
Side effects	Hypoglycemia	Hypoglycemia	GI [‡] effects, Lactic acid	Hepatotoxicity, anemia, Fluid retention	GI [‡] effects (Diarrhea, Bloating)
Contraindications	Cr [†] >2 mg/dl, Liver disease	Renal Failure	Renal dysfunction, Liver disease, CHF [¶] (Class 4)	Liver failure, CHF [¶]	Severe renal disease, Chronic intestinal disease, Liver disease

* The given dose is just for the sample of each group, † Creatinine, ‡ Gastrointestinal, ¶ Congestive heart failure

Table 2. The most available products of insulin in Iran

	Onset time[†]	Peak time[†]	Duration time[†]
Short acting (Regular)	0.5	2-3	4-6
Intermediate acting (NPH*)	1.5	4-8	10-18

* NPH: Neutral Protamine Hagedorn, † All times recorded based on hours

References

- 1- Harris MI, Hadden WC, Knowler WC, et al. Prevalence of diabetes and impaired glucose tolerance and plasma glucose levels in US population aged 20-74. *Diabetes* 1987; 36: 523-34.
- 2- King H, Rewers M. Global estimates for prevalence of diabetes mellitus and impaired glucose tolerance in adults. *Diabetes Care* 1993; 16:157-77.
- 3- World Health Organization. The world health report 1997: Conquering Suffering, enriching humanity. Geneva: World Health Organization: 1997.
- 4- Khatib OMN. Diabetes in the Eastern Mediterranean Region. In: Guidelines for the prevention, management and care of diabetes mellitus. WHO regional office for the Eastern Mediterranean; 2006. p. 9-12.
- 5- Larijani B, Abolhasani F, Mohajeri Tehrani MR, et al. Prevalence of diabetes mellitus in Iran in 2000. *Iranian Journal of Diabetes and Lipid Disorders* 2005; 4(3): 75-83.
- 6- Canadian Diabetes Association. Clinical Practice Guidelines Expert Committee. Diabetes in the Elderly 2003:S106-S109.
- 7- Khoshbin S, Eshrati B, Azizabadi Farahani A, et al. An investigation into the status of the country's aged population in 2002. Iran's Ministry of Health, Treatment, and Medical Education, Deputy Health Office, Family and Population Health Bureau. 2007: 5-10
- 8- Abolhasani F, Mohagerie Tehrani MR, Tabatabaei O, et al. Burden of diabetes mellitus and its complications in Iran in year 2000. *Iranian Journal of Diabetes and Lipid Disorders* 2005; 5(1): 35-48.
- 9- American Diabetes Association. Standards of medical care in diabetes. *Diabetes Care* 2007; 30(Suppl 1): S4-S41.
- 10- Harris MI, Flegal KM, Cowie CC. Prevalence of diabetes, impaired fasting glucose, and impaired glucose tolerance in U.S. adults. The Third National Health and Nutrition Examination Survey, 1988-1994. *Diabetes Care* 1998; 21: 518-24.
- 11- Tessier D, Meneilly G. Diabetes management in the elderly. In: Gerstein HC, Haynes RB. Evidence-based diabetes care. 1st edition, Canada. BC Decker Inc; 2001. p. 370-9.
- 12- Choi KM, Lee J, Kim DR, et al. Comparison of ADA and WHO criteria for the diagnosis of diabetes in elderly Koreans. *Diabet Med* 2002; 19: 853-857.
- 13- McBean AM, Gilbertson DT, Li Sh, et al. Differences in diabetes prevalence, incidence and mortality among the elderly of four racial/ethnic groups: whites, blacks, Hispanics, and Asians. *Diabetes care* 2004; 27: 2317-2324.
- 14- Nakano T, Ito H. Epidemiology of diabetes mellitus in old age in Japan. *Diabetes Research and Clinical Practice* 2007; 77: S76-S81.
- 15- Varas-Lorenzo C, Rueda de Castro AM, Maguire A, et al. Prevalence of glucose metabolism abnormalities and cardiovascular co-morbidity in the US elderly adult population. *Pharmacoepidemiology and drug safety* 2006; 15: 317-326.
- 16- Sloan FA, Bethel A, Jr DR, et al. The growing burden of diabetes mellitus in the US elderly population. *Arch Intern Med* 2008; 168(2): 192-199.
- 17- Azizi F, Emami H, Salehi P, et al. Cardiovascular risk factors in the elderly: Tehran Lipid and Glucose Study (TLGS). *Iranian Journal of Endocrinology and Metabolism* 2003; 17(5): 3-14.
- 18- Dorosty AR, Alavi Naeini AM. Correlation of elderly nutritional status with cardio-vascular disease and diabetes. *Tehran University Medical Journal* 2007; 65(3): 68-71.
- 19- Naghavi M, Jamshidi HR. Receiving Health Services in 2002 in the Islamic

- Republic of Iran, Deputy Health Office, Center for Health Promotion. 2005: 5-58.
- 20- The Iranian Statistics Center, Publication List, National Statistics Portal. Available at: <http://www.sci.org.ir/portal/faces/public/sci/sci.mahsulatvakhadamat/sci.paygah/sci.fehrest> (access: 6 October 2008).
 - 21- Schieber G, Klingen N. Health Financing Reform in Iran: Principles and Possible Next Steps. Social Security Research Institute Health Economic Congress Tehran, Islamic Republic of Iran October 30-November 1, 1999.
 - 22- Motta M, Bennati E, Capri M, et al. Diabetes mellitus in the extreme longevity. *Experimental Gerontology* 2008; 43: 102-105.
 - 23- IDF Asian-Pacific type 2 Diabetes Policy Group, type 2 diabetes. Practical Targets and treatments. Sydney: Health Communications Australia 1999.
 - 24- Scheen AJ. Diabetes mellitus in the elderly: insulin resistance and/or impaired insulin secretion? *Diabetes Metabol* 2005; 31: 5S27-5S34.
 - 25- Chau D, Edelman SV. Clinical Management of Diabetes in the Elderly. *Clinical Diabetes* 2001; 19(4): 172-175.
 - 26- The Italian Longitudinal Study on Aging (ILSA) Working Group. Prevalence of chronic diseases in older Italians: Comparing self-reported and clinical diagnoses. *Int J Epidemiol* 1997; 26: 995-1002.
 - 27- Motta M, Bennati E, Ferlito L, et al. A review on the actual trends of insulin treatment in elderly with diabetes. *Arch Gerontol Geriatr* 2008; 47: 151-61.
 - 28- California Healthcare Foundation/American Geriatrics Society Panel on improving care for elders with diabetes. Guidelines for Improving the Care of the Older Person with Diabetes Mellitus. *JAGS* 2003; 51: S265-S280.
 - 29- Bourdel-Marchasson I, Helmer C, Fagot-Campagna A, et al. Disability and quality of life in elderly people with diabetes. *Diabetes & Metabolism* 2007; 33: S66-S74.
 - 30- Gavard JA, Lustman PJ, Clouse RE. Prevalence of depression in adults with diabetes: an epidemiological evaluation. *Diabetes Care* 1993; 16: 1167-78.
 - 31- Larijani B, Khoramshahi Bayan M, Khalili Gorgani m, et al. Association of depression and diabetes in the doctor Shariati diabetes clinic and Iranian Diabetes Association. *Iranian Journal of Diabetes and Lipid Disorders* 2004; 3(1): 77-82.
 - 32- Maraldi C, Volpato S, Penninx BW, et al. Diabetes mellitus, glycemic control, and incident depressive symptoms among 70- to 79-year-old persons. *Arch Intern Med* 2007; 167: 1137-1144.
 - 33- Egede LE. Diabetes, major depression, and functional disability among U.S. adults. *Diabetes Care* 2004; 27: 421-8.
 - 34- Gregg EW, Engelgau MM, Narayan V. Complications of diabetes in elderly people. *BMJ* 2002; 325: 916-7.
 - 35- Mohsen Khoshniat Niko ‘Farah Madarshahian ‘Mohsen Hassanabadi et al. Comparative study of cognitive status in elderly with and without type 2 diabetes. *Iranian Journal of Diabetes and Lipid Disorders* 2007; 7(1): 71-6.
 - 36- Naslafkih A, Se stier F. Diabetes mellitus related morbidity, risk of hospitalization and disability. *J Insur Med* 2003; 35: 102-13.
 - 37- Chanudet X, Bonnevie L, Bauduceau B. Coronary heart disease and cardiovascular autonomic neuropathy in the elderly diabetic. *Diabetes & Metabolism* 2007; 33: S19-S31.
 - 38- Muggeo M, Verlato G, Bonora E, et al. The Verona Diabetes Study: a population-based survey on known diabetes mellitus prevalence and 5-year all-cause mortality. *Diabetologia* 1995; 38: 318-25.

- 39- Cacciatori V, Bonora E, Travia D, et al. Comparison of elderly diabetes and mature age diabetes: similarities and differences. *Archives of Gerontology and Geriatrics* 1991; 2(Suppl): S223-S226.
- 40- Bethel MA, Sloan FA, Belsky D, et al. Longitudinal incidence and prevalence of adverse outcomes of diabetes mellitus in elderly patients. *Arch Intern Med* 2007; 167: 921-927.
- 41- Gharipour M, Baghei A, Boshtam M, et al. Prevalence of metabolic syndrome among the adults of central of areas of Iran (as part of "Isfahan Healthy Heart Study"). *J of Birjand University of medical Sciences* 2006; 3(13): 56-62.
- 42- Nichols GA, Hillier TA, Erbey JR, et al. Congestive heart failure in type 2 diabetes. *Diabetes Care* 2001; 24:1614-9.
- 43- Solang L, Malmberg K, Ryden L. Diabetes mellitus and congestive heart failure. *Eur Heart J* 1999; 20: 789-95.
- 44- Action to Control Cardiovascular Risk in Diabetes (ACCORD) Trail. Available at: <http://www.nhlbi.nih.gov/health/prof/heart/other/accord>. Updated June 6, 2008.
- 45- Barker WH, Mullooly JP, Getchell W. Changing incidence and survival for heart failure in a well-defined older population, 1970-1974 and 1990-1994. *Circulation* 2006; 113: 799- 805.
- 46- Canadian Diabetes Association Clinical Practice Guidelines Expert Committee. Canadian Diabetes Association 2003 Clinical Practice Guidelines for the Prevention and Management of Diabetes in Canada. *Canadian J Diabetes* 2003; 27(Suppl 2): S1-S152.
- 47- American Diabetes Association. Standards of medical care in diabetes-2009. *Diabetes Care* 2009; 32(Suppl 1): S13-S61
- 48- Chelliah A, Burge MR. Hypoglycaemia in Elderly Patients with Diabetes Mellitus. Causes and Strategies for Prevention. *Drugs Aging* 2004; 21(8): 511-530.
- 49- McAulay V, Frier BM. Hypoglycemia. In: Sinclair AJ, Finucane P.(Editors). *Diabetes in old age*. 2nd Edition. UK. John Wiley & Sons Ltd; 2001. p. 133-152.
- 50- Larijani B, Afshari M, Darvishzadeh F, et al. Lower limb amputation in patients with diabetic foot ulcer: a 22 year review. *MJIRC* 2006; 8(3): 21-4.
- 51- Endocrine and Metabolism Research Center, Tehran University of Medical Sciences; Regional_ National Diabetes Guideline/Diabetic foot. Editors: Seyed Masoud Arzaghi, Ozra Tabatabaei Malazy, Tehran, in press by publisher: Charity Foundation for Special Diseases, 2009. Available in Persian language at <http://emrc.tums.ac.ir/df> lasted update 5/9/2009.
- 52- Lawrence SM, Wraight PR, Campbell DA, et al. Assessment and management of inpatients with acute diabetes-related foot complications: room for improvement. *Internal Medicine Journal* 2004; 34: 229-233.
- 53- Thomson FJ, Masson EA. Can elderly diabetic patients co-operate with routine foot care? *Age and aging* 1992; 21: 333-7.
- 54- Pataky Z, Vischer U. Diabetic foot disease in the elderly. *Diabetes & Metabolism* 2007; 33: S56-S65.
- 55- Forouzandeh F, Aziz Ahari A, Abolhasani f, et al. E valuation of foot neurovascular status in diabetic patients referred to diabetes clinic of Dr. Shariati Hospital, 2003-2004. *Iranian Journal of Diabetes and Lipid Disorders* 2005; 4(4): 43-52.
- 56- Shojaiefard A, Khorgami Z, Larijani B. Independent risk factors for amputation in diabetic foot. *Int J Diab Dev Ctries* 2008; 28: 32-7.
- 57- Shojaiefard A, Khorgami Z, Larijani B. Septic diabetic foot is not necessarily

- an indication for amputation. *J Foot Ankle Surg* 2008; 47: 419-23.
- 58-Booya F, Bandarian F, Larijani B, Pajouhi M, Nooraei M, Lotfi J. Potential risk factors for diabetic neuropathy: a case control study. *BMC Neurol* (Online access) 2005; 5:24.
 - 59-Forouzandeh F, Aziz Ahari A, Abolhasani F, Larijani B. Comparison of different screening tests for detecting diabetic foot neuropathy. *Acta Neurol Scand* 2005; 112:409-13.
 - 60-Butcher A, Dodson P. Visual loss. In: Sinclair AJ, Finucane P. *Diabetes in old age*. 2nd edition, Chichester UK, John Wiley & Sons; 2001. p.119-131
 - 61-.Whiteside MM, Wallhagen M I, Pettengill E. Sensory impairment in older adults: part 2: vision loss. *AJN* 2006; 106(11): 52-61.
 - 62-Massin P, Kaloustian E. The elderly diabetic's eyes. *Diabetes & Metabolism* 2007; 33: S4-S9.
 - 63-Endocrine and Metabolism Research Center, Tehran University of Medical Sciences; *Diabetes Guideline*. Editors: Seyed Masoud Arzaghi, Sasan Sharghi, Tehran, publisher: Charity Foundation for Special Diseases, 2008. Available in <http://vcemrc.tums.ac.ir/> lasted updated 2009, available in Persian language.
 - 64-Wong T Y, Loon S-C, Saw S-M. The epidemiology of age related eye diseases in Asia. *Br J Ophthalmol* 2006; 90:506-11.
 - 65-Manaviat MR, Afkhami Ardekani M. A study of the rate of presence of retinopathy in diabetic type II patients referring to the Yazd Diabetic Center. *J of Shahid Sadoughi University of Medical Sciences and Health Services* 2003; 4(10): 41-6.
 - 66-Cahill M, Halley A, Codd M, et al. Prevalence of diabetic retinopathy in patients with diabetes mellitus diagnosed after the age of 70 years. *Br J Ophthalmol* 1997; 81: 218-222.
 - 67-Kato S, Takemori M, Kitano S, et al. Retinopathy in older patients with diabetes mellitus. *Diabetes Research and Clinical Practice* 2002; 58: 187-192.
 - 68-De Voogd S, Ikarm MK, Wolfs RCW, et al. Is Diabetes Mellitus a Risk Factor for Open-Angle Glaucoma? *Ophthalmology* 2006; 113: 1827-1831.
 - 69-American Diabetes Association. Nephropathy in Diabetes. *Diabetes Care* 2004; 27 (Suppl 1): S79-S83.
 - 70-Sorensen L, Yue DK. Screening and treatment strategies for diabetes complications in the elderly. In: Sinclair AJ, Finucane P. *Diabetes in old age*. 2nd edition, Chichester UK, John Wiley & Sons; 2001. p.41-9.
 - 71-Mohammadzadeh F, Omidfar K, Heshmat R, et al. Comparison of three methods of urine sampling for screening of albuminuria in diabetic patients using immunoturbidometric assay. *Iranian Journal of Diabetes and Lipid Disorders* 2007; 6(3): 219-24.
 - 72-Valizadeh m, Nasser Farahmand A, Mousavi Nasab N, et al. Determining the value of albumin to creatinine ratio in a single morning sample, compared to the 24-Hour urinary albumin excretion test, for determining microalbuminuria in diabetic patients. *Iranian Journal of Diabetes and Lipid Disorders* 2009; 8: 27-36.
 - 73-Nakhjavani M, Azmoudeh Ardalan F, Mazouji A. Relative frequency of microalbuminuria in patients with type 2 diabetes attending the diabetes clinic at Imam Khomeini University Hospital (2001-2). *Iranian Journal of Diabetes and Lipid Disorders* 2002; 2(1): 153-8.
 - 74-Soell M, Hassan M, Miliauskaitė A, et al. The oral cavity of elderly patients in diabetes. *Diabetes & Metabolism* 2007; 33: S10-S18.
 - 75-Rad M, Hashemipour MS, Karimi MR. The relationship between oral manifestations in diabetes and duration of disease and glycemic control. *Iranian*

- Journal of Diabetes and Lipid Disorders 2006; 6(2): 159-67.
- 76- Diabetes and sexual function. Available in <http://diabetes.ygoy.com> lasted update October 2007.
- 77- Tabatabaei Malazy O, Shariat M, Heshmat R, et al. Vulvovaginal Candidiasis and its influencing factors in diabetes clinic of Dr. Shariati Hospital 2002-2005. Iranian Journal of Diabetes and Lipid Disorders 2006; 5(4): 345-352.
- 78- Vinik A, Richardson D. Erectile dysfunction. In: Sinclair AJ, Finucane P. Diabetes in old age. 2nd edition, Chichester UK, John Wiley & Sons; 2001. p.89-102.
- 79- Mofid A, Zandieh S, seyed Alinaghi SA, et al. Prevalence of erectile dysfunction in Tehranian diabetic men. Iranian South Medical J 2007; 1(10): 46-53.
- 80- Cayea D, Boyd C, Durso SC. Individualizing therapy for older adults with diabetes mellitus. Drugs Aging 2007; 24(10): 851-863.
- 81- Niefeld MR, Braunstein JB, Wu AW, et al. Preventable hospitalization among elderly medicare beneficiaries with type 2 diabetes. Diabetes Care 2003; 26: 1344-1349.
- 82- Wolff JL, Starfield B, Anderson G. Prevalence, expenditures, and complications of multiple chronic conditions in the elderly. Arch Intern Med 2002; 162 (20): 2269-76.
- 83- Piette JD, Kerr EA. The impact of comorbid chronic conditions on diabetes care. Diabetes Care 2006; 29 (3): 725-31.
- 84- Tessier DM, Lassmann-Vague VJR. Diabetes and education in the elderly. Diabetes & Metabolism 2007; 33: S75-S78.
- 85- Tabatabaei Malazy O, Heshmat R, Taheri E, et al. Comparison of a generic and a brand metformin product in type II diabetes: A double blind randomized clinical trial study. DARU 2007; 15(2): 113-7.
- 86- Song MK, Rosenthal MJ, Naliboff BD. Effects of bovine prostate powder on zinc, glucose and insulin metabolism in old patients with non-insulin-dependent diabetes mellitus. Metabolism 1998; 47: 39-43.
- 87- Paolisso G, D'Amore A, Galzerano D. Daily vitamin E supplements improve metabolic control but not insulin secretion in elderly type 2 diabetic patients. J Clin Endocrinol Metab 1994; 78: 1510-4.
- 88- Paolisso G, Scheen A, Cozzolino D. Changes in glucose turnover parameters and improvement of glucose oxidation after 4 week magnesium administration in elderly noninsulin-dependent (type 2) diabetic patients. J Clin Endocrinol Metab 1994; 78: 1510-4.
- 89- Paolisso G, D'Amore A, Balbi V. Plasma vitamin C affects glucose homeostasis in healthy subjects and in non-insulin-dependent diabetics. Am J Physiol 1994; 266: E261-8.
- 90- Ligtenberg PC, Hoekstra JBL, Bol E. Effects of physical training on metabolic control in elderly type 2 diabetes mellitus patients. Clin Sci 1997; 93: 127-35.
- 91- Agurs-Collins TD, Kumanyika SK, Ten Have TR. A randomized controlled trial of weight reduction and exercise for diabetes management in older African-American subjects. Diabetes Care 1997; 20: 1503-11.
- 92- Petterson T, Lee P, Hollis S, et al. Wellbeing and treatment satisfaction in older people with diabetes. Diabetes Care 1998; 21: 930-5.
- 93- Esteghamati A, Hassabi M, Halabchi F, et al. Exercise prescription in patients with diabetes type 2. Iranian Journal of Diabetes and Lipid Disorders 2008; 7 : 1-15.
- 94- American Diabetes Association. Diabetes mellitus and exercise. Diabetes Care 2002; 25(Suppl 1): S64-68.

- 95- Wilson M, Moore M P, Lunt H. Treatment satisfaction after commencement of insulin in Type 2 diabetes. *Diabetes Res Clin Pract* 2004; 66: 263-7
- 96- Gilden JL, Hendryx M, Casia C. The effectiveness of diabetes education programs for older patients and their spouses. *J Am Geriatr Soc* 1989; 37: 1023-30.
- 97- Kronsbein P, Jorgens V, Venhaus A. Evaluation of a structured treatment and teaching program on non-insulin-dependent diabetes. *Lancet* 1988; 2: 1407-11.
- 98- Meltzer S, Leiter L, Daneman D, et al. 1998 clinical practice guidelines for the management of diabetes in Canada. *Can Med Assoc J* 1998; 159(Suppl8): S1-29.
- 99- Salas M, Bubolz T, Caro JJ. Impact of physical functioning of health status on hospitalizations, physician visits, and costs in diabetic patients. *Arch Med Res* 2000; 31: 223-7.
- 100- Chehade JM, Mooradian AD. Drug therapy: Current and emerging agents. In: Sinclair AJ, Finucane P (Editors). *Diabetes in old age*, 2nd edition. UK. John Wiley & Sons, Ltd; 2001. p.199-214.