

## CENTILE CHARTS OF MOTOR DEVELOPMENT IN GIRLS AGED BETWEEN 7.5-19.5 FROM THE KUJAWSKO-POMORSKIE DISTRICT

**Marek Napierała**

*Faculty of Physical Education, Kazimierz Wielki University in Bydgoszcz, Poland*

Submitted in May, 2006

---

The aim of the research was to present motor development with the help of centile charts, where the centile curves determine particular parts of the population: 3, 10, 25, 50, 75, 90 and 97 percentile.

The research was conducted in the spring of 2001. A survey of 11 803 girls aged between 7.5 and 19.5 years old, from various types of rural and urban schools of the Kujawsko-Pomorskie district, was conducted. Centile charts make it possible to observe the process of the development of the examined characteristic, to answer the question as to whether an individual is at the level which is characteristic of a certain age, and, in case of aberrations, to come to a conclusion that the process of growth is disturbed. While determining the centile position one can become familiar with the level of development of an individual against calendar individuals of the same age. The value of C50 determines the calendar age of an individual. The results of the empiric research shows that the curve of the individual development of a child "wanders" within some channels, which is quite normal and does not mean a distortion of development.

---

*Keywords: Motor fitness of girls, centile charts.*

### INTRODUCTION

Political changes have brought about social and economical changes in Poland. This situation can influence the biological development of children and teenagers. In order to present individual development one can use centile charts, which are widely used by doctors and teachers. They are also being developed by the employees of the International Institute of Mothers and Children in Warsaw (Bożilow et al., 2004).

The development of an individual at every stage is a function of genetic factors, environment and time. As a result of interaction between genotype and environment, development progresses in accordance with its specific developmental line (Stein & Julian, 2003).

### METHOD

Motor development was presented with the help of centile charts, where the centile curves determine particular parts of the population: 3, 10, 25, 50, 75, 90 and 97 percentile (Stupnicki et al., 2003).

The research was conducted in the spring (at the turn of April and May) of 2001 and covered 11 803 girls aged between 7.5 and 19.5 from various types of rural and urban schools of the Kujawsko-Pomorskie district. The number of girls in particular age groups as well as the environment they lived in were similar.

The calendar groups were created in accordance with the terms generally used in tests of the developmen-

tal type (Malinowski, 1978; Kemper & Van Mechelen, 1996). It was assumed that a 7.5 year old child was one who, on the day the research was done, was age between 7 years and eight years without one day. Considering the 24 hour long biological rhythm, the research on motor activity was conducted between 10 a. m. and 1 p. m.

Both daytime and extramural students of the Academy of Bydgoszcz (now Kazimierz Wielki University in Bydgoszcz), as well as the teachers of physical education and the medical personnel from the examined schools proved very helpful. All examiners were trained in accordance with the research instructions.

In the opinion of the author motor ability is mainly seen at the level of manifestation of abilities and motor skills of an individual during a particular act of movement, including fitness tests. Fitness as a characteristic was adopted as the ability to solve motor tasks and the level of the fitness of the population was expressed in average values obtained in motor tests.

In order to determine motor abilities, eight tests from the International Committee on the Standardisation of Physical Fitness were used. The test includes universal assessment of groups of muscles of the whole body. The technical elements are not aimed at any of the basic sports disciplines.

Before the tests, the examined parties did a warm-up typical for intensive physical training. The sports outfit used during research was to consist of a t-shirt and shorts (alternatively a light track-suit) and some sports shoes without spikes and pins, with a non-slippery sole. The tests of overhang and forward bend were to be done


without any footwear. Instructions, helpful while doing particular tests, from the International Committee on the Standardisation of Physical Fitness, as described by Larson (1974) were adopted.

The centile charts were developed by the author on the basis of the obtained results.


**RESULTS**

The results are presented in graphic form, on centile charts, Figure 1-8.


**Fig. 1**  
The centile chart of the girls' 50 m run




**Fig. 2**  
The centile chart of the girls' 600m for girls aged 7.5-11.5 and 800 m age 12.5-19.5 run




**Fig. 3**  
The centile chart of the girls' 4 × 10 m shuttle run



**Fig. 4**  
The centile chart of the girls' forward bend of the trunk from the standing position




**Fig. 5**  
The centile chart of the girls' - sit-ups within 30 sec.




**Fig. 6**


The centile chart of the girls' horizontal bar bent arm hang

**Fig. 7**

The centile chart of the girls' dynamometer hand grip squeeze

**Fig. 8**

The centile chart of the girls' standing long jump



## DISCUSSION

The centile charts make it possible to observe the process of the development of the examined characteristic, to answer the question as to whether an individual is at the level which is characteristic of a certain age, and, in case of aberrations, to come to a conclusion that the process of growth is disturbed (Stupnicki et al., 2003).

While determining the centile position one can become familiar with the level of development of an individual against calendar individuals of the same age. The value of C50 determines the calendar age of an individual.

## CONCLUSION

The results of the empiric research show that the curve of individual development of a child "wanders" within some channels, which is quite normal and does not mean a distortion of development (Cieřlik et al., 1994). The developmental standard is a biological point of a reference which is used to determine the physical development of the population at the developmental age. The present evidence is a part of broader research into somatic and motor development of children and teenagers from the Kujawsko-Pomorskie district.

## REFERENCES

- Bořilow, W. et al. (2004). Siatki centylowe dla dzieci i młodzieży z regionów podkarpackiego: Pomorza Środkowego i radomskiego. In J. Zagórski, H. Popławska, & M. Skład (Eds.), *Uwarunkowania rozwoju dzieci i młodzieży wiejskiej*. Lublin: Instytut Medycyny Wsi.
- Cieřlik, J. et al. (1994). *Dziecko Poznańskie '90: Wzrastanie, dojrzewanie, normy i metody oceny rozwoju*. Poznań: Wydawnictwo Naukowe Bogucki.
- Kemper, H., & Van Mechelen, W. (1996). *Physical fitness testing of children: A European perspective*. Champaign, IL: Human Kinetics.
- Larson, L. A. (Ed.). (1974). *Fitness, health and work capacity: International standards for assessment*. New York: Macmillan.
- Malinowski, A. (Ed.). (1978). *Dziecko Wielkopolskie: Normy rozwoju fizycznego dzieci i młodzieży z różnych środowisk Wielkopolski*. UAM Seria Antropologia: Poznań.
- Stein, D., & Julian, U. (2003). Should the main objective of adapted physical education be the development of motor skills or the development of self-esteem. *Journal of Physical Education, Recreation & Dance*, 74.
- Stupnicki, R. et al. (2003). *Centylowe siatki sprawności fizycznej polskiej młodzieży wg testów Eurofit*. Studia i Monografie. Warszawa: AWF.

**CENTILOVÉ STUPNICE  
MOTORICKÉHO VÝVOJE  
U DÍVEK VE VĚKU 7,5-19,5 ROKU  
Z KUJAVSKO-POMOŘSKÉHO KRAJE**  
(Souhrn anglického textu)

Cílem výzkumů bylo představit motorický vývoj pomocí centilových stupnic, kdy centilové křivky určují jisté části populace: 3, 10, 25, 50, 75, 90 a 97 percentil.

Výzkumy byly prováděny na jaře roku 2001. Týkaly se 11 803 dívek ve věku mezi 7,5 a 19,5 lety z různých typů venkovských a městských škol kujavsko-pomořského kraje. Centilové stupnice umožňují sledovat proces vývoje zkoumaných charakteristik, zodpovědět na otázku, zda se jednatlivec nachází na úrovni charakteristické pro určitý věk a – v případě odchylek – dospět k závěru, že proces růstu je narušený. Při stanovování centilové pozice je možno poznat úroveň vývoje jednatlivce ve srovnání s jednatlivci téhož kalendářního věku. Kalendářní věk jednatlivce určuje hodnota C50. Výsledky empirických výzkumů ukazují, že křivka individuálního vývoje dítěte se klikatí v mezích určitých kanálů; to je však normální a neznamena to deformaci vývoje.

*Klíčová slova: motorická zdatnost dívek, centilové stupnice.*

**Dr. Marek Napierała**



Kazimierz Wielki University  
in Bydgoszcz  
Institute of Theory  
and Methodology  
of Physical Education  
ul. Chodkiewicza 30  
Bydgoszcz 85-064  
Poland

***Education and previous work experience***

Head of Physical Culture Department

***Scientific orientation***

Dynamics of development of somatic and motor features of children and youth in Kujawy-Pomorze region and conditioning of the development; motor activity of children and youth and its influence on their health.

***First-line publication***

- Napierała, M. (2002). State of motor development of school children aged 7.5–19.5 from Kujawy-Pomorze region. *Fizicheskoje vospitanie studentov tvorcheskikh specialnostei*, 6, 82–91.
- Napierała, M. (2002). Activity of elder the people. *Research yearbook: Academy of Physical Education and Sport*, 8, 129–143.
- Napierała, M. (2003). Increase of physical development of boys aged 7.5–19.5 from Kujawy-Pomorze region. *Research yearbook: University School of Physical Education*, 9, 127–132.
- Napierała, M. (2004). Environmental conditioning of the motor efficiency of girls from the Kujawy region. In F. Vaverka (Ed.), *Movement and Health* (pp. 216–220). Olomouc: Univerzita Palackého.
- Napierała, M. (2005). Motor development of children and youth in the Kujawy-Pomorze region. *Research yearbook: Studies in Physical Education and Sport*, 11, 73–80.
- Napierała, M. (2006). Infect with recreation. In J. Kosiewicz (Ed.), *Movement recreation for all* (pp. 37–44). Warszawa: Wydawnictwo i Księgarnie.