

DIVING DOCTOR'S DIARY

PULMONARY OEDEMA, DYSPNOEA AND DIVING

Carl Edmonds

Key Words

Cardiovascular, case reports, drugs, equipment, immersion, medical conditions and problems.

Introduction

Over a decade ago I was contacted by the physician on St Thomas Island in the Caribbean and asked: Why was it that there were so many surface snorkellers developing pulmonary oedema, and who also had hypertension treated with beta-blockers?

I explained to him the significance of the beta-blockers, i.e. the production of asthma-like states with obstructive airways disease, the reduction in effort tolerance and the possible detrimental cardiac effects of bradycardia of immersion. I was not then very interested in snorkellers. We certainly had observed the cardiac death risk in divers who were also on these drugs.¹

I was reminded of this association by the most recent *Alert Diver*, which described two scuba divers with pulmonary oedema, in two separate articles.^{2,3} The first was in a hypertensive female tourist taking beta-blockers (with the implication that if she had been seen by an Australian or New Zealand diving doctor, this would have been avoided).² The second was in a middle aged male who went on to cardiac surgery once the cause was clarified.³

I was impressed with the coincidence of recently reviewing two similar cases of cardiac induced pulmonary oedema, but with some interesting variations. These were similarly dyspnoeic, but in swimmers, not divers, and both were doctors. Each was identified as pulmonary oedema by the treating institution.

These two physicians present a not uncommon problem, similar to those of St Thomas', but one not well recognised; the production of dyspnoea associated with immersion per se.

Case reports

CASE 1

Late last year I received an e-mail from a Swedish pulmonary physician. She wrote:

"I found your interesting article via the Internet, on snorkelling deaths. I am 61 years old and visited Australia in October for a conference. We went to Cairns and tried snorkelling on the reef. In the water I got pulmonary oedema and had to be transported to hospital by helicopter and stay there for 4 days. I have never had problems like that before although I have some hypertension. I did not aspirate seawater and was not diving. I am not a good swimmer and it has made me a little nervous although I have snorkelled before. That cannot explain what happened. My question: Could it have been caused by a tight wetsuit or the dead space in the snorkel? I do not have large lung volumes, but I am physically active for my age.

Well, I survived and am not going to snorkel any more, but it would be nice to hear your opinion".

I am not enthusiastic about answering questions over the Internet, especially when I know that I am not going to get the full history and am denied the luxury of a physical examination, let alone appropriate investigations. I replied as follows:

"Before I respond to your questions, I need to know:-

- 1 What medication were you on (especially beta-blockers)?
- 2 Have you had a stress ECG? If yes, give the full report.
- 3 Any history of heart or lung disease (e.g. asthma)?"

The reply came almost before I had finished sending my questions!

"Dear Colleague,

Thank you for reading my question (Internet is fantastic).

- 1 I am on a beta-blocker, metoprolol tartrate (Metoprolol) 100 mg, and an angiotensin 2 receptor antagonist, candesartan cilexetil (Atacand) 16 mg.
- 2 No stress test was done before the snorkelling but it was before I left the Hospital in Cairns and that was without any problem. It was made because of a raised troponin level.
- 3 No history of heart or lung disease before. I am physically rather active for my age and my ECG at rest has been normal."

A discharge summary from Cairns Hospital indicated that the diagnosis was a cardiac induced pulmonary oedema, with myocardial ischaemia as the cause. She was commenced on Simvastatin and aspirin. The exercise stress test, while on beta-blockers, produced a maximum blood pressure of 220/100 and a maximum workload of 10.4 mets. There were no ECG changes or chest pains during exercise, but a U-shaped ST depression infero-laterally became more

horizontal during the recovery phase and was considered to be probably significant.

As so often happens in these cases, the patient's description was not completely accurate.

I informed her that I considered further investigations should be carried out regarding her cardiac status and that full lung function tests be performed. The latter were undertaken and there was no sign of any bronchial obstruction, even when on the beta-blockers.

CASE 2

This was more informative, because of the subsequent developments.

A 55 year old male had mild hypertension, was taking a beta-blocker (metoprolol) and aspirin, had no problems associated with scuba diving. But with surfing he regularly developed mild dyspnoea, 5-10 minutes after entering the surf, irrespective of the sea state. It would be relieved as he continued his surfing activities. Over 5 years this observation was verified on hundreds of occasions, but the dyspnoea seemed to be increasing in intensity and was affecting his aquatic fitness.

By the time he had his angiogram, which demonstrated multiple coronary vessel obstruction, he was experiencing dyspnoea not only during surfing but also with other exertion. He did not suffer angina pectoris in the aquatic environment, but this was becoming pronounced with terrestrial activities, being provoked by cold temperature, mild to moderate exertion, emotion, excessive food intake, etc.

Following a successful coronary artery bypass graft, all clinical evidence of cardiac ischaemia disappeared and he had no evidence of ischaemia on either the stress ECG or thallium scan.

When he resumed surfing he experienced no further dyspnoea with immersion and no apparent impairment of physical fitness during this activity.

This case demonstrates the successful influence of coronary flow restoration in the prevention of episodes of dyspnoea and pulmonary oedema during immersion.

Discussion

The pathophysiology underlying these cases was relatively clear because of the absence of any diving activity. With scuba diving there is a greater complexity in the development of dyspnoea and pulmonary oedema, as described in a recent diving medical text.⁴

During the 19th century immersion in a bath was one treatment given for "dropsy", demonstrating the effects of immersion on the redistribution of peripheral fluid.

Linnarsson assessed subjects with cardiac disease during an increasing work load, both in the water and on land, until symptoms or ECG abnormalities developed. The aquatic environment consistently masked the anginal symptoms of myocardial ischaemia.⁵

Diving diseases

Pulmonary oedema has been described in a variety of diving diseases. In some of these, it is a consequence of other diving respiratory pathology, such as in;

- the drowning syndromes, including salt water aspiration,
- pulmonary barotrauma,
- decompression sickness,
- underwater blast,
- some gas contaminants,
- pulmonary oxygen toxicity,
- some marine animal envenomations (e.g. the Irukandji syndrome).

In other instances pulmonary oedema has been described as the primary manifestation from a diving exposure.⁶ Sometimes dyspnoea occurs in otherwise healthy individuals during scuba diving. The disorders occur while swimming or diving in shallow water, sometimes in the head-out position. The symptoms include dyspnoea, cough, haemoptysis and expectoration of frothy sputum.

The pathophysiological explanations advanced are often not completely convincing. Nevertheless there are sufficient such cases to warrant concern. The various presumed aetiological types are as follows.

Pulmonary oedema of immersion.

Immersion of the body, with the head above water, will have significant pulmonary effects.⁸ Because of the hydrostatic pressure exerted by the water there is a redistribution of the blood (over half a litre) into the large intrathoracic blood vessels and, to a lesser extent, into the pulmonary microcirculation. Immersion also greatly affects the structure and function of the thoracic cavity. The following changes occur;

- reduced vital capacity (approximately 5%),
- reduced static lung compliance,
- increasing closing volumes leading to,
- increase in functional air trapping,
- increase in diffusing capacity,
- increased flow resistance in airways,
- altered ventilation, increasing in the apical regions and reduction in functional residual capacity.

Pulmonary oedema and dyspnoea associated with cardiac disease.

This may be related to pulmonary oedema of immersion. Wilmshurst recorded the observation of pulmonary oedema, inducing dyspnoea in divers, associated with hypertension and postulated a number of aetiologies.⁹ The stress factors which combine to induce this disorder include; increased cardiac loading due to the effect of water immersion, the respiratory changes noted above and cold induced peripheral vasoconstriction increasing the workload on the heart.

The effect therefore included increased cardiac pre-load, increased breathing resistance and exaggerated cardiac after-load.¹⁰

Some older divers, especially males, will develop dyspnoea 5 to 10 minutes after immersion. In these cases there is probably also a precipitation of cardiac ischaemia or arrhythmias because of the many physiological effects of immersion.

The following cardiovascular effects of immersion have been observed;

- increased central blood volume, with blood passing from the periphery to the lungs due to the hydrostatic gradient,
- increased stroke volume with increased cardiac output,
- increased peripheral perfusion,
- extrasystoles during the early phase of immersion (due to cardiac distension),
- the diving reflex, when facial immersion and trigeminal stimulation induce bradycardia, shunting of the blood to the heart/brain axis from the periphery and viscera, and peripheral vasoconstriction,
- the cardiac effects of cold exposure
- and hyperbaric bradycardia, usually by 10 beats per minute, which is possibly related to increased oxygen partial pressure, increased gas density, increased hydrostatic pressure and a narcotic effect of inert gas.

Each or all of these may play a part in inducing mild pulmonary oedema and the respiratory symptom of dyspnoea. Subsequent investigation will demonstrate the presence of coronary artery insufficiency or arrhythmia and the problem will disappear when this is corrected.

Troponin estimations may supplement the traditional ECG and cardiac enzymes (CK and CK-MB) in determining cardiac involvement.¹⁰

Asthma provocation.

The scuba situation is likely to induce asthma in those so predisposed. There are multiple provoking factors, including;

- exercise, especially if swimming against a significant tidal current,
- breathing against a resistance (the demand valve),
- breathing cold dry air (decompressed air),
- breathing dense air (related to depth),
- salt water spray inhalation (leaky demand valve) and
- psychological stress and hyperventilation.

Cold urticaria.

Cold urticaria is a specific disease due to histamine release after exposure to cold. It may affect the skin, gastrointestinal tract and the lungs.

In scuba diving there is a reduction in breathing air temperature as the air pressure drops during its passage from the high pressure cylinder to the diver (Charles Law) and by-passes the nasal cavity with its warming and humidification effects.

Deep diving dyspnoea.

Cold gas inhalation at depth makes all divers susceptible to dyspnoea at great depths, due to convective heat loss in the airways and the local response to this.

Other non-diving disorders

Other non-diving disorders may present as dyspnoea while diving, because of the occasional exceptional physical demands of this activity. Examples are lung disease, drug effects from beta-blockers or irritants such as cannabis.

References

- 1 Edmonds C and Walker D. Scuba diving fatalities in Australia and New Zealand. The human factor. *SPUMS J* 1989; 19 (3): 94-104
- 2 Taylor L. Fitness to dive. *Alert Diver SEAP Edition*. 2000; Oct-Dec: 4-7
- 3 Allen J. A cardiac near-catastrophe. *Alert Diver SEAP Edition*. 2000; Oct-Dec: 17-18
- 4 Edmonds C, Lowry C, Pennefather J and Walker R. *Diving and Subaquatic Medicine. 4th Edition*. London: Arnold Publication, 2001 in press
- 5 Linnarsson D. Karolinska Institute. Reported by Lourey CJ in The cardiac reflexes revisited. *SPUMS J* 1981; 11 (Suppl): 11-16
- 6 Lundgren CEG and Miller JN. *The Lung at Depth*. New York: Marcel Dekker, 1999
- 7 Hong SK. Breath-hold diving. Chapter 6 in *Diving Medicine*. Bove AA and Davis J. Eds. Philadelphia: Saunders, 1990
- 8 Wilmshurst PD. Cold induced pulmonary oedema in

- scuba divers and swimmers and subsequent development of hypertension. *Lancet* 1989; 1: 62-65
- 9 Elliott D and Moon R. In chapter 21 of *The Physiology and Medicine of Diving. 4th Edition.* Bennett PB and Elliott DH. Eds. London: Saunders, 1993
- 10 Vasikaran SD, Hitchcock T, Burnett JR and Clugston RA. Measuring myocardial damage. *Med J Aust* 2001; 174: 163-164

Additional references

- Bonneau A, Friemel F and Lapierre D. Electrocardiographic aspects of skin diving. *European J Appl Physiol* 1989; 58 (5): 487-493
- Eckenhoff RG and Knight DR. Cardiac arrhythmias and heart rate changes in prolonged hyperbaric air exposures. *Undersea Biomed Res* 1984; 11 (4): 335-367
- Eldridge L. Sudden unexplained death syndrome in cold water scuba diving. *Undersea Biomed Res* 1979; 6 (Suppl); 41
- McDonough JR, Barutt BS and Saffron RN. Cardiac arrhythmias as a precursor to drowning accidents. In *The Physiology of Breathhold Diving.* Lundgren CEG and Ferrigno M. Eds. Undersea and Hyperbaric Medical Society, Washington. 1987
- Pons M. Pulmonary oedema in healthy persons during scuba-diving and swimming. *European Respiratory J* 1995; 8: 762-767
- Weiler-Ravell D and Shupac A. Pulmonary oedema and haemoptysis induced by strenuous swimming. *Brit Med J* 1995; 311: 361-362
- Hampson NB and Dunsford RG. Pulmonary edema of scuba divers. *Undersea Hyperbaric Med* 1997; 24 (1): 29-33.

Dr Carl Edmonds' address is Diving Medical Centre, 66 Pacific Highway, St Leonards, New South Wales 2065, Australia. E-mail <divmed@ozemail.com.au>.


EUBS 2001

27TH ANNUAL MEETING OF THE EUROPEAN UNDERWATER AND BAROMEDICAL SOCIETY ON DIVING AND HYPERBARIC MEDICINE

September 12th to 16th 2001

Hotel Inter-Continental, Hamburg

For further details contact the EUBS website
<http://www.eubs.org>



The Hyperbaric Technicians and Nurses Association

with
Australian & New Zealand Hyperbaric Medical Group

is proud to present

The 9th Annual Scientific Meeting on Diving and Hyperbaric Medicine

19th- 22nd SEPTEMBER 2001

Conference Venue

THE ESPLANADE HOTEL

Located right in the heart of Fremantle, Western Australia

Guest Speakers

Dr Neil Hampson

Medical Director, Hyperbaric Department. Chief, Division of Pulmonary and Critical Care Medicine, Virginia Mason Medical Centre, Seattle, USA. President-Elect, Undersea and Hyperbaric Medical Society.

Dr Mike Bennett

Medical Director Department of Diving and Hyperbaric Medicine, Prince of Wales Hospital, Sydney, Australia. Chairman, ANZHMG. Member of Hyperbaric Oxygen Therapy Committee, Undersea and Hyperbaric Medical Society.

Mr Ron Nishi

Senior Scientist, Experimental Diving Unit, DCIEM, Toronto, Canada. Developer of the DCIEM Diving Decompression Tables.

FOR MORE INFORMATION CONTACT

Web Site at <http://www.users.bigpond.com/htna>
or Fremantle 2001 Organising Committee
Dept of Diving and Hyperbaric Medicine
Feemantle Hospital and Health Service
PO Box 480, Feemantle WA 6959
Phone 61-(0)8-9431-2233
Fax 61-(0)8-9431 2235
E-mail <jeff.pinkham@health.wa.gov.au>

SPUMS NOTICES

SOUTH PACIFIC UNDERWATER MEDICINE SOCIETY DIPLOMA OF DIVING AND HYPERBARIC MEDICINE

Requirements for candidates

In order for the Diploma of Diving and Hyperbaric Medicine to be awarded by the society, the candidate must comply with the following conditions:

- 1 The candidate must be a financial member of the Society.
- 2 The candidate must supply evidence of satisfactory completion of examined courses in both Basic and Advanced Course in Diving and Hyperbaric Medicine at an approved institution.
- 3 The candidate must have completed the equivalent (as determined by the Education Officer) of at least six months full time training in an approved Hyperbaric Medicine Unit.
- 4 The candidate must submit a written research proposal in a standard format for approval by the Education Officer before commencing their research project.
- 5 The candidate must produce, to the satisfaction of the Education Officer, a written report on the approved research project, in the form of a scientific paper suitable for publication.

Additional information

The candidate must contact the Education Officer to advise of their intended candidacy, seek approval of their courses in Diving and Hyperbaric Medicine and training time in the intended Hyperbaric Medicine Unit, discuss the proposed subject matter of their research proposed, and obtain instructions before submitting any written material or commencing a research project.

All research reports must clearly test a hypothesis. Preference will be given to reports of original basic or clinical research. Case series reports may be acceptable if thoroughly documented, subject to quantitative analysis, and the subject is extensively researched and discussed in detail. Reports of a single case are insufficient. Review articles may be acceptable if the world literature is thoroughly analysed and discussed, and the subject has not recently been similarly reviewed. Previously published material will not be considered.

It is expected that all research will be conducted in accordance with the "Joint NH&MRC/AVCC statement and guidelines on research practice" (available at <http://www.health.gov.au/nhmrc/research/nhmrcavc.htm>). All research involving humans or animals must be accompanied

by documentary evidence of approval by an appropriate research ethics committee. It is expected that research project and the written report will be primarily the work of the candidate.

The Education Officer reserves the right to modify any of these requirements from time to time.

The Education Officer's address is Dr David Doolette, Department of Anaesthesia and Intensive Care, The University of Adelaide, Adelaide, South Australia 5005. Telephone(0)8-8303-6382. Fax (0)8-8303-3909. E-mail <David.Doolette@adelaide.edu.au>

Key Words

Qualification.

REGISTRAR PLACES FOR THE DIPLOMA OF DIVING AND HYPERBARIC MEDICINE

New South Wales

Department of Diving and Hyperbaric Medicine Prince of Wales Hospital

Dr Mike Bennett <M.Bennett@unsw.edu.au>
High Street, Randwick, New South Wales 2031
Phone +61-(0)2-9382-3881
Fax +61-(0)2-9382-3882
One 6 month full time (Anaesthetic Provisional Fellow).
One 3 month part (50%) time rotation of anaesthetic registrar.

Submarine and Underwater Medicine Unit

Royal Australian Navy

Captain Robert Green <Robert.Green2@defence.gov.au>
Officer in Charge SUMU, HMAS PENGUIN
Middle Head Road, Mosman, New South Wales 2088
Phone +61-(0)2-9960-0333
Fax +61-(0)2-9960-4435
Paid position only available for RAN MOs.
No objection to someone getting self-funded work experience through this unit.

New Zealand

Slark Hyperbaric Unit

Royal New Zealand Navy

Dr Alison Drewry <navyhosp@ihug.co.nz>
Naval Base
Private Bag 32901, Auckland
Phone +64-(0)9-445-5972
Fax +64-(0)9-445-5973
One full-time Military Medical Officer.
Up to two self-funded positions for doctors (>3 years post graduation).

Queensland

Wesley Hospital Centre for Hyperbaric Medicine

Dr Simon Mitchell <smitchell@wesley.com.au>
 Suite 53, Sandford Jackson Building
 Auchenflower, Queensland 4066
 Phone + 61-(0)7-3371-6033
 Mobile + 61-0413-315-135
 Fax + 61-(0)7-3371-1566
 No funded training post.
 No objection to someone getting self funded work experience through this unit.

South Australia

Hyperbaric Medicine Unit

Royal Adelaide Hospital

Dr David Wilkinson <dwilkins@mail.rah.sa.gov.au>
 North Terrace, Adelaide, South Australia 5000
 Phone +61-(0)8-8222-5116
 Fax +61-(0)88232-4207
 One 6 month anaesthetic rotation as part of a Provisional Fellowship year.

Tasmania

Hyperbaric Medicine Unit

Royal Hobart Hospital

Dr Margaret Walker <margaret.walker@dchs.tas.gov.au>
 Hobart, TAS 7000
 Phone +61-(0)3-6222-8193
 Fax +61-(0)3-6222-8322
 One anaesthetic registrar post, half time anaesthesia, half-time hyperbaric medicine.

Victoria

The Alfred Hyperbaric Service

Dr Ian Millar <I.Millar@alfred.org.au>
 Alfred Hospital, Commercial Road, Prahran, Victoria 3181
 Phone +61-(0)3-9276-2269
 Fax +61-(0)3-9276 3052
 One 6 month full-time registrar position available to those in anaesthesia, intensive care physician or emergency medicine training programs.

Western Australia

Department of Diving and Hyperbaric Medicine

Fremantle Hospital

Dr Robert Wong <Robert.Wong@health.wa.gov.au>
 Fremantle Hospital, Fremantle Western Australia 6160
 Phone +61-(0)8-9431-2233
 Fax +61-(0)8-9431-2819
 One full-time registrar, one part-time anaesthetic registrar (6 months part-time rotation) and one emergency medicine registrar (12 months part time).

Key Words

Qualifications.

EMBASE

Readers will be aware that the South Pacific Underwater Medicine Society Journal is now indexed in the Elsevier Science medical data base, EMBASE. The information below has been gathered from pamphlets provided by Elsevier Science Secondary Publishing Division.

EMBASE is a current and comprehensive pharmacological and biomedical bibliographic database. It has extensive coverage of the drug-related literature, the database gathers information from more than 5,000 journals published in 70 countries. EMBASE contains more than 13 million bibliographic records from 1974 to the present, and over 555,000 records will be added annually. There are 5 million unique MEDLINE records included in EMBASE which creates an unparalleled single-file bibliographic resource. More than 80% of recent records contain full author abstracts. All bibliographic record source material has been subjected to a stringent peer-review process, while all other content on the site is continuously screened and reviewed. The result is high-quality, validated information. Fully indexed citations and complete author abstracts appear on average within 10 working days after receipt of the journal.

Each record contains the full bibliographic citation as well as indexing terms and codes. EMBASE records are indexed using EMTREE, a highly-sophisticated, hierarchically-ordered thesaurus with over 42,000 drug and medical terms (preferred terms), approximately 10,000 codes and close to 180,000 synonyms, including alternative drug names and MeSH subject heading used by the USA National Library of Medicine. Unique fields, such as drug trade and manufacturer names and device trade and manufacturer names, provide searchers with powerful record retrieval tools.

EMBASE.com is designed with a highly intuitive interface so there is little or no learning curve. Which allows a novice or experienced searcher to pinpoint the information needed quickly and easily. A system of powerful links allows navigation with ease through multiple layers of rich content so searches are both cost-effective and time-efficient. Records are displayed in an easy-to-read format, and special characters (such as Greek letters and superscripts) are displayed as in the original document, enhancing comprehension.

EMBASE can be accessed at <http://www.embase.com> which offers a free demonstration to experience the full functionality of EMBASE using a limited data set (approximately 30,000 records).

Key Words

Research