

L^1 –NORM OF INFINITELY DIVISIBLE RANDOM VECTORS AND CERTAIN STOCHASTIC INTEGRALS

MICHAEL B. MARCUS

Department of Mathematics

The City College of CUNY, New York, NY 10031

email: mbmarcus@earthlink.net

JAN ROSIŃSKI

Department of Mathematics

University of Tennessee, Knoxville, TN 39996

email: rosinski@math.utk.edu

submitted September 10, 2000 *Final version accepted* January 10, 2001

AMS 2000 Subject classification: 60E07, 60E15, 60H05

Infinitely divisible random variables, stochastic integrals

Abstract

Equivalent upper and lower bounds for the L^1 norm of Hilbert space valued infinitely divisible random variables are obtained and used to find bounds for different types of stochastic integrals.

1 L^1 –norm of infinitely divisible random vectors

Let X be an infinitely divisible random vector in a separable Hilbert space H . (See e.g. [3], [7].) Assume that $E\|X\| < \infty$, $EX = 0$ and that X does not have a Gaussian component. The characteristic function of X can be written in the form

$$E \exp i\langle y, X \rangle = \exp \left[\int_H (e^{i\langle y, x \rangle} - 1 - i\langle y, x \rangle) Q(dx) \right] \quad (1)$$

where Q is a unique σ –finite Borel measure on H with $Q(\{0\}) = 0$ and

$$\int_H \min\{\|x\|^2, \|x\|\} Q(dx) < \infty. \quad (2)$$

Q is called the Lévy measure of X . It is one of the principle entities used to describe the distribution of X .

We obtain bounds for $E\|X\|$ in terms of a functional of Q . Assume that $Q(H) > 0$ and let $l = l(Q) > 0$ be a solution of the equation

$$\xi(l) \stackrel{def}{=} \int_H \min\{l^{-2}\|x\|^2, l^{-1}\|x\|\} Q(dx) = 1. \quad (3)$$

It follows from (2) that $l(Q)$ is uniquely defined.

Note that (3) only depends on the projection of Q on \mathbf{R}^+ , which we denote by Q_r , i.e.

$$Q_r([t, \infty)) \stackrel{\text{def}}{=} Q(\{x \in H : \|x\| \geq t\}), \quad t > 0. \quad (4)$$

Thus we can also write (3) in the form

$$\xi(l) = \int_0^\infty \min\{l^{-2}x^2, l^{-1}x\} Q_r(dx) \quad (5)$$

and define $l(Q_r)$ to be the value of l for which the integral in (5) is equal to one. Clearly $l(Q) = l(Q_r)$.

The following theorem is the main result of this paper.

Theorem 1.1 *Under the above assumptions we have*

$$(0.25)l(Q_r) \leq E\|X\| \leq (2.125)l(Q_r). \quad (6)$$

If Q is symmetric, the constant in the upper bound of (6) can be decreased to 1.25.

In preparation for the proof of this theorem we consider a decomposition of X . We write $X = Y + Z$, where Y and Z are independent mean zero random vectors with characteristic functions given by

$$\begin{aligned} E \exp i\langle u, Y \rangle &= \exp \left(\int_{\|x\| < l} (e^{i\langle u, x \rangle} - 1 - i\langle u, x \rangle) Q(dx) \right) \\ E \exp i\langle u, Z \rangle &= \exp \left(\int_{\|x\| \geq l} (e^{i\langle u, x \rangle} - 1 - i\langle u, x \rangle) Q(dx) \right). \end{aligned} \quad (7)$$

where $l = l(Q)$.

The next lemma considers some moments of Y and Z . It is well known when $H = \mathbf{R}$.

Lemma 1.1 *Let $\lambda = Q(\{x : \|x\| \geq l\})$. Then*

$$E\|Y\|^2 = \int_{\|x\| < l} \|x\|^2 Q(dx) \quad (8)$$

$$\begin{aligned} E\|Y\|^4 &= \int_{\|x\| < l} \|x\|^4 Q(dx) + \left(\int_{\|x\| < l} \|x\|^2 Q(dx) \right)^2 \\ &+ 2 \int_{\|x\| < l} \int_{\|y\| < l} \langle x, y \rangle^2 Q(dx) Q(dy) \end{aligned} \quad (9)$$

and

$$\frac{1 - e^{-2\lambda}}{2\lambda} \int_{\|x\| \geq l} \|x\| Q(dx) \leq E\|Z\| \leq 2 \int_{\|x\| \geq l} \|x\| Q(dx). \quad (10)$$

When Q is symmetric, each number 2 in (10) can be replaced by 1.

Proof Let $\{e_j\}$ be an orthonormal basis in H . Let \bar{Q}_l denote Q restricted to $\{\|x\| < l\}$, the Lévy measure of Y . Since $\langle e_j, Y \rangle$ is a mean-zero real infinitely divisible random variable

$$\begin{aligned} E\|Y\|^2 &= \sum_j E\langle e_j, Y \rangle^2 \\ &= \sum_j \int_H \langle e_j, x \rangle^2 \bar{Q}_l(dx) \\ &= \int_H \|x\|^2 \bar{Q}_l(dx) \end{aligned}$$

which proves (8).

To prove (9) we consider an infinitely divisible random vector in \mathbf{R}^2 given by $(\langle e_j, Y \rangle, \langle e_k, Y \rangle)$. This random vector has characteristic function

$$E e^{is\langle e_j, Y \rangle + it\langle e_k, Y \rangle} = \exp \left(\int_H [e^{is\langle e_j, x \rangle + it\langle e_k, x \rangle} - 1 - i(s\langle e_j, x \rangle + t\langle e_k, x \rangle)] \bar{Q}_l(dx) \right).$$

Differentiating it twice with respect to s and t and then setting s and t equal to zero gives

$$\begin{aligned} E\langle e_j, Y \rangle^2 \langle e_k, Y \rangle^2 &= \int_H \langle e_j, x \rangle^2 \langle e_k, x \rangle^2 \bar{Q}_l(dx) + \int_H \langle e_j, x \rangle^2 \bar{Q}_l(dx) \int_H \langle e_k, x \rangle^2 \bar{Q}_l(dx) \\ &\quad + 2 \left(\int_H \langle e_j, x \rangle \langle e_k, x \rangle \bar{Q}_l(dx) \right)^2. \end{aligned}$$

Therefore

$$\begin{aligned} E\|Y\|^4 &= \sum_{j,k} E\langle e_j, Y \rangle^2 \langle e_k, Y \rangle^2 \\ &= \int_H \|x\|^4 \bar{Q}_l(dx) + \left(\int_H \|x\|^2 \bar{Q}_l(dx) \right)^2 \\ &\quad + 2 \sum_{j,k} \left(\int_H \langle e_j, x \rangle \langle e_k, x \rangle \bar{Q}_l(dx) \right)^2. \end{aligned}$$

Since the last term of this equation can be written as

$$\begin{aligned} &\sum_{j,k} \left(\int_H \langle e_j, x \rangle \langle e_k, x \rangle \bar{Q}_l(dx) \right)^2 \\ &= \sum_{j,k} \int_H \langle e_j, x \rangle \langle e_k, x \rangle \bar{Q}_l(dx) \int_H \langle e_j, y \rangle \langle e_k, y \rangle \bar{Q}_l(dy) \\ &= \int_H \int_H \left(\sum_j \langle e_j, x \rangle \langle e_j, y \rangle \right)^2 \bar{Q}_l(dx) \bar{Q}_l(dy) \\ &= \int_H \int_H \langle x, y \rangle^2 \bar{Q}_l(dx) \bar{Q}_l(dy) \end{aligned}$$

the proof of (9) is complete.

Let $\{W_i\}$ be independent identically distributed random vectors in H such that $P(W_i \in A) = \lambda^{-1}Q(A \cap \{\|x\| \geq l\})$ for all $A \in \mathcal{B}(H)$. Let N be a Poisson random variable with mean λ and independent of $\{W_i\}$. Then

$$Z \stackrel{\mathcal{D}}{=} \sum_{i=1}^N W_i - \lambda E W_1. \quad (11)$$

Consequently

$$E\|Z\| \leq E \sum_{i=1}^N \|W_i\| + \lambda E\|W_1\| = 2 \int_{\|x\| \geq l} \|x\| Q(dx)$$

which proves the upper bound in (10). (When Q is symmetric, $E W_1 = 0$ which gives the factor 1 instead of 2 in (10).)

To obtain the lower bound in (10) we first assume that Q is symmetric. Consequently the W_i 's are symmetric and we get

$$\begin{aligned} E\|Z\| &= \sum_{k=1}^{\infty} \left(E \left\| \sum_{i=1}^k W_i \right\| \right) P(N = k) \\ &\geq \sum_{k=1}^{\infty} (E\|W_1\|) P(N = k) \\ &= \frac{1 - e^{-\lambda}}{\lambda} \int_{\|x\| \geq l} \|x\| Q(dx). \end{aligned} \quad (12)$$

If Q is not symmetric, then we consider $\tilde{Z} = Z - Z'$, where Z' is an independent copy of Z . \tilde{Z} has symmetric Lévy measure $\tilde{Q}(A) = Q(A) + Q(-A)$ for $A \in \mathcal{B}(H)$ and $\tilde{Q}(\{\|x\| \geq l\}) = 2\lambda$. Applying (12) we get

$$\begin{aligned} 2E\|Z\| \geq E\|\tilde{Z}\| &\geq \frac{1 - e^{-2\lambda}}{2\lambda} \int_{\|x\| \geq l} \|x\| \tilde{Q}(dx) \\ &= \frac{1 - e^{-2\lambda}}{\lambda} \int_{\|x\| \geq l} \|x\| Q(dx). \end{aligned}$$

This completes the proof of Lemma 1.1.

Proof of Theorem 1.1 We first obtain the upper bound in (6). Using (8) and (10) we see that

$$\begin{aligned} E\|X\| &\leq (E\|Y\|^2)^{1/2} + E\|Z\| \\ &\leq \left(\int_{\|x\| < l} \|x\|^2 Q(dx) \right)^{1/2} + 2 \int_{\|x\| \geq l} \|x\| Q(dx) \\ &\leq (2.125)l. \end{aligned} \quad (13)$$

The last bound follows from the definition of $l(Q)$ by elementary calculus. Since the factor 2 can be dropped in (10) when Q is symmetric, the number 2.125 in (13) can be replaced by 1.25 in this case.

We now obtain the lower bound. It follows from (3) that either

(a) $\int_{\|x\|<l} \|x\|^2 Q(dx) \geq (0.5)l^2$ or

(b) $\int_{\|x\|\geq l} \|x\| Q(dx) \geq (0.5)l$.

Suppose that (a) holds. By Hölder's inequality we have

$$\begin{aligned} E\|Y\|^2 &= E(\|Y\|^{2/3}\|Y\|^{4/3}) \\ &\leq (E\|Y\|)^{2/3} (E\|Y\|^4)^{1/3} \end{aligned} \tag{14}$$

and by (8) and (9)

$$\begin{aligned} E\|Y\|^4 &\leq \int_{\|x\|<l} \|x\|^4 Q(dx) + 3\left(\int_{\|x\|<l} \|x\|^2 Q(dx)\right)^2 \\ &\leq l^2 E\|Y\|^2 + 3(E\|Y\|^2)^2. \end{aligned} \tag{15}$$

Combining (14) and (15) we get

$$\begin{aligned} E\|Y\| &\geq \frac{(E\|Y\|^2)^{3/2}}{\left(l^2 E\|Y\|^2 + 3(E\|Y\|^2)^2\right)^{1/2}} \\ &= \frac{E\|Y\|^2}{\left(l^2 + 3E\|Y\|^2\right)^{1/2}}. \end{aligned}$$

Using assumption (a) and (8) we see that

$$E\|Y\| \geq \frac{(0.5)l^2}{\left(l^2 + 3(0.5)l^2\right)^{1/2}}$$

because the function $y = t/(l^2 + 3t)^{1/2}$ is increasing for $t > 0$. Therefore

$$E\|X\| \geq E\|Y\| > (0.25)l.$$

This gives the lower bound in (6) when (a) holds.

Suppose now that (b) holds. Then

$$\begin{aligned} \int_{\|x\|\geq l} \left(\|x\| + (\|x\| - l)\right) Q(dx) \\ \geq \int_{\|x\|\geq l} \|x\| Q(dx) \geq (0.5)l. \end{aligned}$$

Consequently

$$\int_{\|x\|\geq l} \|x\| Q(dx) \geq (0.25)(1 + 2\lambda)l.$$

Using this in (10) we get

$$\begin{aligned} E\|X\| &\geq E\|Z\| \geq (0.25)\frac{(1 + 2\lambda)(1 - e^{-2\lambda})}{2\lambda}l \\ &> (0.25)l. \end{aligned}$$

This completes the proof of Theorem 1.1.

Corollary 1.1 *Let $\xi(l)$ be as in (3). Then*

$$(0.25) \min\{\xi(1), \sqrt{\xi(1)}\} \leq E\|X\| \leq (2.125) \max\{\xi(1), \sqrt{\xi(1)}\}. \quad (16)$$

Proof Observe that if $l(Q_r) < 1$, then $\xi(1) \leq l(Q_r) \leq \sqrt{\xi(1)}$ and if $l(Q_r) \geq 1$, then $\sqrt{\xi(1)} \leq l(Q_r) \leq \xi(1)$. Combining this observation with Theorem 1.1 gives (16).

Remark 1.1 Note that (6) is homogeneous in X , i.e. if X is replaced by cX , for some constant $c > 0$ then the bounds also change by a factor of c . Corollary 1.1 is often useful but (16) is not homogeneous in X .

Remark 1.2 There is no reason to believe that the constants in (6) of Theorem 1.1 are best possible. However, they provide good estimates for the absolute moment of an infinitely divisible random variable. To illustrate this point, consider $X = N - \lambda$, where N is a Poisson random variable with parameter λ . Since $Q = \lambda\delta_1$, we obtain $l(Q) = \lambda$ when $\lambda \in (0, 1)$ and $l(Q) = \lambda^{1/2}$ when $\lambda > 1$. A direct computation for $\lambda \in (0, 1)$ gives $E|X| = 2\lambda e^{-\lambda} = 2e^{-\lambda}l(Q)$. Hence the smallest possible constant on the right hand side of (6) must be at least 2. (In Theorem 1.1 we get 2.125). If λ is a positive integer, then $E|X| = 2\lambda^{\lambda+1}e^{-\lambda}/\lambda! \sim (2/\pi)^{1/2}l(Q)$. Hence the largest possible constant on the left hand side of (6) must smaller than 0.8. One can lower this bound further by considering the first moment of the symmetrization of X .

Remark 1.3 Let $X = \sum_{i=1}^N Y_i - \lambda EY_1$ be a centered compound Poisson random vector; here $\{Y_i\}$ are independent identically distributed random vectors in H with $E\|Y_1\| < \infty$ and N is a Poisson random variable with parameter λ independent of the $\{Y_i\}$. In this case $Q = \lambda\mathcal{L}(Y_1)$ and $l = l(Q)$ is the unique solution of the equation

$$\lambda E \min\{\|Y_1\|^2, l\|Y_1\|\} = l^2. \quad (17)$$

One sees from this that $l(Q) = K_{Y_1}(\lambda)$, where K_{Y_1} is the K -function of M. Klass, [4]. The proof of Theorem 1.1 parallels, to a certain extent, the estimates given in Section 1.4.3, [1] for the absolute moment of sums of independent identically distributed random variables in terms of the K -function. However our results seem to be easier to use when dealing with infinitely divisible random vectors. For example let $EY_1 = 0$ and $H = \mathbf{R}$. A straightforward application of Proposition 1.4.10, [1] gives the bounds for $E|X|$ in terms of $EK_{Y_1}(N)$. But $l(Q) = K_{Y_1}(EN)$. Since K_{Y_1} is nonlinear, the passage from $EK_{Y_1}(N)$ to $K_{Y_1}(EN)$ is far from obvious. By working with infinitely divisible random vectors directly we avoid this difficulty.

We give another set of bounds for $E\|X\|$ which, in certain circumstances, may be easier to compute. Let

$$\psi_r(u) \stackrel{def}{=} \int_0^\infty (1 - \cos uv) Q_r(dv), \quad u \geq 0. \quad (18)$$

(ψ_r is the Lévy exponent of a real valued symmetric infinitely divisible random variable. Indeed, when X itself is real valued and symmetric, ψ_r is the Lévy exponent of X). Let

$$\zeta(t) \stackrel{def}{=} \frac{1}{t} \int_0^{1/t} \frac{\psi_r(u)}{u^2} du. \quad (19)$$

Theorem 1.2 *Let X , Q and Q_r be as in Theorem 1.1 and let ψ_r be as given in (18). Let $t^* = t^*(\psi_r)$ be such that $\zeta(t^*) \leq 1/4$ and $t_* = t_*(\psi_r)$ be such that $\zeta(t_*) \geq 5/2$. Then*

$$(0.25)t_*(\psi_r) \leq E\|X\| \leq (2.125)t^*(\psi_r). \quad (20)$$

If Q is symmetric the constant in upper bound in (20) can be replaced by 1.25.

The proof of this theorem follows easily from the following lemma.

Lemma 1.2

$$\frac{1}{4}\xi(l) \leq \zeta(l) \leq \frac{5}{2}\xi(l) \quad (21)$$

where $\xi(l)$ is given in (3).

Proof We have

$$\begin{aligned} \zeta(l) &= (1/l) \int_{x < l} \int_0^{1/l} \frac{1 - \cos ux}{u^2} du Q_r(dx) \\ &\quad + (1/l) \int_{x \geq l} \int_0^{1/x} \frac{1 - \cos ux}{u^2} du Q_r(dx) \\ &\quad + (1/l) \int_{x \geq l} \int_{1/x}^{1/l} \frac{1 - \cos ux}{u^2} du Q_r(dx) \\ &\stackrel{def}{=} I + II + III. \end{aligned} \quad (22)$$

Note that when $|\theta| \leq 1$, $\theta^2/4 \leq 1 - \cos \theta \leq \theta^2/2$. Consequently,

$$\frac{x}{4} \leq \int_0^{1/x} \frac{1 - \cos ux}{u^2} du \leq \frac{x}{2} \quad (23)$$

and when $x < l$

$$\frac{x^2}{4l} \leq \int_0^{1/l} \frac{1 - \cos ux}{u^2} du \leq \frac{x^2}{2l}. \quad (24)$$

Using (24) and (23) in (22) we see that

$$\frac{1}{4}\xi(l) \leq I + II \leq \frac{1}{2}\xi(l) \quad (25)$$

which gives the lower bound in (21). Also

$$\begin{aligned} III &\leq (2/l) \int_{x \geq l} \int_{1/x}^{1/l} \frac{du}{u^2} Q_r(dx) \\ &\leq (2/l) \int_{x \geq l} |x| Q_r(dx). \end{aligned} \quad (26)$$

Combining (25) and (26), gives the upper bound in (21).

Proof of Theorem 1.2 It follows from (21) that

$$\frac{2}{5}\zeta(l) \leq \xi(l) \leq 4\zeta(l). \quad (27)$$

This implies that if $\zeta(t_*) \geq 5/2$ then $\xi(t_*) \geq 1$ and consequently $t_* \leq l(Q_r)$. Thus the lower bound in (20) follows from the lower bound in (6). The proof of the upper bound in (20) follows similarly.

It is particularly useful to know that the first moment of an infinitely divisible random variable is finite. Then we can obtain bounds for random variables with mean zero by first obtaining bounds for their symmetrized version. We use this technique in [6] to study the continuity of a wide class of infinitely divisible processes. Of course it would also be interesting to obtain bounds for other moments. The proof we give here does not readily extend to higher moments.

In the next section we consider stochastic integrals. In Section 3 we give some simple examples of how to use Theorems 1.1 and 1.2 to estimate the expected value of the norm of Hilbert space valued infinitely divisible random variables and certain stochastic integrals.

2 Stochastic integrals with respect to infinitely divisible random measures

Consider the infinitely divisible H -valued random variable X defined in (1). Given X there are two natural objects to consider, $\langle f, X \rangle$ where $f \in H$ and FX where F is an operator from H to another Hilbert space K . In Theorem 1.1 we obtain bounds for $E\|X\|$. As an extension of this result and as a corollary of this theorem we can obtain bounds for $E|\langle f, X \rangle|$ and $E\|FX\|$. However, it is natural to consider this extension in greater generality.

The “products” $\langle f, X \rangle$ and FX constitute natural bilinear forms taking values in $L^1(\Omega; \mathbf{R})$ and $L^1(\Omega; K)$, respectively. Therefore, for any measure M , defined on a measurable space S , and taking values in $L^1(\Omega; H)$, integrals of the form $\int_S \langle f(s), M(ds) \rangle$ and $\int_S F(s)M(ds)$ are well defined as Bartle type integrals. It is known that bounded deterministic functions are integrable in this setting. (See e.g. [2]).

Suppose that M , in addition, is an independently scattered infinitely divisible random measure. In this section we use Theorem 1.1 to obtain nice two-sided estimates for $E|\int_S \langle f(s), M(ds) \rangle|$ and $E\|\int_S F(s)M(ds)\|$ in terms of an Orlicz norm of f and F . Such bounds are relevant to our previous work [6] and can also be useful in modeling based on time–space infinitely divisible random noise. Furthermore, Theorem 1.1 gives a complete characterization of the class of L^1 –integrable deterministic functions with respect to independently scattered infinitely divisible random measures.

The L^1 bounds obtained in Theorem 1.1 can be generalized further to the case of stochastic integrals of random predictable integrands. This step is possible by applying decoupling inequalities along the lines of [5]. Such a generalization is not immediate, especially if one requires specific constants in the bounds. We will not consider it here. However, the L^1 bounds obtained in this section constitute a base for such a generalization.

The infinitely divisible random measures, that we integrate with respect to, are defined as follows: Let S be a Borel space equipped with a σ –finite Borel measure m . Let $\{\theta(\cdot, s)\}_{s \in S}$ be a measurable family of Lévy measures on H with the property that

there exists a sequence $\{S_n\}$ of Borel sets increasing to S such that for every n , $m(S_n) < \infty$ and

$$\int_{S_n} \int_H \min\{\|x\|^2, \|x\|\} \theta(dx, s) m(ds) < \infty.$$

Let $\mathcal{S} = \{A \in \mathcal{B}(S) : A \subset S_n \text{ for some } n\}$. It follows from Kolmogorov's Consistency Theorem that there exists an H -valued independently scattered random measure $M = \{M(A) : A \in \mathcal{S}\}$ such that, for every $A \in \mathcal{S}$

$$E \exp(i\langle y, M(A) \rangle) = \exp\left(\int_A \int_H (e^{i\langle y, x \rangle} - 1 - i\langle y, x \rangle) \theta(dx, s) m(ds)\right) \quad (28)$$

with $E\|M(A)\| < \infty$ and $EM(A) = 0$.

Let $f : S \rightarrow H$ be a simple function with support in \mathcal{S} , i.e. $f = \sum_{j=1}^n a_j I_{A_j}$, $A_j \in \mathcal{S}$. Consider the stochastic integral

$$\int_S \langle f, dM \rangle \stackrel{\text{def}}{=} \sum_{j=1}^n \langle a_j, M(A_j) \rangle. \quad (29)$$

Clearly, $E \int_S \langle f, dM \rangle = 0$.

To define the Orlicz space pseudo-norm we consider the function

$$\phi(y, s) \stackrel{\text{def}}{=} \int_H \phi_0(|\langle y, x \rangle|) \theta(dx, s), \quad y \in H, \quad s \in S \quad (30)$$

where

$$\phi_0(u) = u^2 I(u < 1) + (2u - 1) I(u \geq 1), \quad u \geq 0. \quad (31)$$

Since $\phi_0(u) = 2 \int_0^u \min\{v, 1\} dv$, ϕ_0 is a convex increasing function starting from 0 such that $\phi_0(2u) \leq 4\phi_0(u)$. Thus $\phi(\cdot, s)$ is a convex symmetric function with $\phi(0, s) = 0$ satisfying the Δ_2 -condition, i.e.

$$\phi(2y, s) \leq 4\phi(y, s), \quad y \in H.$$

Let $g : S \rightarrow H$ be a measurable function. We define an Orlicz space pseudo-norm of g by

$$\|g\|_\phi \stackrel{\text{def}}{=} \inf\{c > 0 : \int_S \phi(c^{-1}g(s), s) m(ds) \leq 1\}. \quad (32)$$

Theorem 2.1 *Let $\tilde{f} : S \rightarrow H$ be a measurable function satisfying*

$$\int_S \int_H \min\{|\langle \tilde{f}(s), x \rangle|^2, |\langle \tilde{f}(s), x \rangle|\} \theta(dx, s) m(ds) < \infty. \quad (33)$$

The definition of the stochastic integral in (29) can be extended to

$$\int_S \langle \tilde{f}, dM \rangle \quad (34)$$

where $E \int_S \langle \tilde{f}, dM \rangle = 0$ and

$$(0.125)\|\tilde{f}\|_\phi \leq E \left| \int_S \langle \tilde{f}, dM \rangle \right| \leq (2.125)\|\tilde{f}\|_\phi. \quad (35)$$

Proof Consider (29) and note that

$$\begin{aligned} & E \exp \left(iu \int_S \langle f, dM \rangle \right) \\ &= \exp \left(\int_S \int_H [e^{iu \langle f(s), x \rangle} - 1 - iu \langle f(s), x \rangle] \theta(dx, s) m(ds) \right). \end{aligned}$$

Hence $X = \int_S \langle f, dM \rangle$ has the same form as (1) where Q is the image of the measure $\theta(dx, s)m(ds)$ under the map $(s, x) \rightarrow \langle f(s), x \rangle$. $\xi(l)$ computed for this Q (or Q_r) in (3) also can be written as

$$\xi(l) = \int_S \int_H \min\{\langle l^{-1}f(s), x \rangle^2, |\langle l^{-1}f(s), x \rangle|\} \theta(dx, s)m(ds). \quad (36)$$

Comparing (36) and (30) we see that

$$\xi(l) \leq \int_S \phi(l^{-1}f(s), s) m(ds) \leq \xi(l/2).$$

Consequently

$$l(Q_r) \leq \|f\|_\phi \leq 2l(Q_r).$$

This bound combined with Theorem 1.1 yields

$$(0.125)\|f\|_\phi \leq E \left| \int_S \langle f, dM \rangle \right| \leq (2.125)\|f\|_\phi. \quad (37)$$

Since ϕ satisfies the Δ_2 -condition the results for f can be extended to hold for \tilde{f} .

If $H = R$ the integral (34) is the usual stochastic integral and $\|\cdot\|_\phi$ is a norm.

The integral in (34) is a real valued random variable. We now consider a Hilbert space valued stochastic integral. Let $F : S \rightarrow L(H, K)$, where $L(H, K)$ is the space of bounded linear operators from H into another separable Hilbert space K . We assume that for each $x \in H$ and $y \in K$, the function $s \rightarrow \langle y, F(s)x \rangle$ is measurable and that

$$\int_S \int_H \min\{\|F(s)x\|^2, |\langle F(s)x, x \rangle|\} \theta(dx, s)m(ds) < \infty. \quad (38)$$

We consider the stochastic integral

$$\int_S F dM \stackrel{def}{=} \sum_j \int_S \langle F^* e_j, dM \rangle e_j \quad (39)$$

where $F^* : S \rightarrow L(K, H)$, is the transpose of the operator $F(s)$, and $\{e_j\}$ is an orthonormal basis for K . (This integral could be defined via simple functions as outlined in the introduction to this section, however, since we have already established the existence of the inner product integrals, it is easier to use (39)).

The integral in (39) is well defined. To see this we show that the series in (39) converges and the limit does not depend on the choice of $\{e_j\}$. To begin note that

$$|\langle F^*(s)e_j, x \rangle| = |\langle e_j, F(s)x \rangle| \leq \|F(s)x\| \quad (40)$$

implying that (33) holds and $\int_S \langle F^* e_j, dM \rangle$ is well-defined. Next consider the random vector

$$S_{k,n} = \sum_{j=k}^n \int_S \langle F^* e_j, dM \rangle e_j.$$

Let $P_{k,n}y = \sum_{j=k}^n \langle y, e_j \rangle e_j$ and

$$\xi_{k,n}(1) \stackrel{def}{=} \int_S \int_H \min\{\|P_{k,n}F(s)x\|^2, \|P_{k,n}F(s)x\|\} \theta(dx, s) m(ds).$$

It follows from Corollary 1.1 that

$$E\|S_{k,n}\| \leq (2.125) \max\{\xi_{k,n}(1), \sqrt{\xi_{k,n}(1)}\}.$$

By (38) and the Dominated Convergence Theorem we see that the series in (39) converges in L^1_H .

To show that the right hand side of (39) does not depend on the choice of basis we perform the following formal computation which can be justified in the same way as the above proof of convergence. Let $\{g_j\}$ be another orthonormal basis in K . Then

$$\begin{aligned} & \sum_k \int_S \langle F^* g_k, dM \rangle g_k \\ &= \sum_{i,j} \sum_k \langle g_k, e_i \rangle \langle g_k, e_j \rangle \int_S \langle F^* e_i, dM \rangle e_j \\ &= \sum_{i,j} \langle e_i, e_j \rangle \int_S \langle F^* e_i, dM \rangle e_j \\ &= \sum_j \int_S \langle F^* e_j, dM \rangle e_j. \end{aligned}$$

We proceed to develop bounds for the L^1 norm of the stochastic integral. Let

$$\Phi(U, s) \stackrel{def}{=} \int_H \phi_0(\|Ux\|) \theta(dx, s), \quad U \in L(H, K), \quad s \in S \quad (41)$$

where ϕ_0 is given by (31), and define a pseudo-norm

$$\|F\|_\Phi \stackrel{def}{=} \inf\{c > 0 : \int_S \Phi(c^{-1}F(s), s) m(ds) \leq 1\}. \quad (42)$$

It is easy to see that $X = \int_S F dM$ can be described by (1) in which Q is the image of the measure $\theta(dx, s)m(ds)$ under the map $(s, x) \rightarrow F(s)x$. By the same argument used in the proof of (37) we get

$$(0.125)\|F\|_\Phi \leq E\| \int_S F dM \| \leq (2.125)\|F\|_\Phi. \quad (43)$$

for every F satisfying (38).

Remark 2.1 Stochastic integrals in which the integrand is real valued and the random measure is Hilbert space valued and vice versa are special cases of (39), so that (43) holds for them also.

3 Examples

Let X , Q and Q_r be as given in Section 1. In particular let

$$Q_r(dx) = C \left(\frac{1}{x^{p+1}} I_{[0 < x < 1]} + \frac{1}{x^{q+1}} I_{[1 \leq x < \infty]} \right) dx \quad (44)$$

$p, q \in (1, 2)$. It is elementary to verify that the value of l for which (3) and (5) is satisfied is given by the solution of one of the following equations

$$\frac{1}{l^2} \left(\frac{p-q}{(2-p)(2-q)} \right) + \frac{1}{l^q} \left(\frac{1}{(q-1)(2-q)} \right) = \frac{1}{C} \quad (45)$$

or

$$\frac{1}{l} \left(\frac{p-q}{(p-1)(q-1)} \right) + \frac{1}{l^p} \left(\frac{1}{(p-1)(2-p)} \right) = \frac{1}{C}. \quad (46)$$

(The value of l is unique. If it turns out that $l > 1$, then this value satisfies (45) and (46) has no solution. If $l < 1$ the opposite occurs. When $l = 1$, both equations are the same.)

In particular, if $p = q$ and X a Hilbert space valued infinitely divisible random variable with $Q_r(dx)$ given by (44), then by Theorem 1.1

$$(0.25) \left(\frac{C}{(2-p)(p-1)} \right)^{1/p} \leq E\|X\| \leq (2.125) \left(\frac{C}{(2-p)(p-1)} \right)^{1/p}. \quad (47)$$

If X is symmetric, the constant 2.125 in the upper bound of (47) can be decreased to 1.25.

There is a well known canonical way to define a mean zero, strictly p -stable, H -valued random vector X so that it has a nice characteristic function. For $1 < p < 2$ we set

$$E \exp i \langle y, X \rangle = \exp \left(- \int_{\partial U} |\langle y, \theta \rangle|^p \left(1 - i \tan \frac{\pi p}{2} \operatorname{sgn} \langle y, \theta \rangle \right) \sigma(d\theta) \right) \quad (48)$$

where σ is a finite measure on the unit sphere ∂U of H , called spectral measure of X . This equation comes from (1) with the Lévy measure Q of X written in spherical coordinates as

$$Q(dr, d\theta) = c_p r^{-p-1} dr \sigma(d\theta) \quad r > 0 \quad \theta \in \partial U. \quad (49)$$

Using (1), (49) and the identities

$$\int_0^\infty (e^{ir} - 1 - ir) r^{-1-p} dr = \Gamma(-p) e^{-i\pi p/2} \quad (50)$$

and

$$\int_0^\infty (e^{-ir} - 1 + ir) r^{-1-p} dr = \Gamma(-p) e^{i\pi p/2} \quad (51)$$

one can easily derive (48) and see that

$$c_p = - \frac{1}{\cos(\pi p/2) \Gamma(-p)}. \quad (52)$$

It follows from (47) that

$$(0.25)k_p[\sigma(\partial U)]^{1/p} \leq E\|X\| \leq (2.125)k_p[\sigma(\partial U)]^{1/p} \quad (53)$$

where

$$\begin{aligned} k_p &= \left(\frac{c_p}{(2-p)(p-1)} \right)^{1/p} \\ &= \left(\frac{p}{(2-p)\Gamma(2-p) \sin \frac{\pi(p-1)}{2}} \right)^{1/p}. \end{aligned} \quad (54)$$

Note that k_p is a continuous function of $p \in (1, 2)$; $k_p \sim (2/\pi)(p-1)^{-1}$ as p decreases to 1 and $k_p \rightarrow \sqrt{2}$ as p increases to 2.

We now use Theorem 1.2 to obtain a lower bound for $E\|X\|$ which holds under fairly general assumptions.

Lemma 3.1 *Let X and ψ_r be associated as in (18). Assume that $\psi_r(u)/u^2$ is decreasing as u increases and that $\psi(u)$ is increasing as u increases. Then*

$$E\|X\| \geq \frac{1}{4\psi_r^{-1}(5/2)}. \quad (55)$$

Proof Under the hypotheses on ψ_r we see that $\zeta(t) \geq \psi_r(1/t)$. Therefore, if $\psi_r(1/t_*) = 5/2$, the result follows from Theorem 1.2.

Suppose that ψ_r is regularly varying at zero with index $1 < p < 2$. Then

$$\lim_{t \rightarrow \infty} \frac{\zeta(t)(p-1)}{\psi_r(1/t)} = 1. \quad (56)$$

Therefore, heuristically, when $\psi_r^{-1}(5(p-1)/2)$ is very small

$$E\|X\| \geq \frac{1-\epsilon}{4\psi_r^{-1}(5(p-1)/2)} \quad (57)$$

for all $\epsilon > 0$. A similar statement can be made when ψ_r is regularly varying at infinity with index $1 < p < 2$. In this case

$$\lim_{t \rightarrow 0} \frac{\zeta(t)(p-1)}{\psi_r(1/t)} = 1. \quad (58)$$

Therefore, heuristically, when $\psi_r^{-1}(5(p-1)/2)$ is very large

$$E\|X\| \geq \frac{1-\epsilon}{4\psi_r^{-1}(5(p-1)/2)} \quad (59)$$

for all $\epsilon > 0$. Both these statement can be made precise with specific examples.

Consider the stochastic integral in (34). Define

$$\phi_1(u) = u^2 I(u < 1) + u I(u \geq 1), \quad u \geq 0. \quad (60)$$

Clearly

$$\phi_1(u) \leq \phi_0(u) \leq 2\phi_1(u) \quad (61)$$

where ϕ_0 is given in (31). Set

$$\tilde{\phi}(y, s) \stackrel{def}{=} \int_H \phi_1(|\langle y, x \rangle|) \theta(dx, s), \quad y \in H, \quad s \in S. \quad (62)$$

It follows from Theorem 2.1 that

$$(0.125)\|\tilde{f}\|_{\tilde{\phi}} \leq E|\int_S \langle \tilde{f}, dM \rangle| \leq (4.25)\|\tilde{f}\|_{\tilde{\phi}} \quad (63)$$

where $\|\tilde{f}\|_{\tilde{\phi}}$ is defined as in (32).

We proceed to evaluate $\|\tilde{f}\|_{\tilde{\phi}}$ for M in (28) with $\theta(dx, s)m(ds) = \theta(dr, d\theta, s)m(ds) = Cr^{-p-1}dr\sigma(d\theta)m(ds)$, similar to (49). A simple calculation shows that

$$\|\tilde{f}\|_{\tilde{\phi}} = \inf\{v > 0 : D_p \int_S \int_{\partial U} |\langle \frac{\tilde{f}(s)}{v}, \theta \rangle|^p d\sigma(\theta) m(ds) \leq 1\} \quad (64)$$

where $D_p = C/((2-p)(p-1))$. Consequently

$$\|\tilde{f}\|_{\tilde{\phi}} = \left(D_p \int_S \int_{\partial U} |\langle f(s), \theta \rangle|^p d\sigma(\theta) m(ds) \right)^{1/p} \quad (65)$$

or, equivalently

$$\|\tilde{f}\|_{\tilde{\phi}} = \left(D_p \int_S \|f(s)\|^p \left(\int_{\partial U} |\langle \frac{f(s)}{\|f(s)\|}, \theta \rangle|^p d\sigma(\theta) \right) m(ds) \right)^{1/p}. \quad (66)$$

We give two specific examples. Let $f_k(s)$ be the k -th coordinate of $f(s)$ and $e_k = (0, \dots, 1, \dots)$ be the unit vector in the k -th coordinate. Suppose that the measure σ assigns mass $p_k/2$ to both e_k and $-e_k$. Then it follows from (65) that

$$\|\tilde{f}\|_{\tilde{\phi}} = \left(D_p \sum_{k=1}^{\infty} \left(\int_S |f_k(s)|^p m(ds) \right) p_k \right)^{1/p}. \quad (67)$$

For the second example let σ be uniform measure of mass one on the unit sphere in \mathbf{R}^n . It follows from (66) that

$$\|\tilde{f}\|_{\tilde{\phi}} = \left(\frac{D_p E|g|^p}{E|g|^{n+p-1} E|g|^{n-1}} \int_S \|f(s)\|^p m(ds) \right)^{1/p} \quad (68)$$

where g is a normal random variable with mean zero and variance one.

To derive (68) we calculate $\int_{\partial U} |\langle \frac{f(s)}{\|f(s)\|}, \theta \rangle|^p d\sigma(\theta)$. Clearly this integral remains the same if we rotate the unit sphere in \mathbf{R}^n . Thus it remains the same for any unit vector $f(s)/\|f(s)\|$ in \mathbf{R}^n . We take e_1 for this vector.

Let $\xi = (\xi_1, \dots, \xi_n)$ where $\{\xi_i\}_{i=1}^n$ are independent normal random variables with mean zero and variance one. Note that

$$\begin{aligned} E|\xi_1|^p &= (2\pi)^{-n/2} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} |\langle e_1, \xi \rangle|^p e^{-\|\xi\|^2/2} d\xi \\ &= (2\pi)^{-n/2} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \|\xi\|^p |\langle e_1, \frac{\xi}{\|\xi\|} \rangle|^p e^{-\|\xi\|^2/2} d\xi. \end{aligned} \quad (69)$$

Writing this in polar coordinates we get

$$\begin{aligned} E|\xi_1|^p &= (2\pi)^{-n/2} \int_0^{\infty} r^{p+n-1} e^{-r^2/2} dr \int_{\partial U} |\langle e_1, \theta \rangle|^p d\theta \\ &= 2^{-(n+1)/2} \pi^{-(n-1)/2} E|\xi_1|^{n+p-1} \int_{\partial U} |\langle e_1, \theta \rangle|^p d\theta. \end{aligned} \quad (70)$$

Consequently

$$\int_{\partial U} |\langle e_1, \theta \rangle|^p d\theta = \frac{2^{(n+1)/2} \pi^{(n-1)/2} E|\xi_1|^p}{E|\xi_1|^{n+p-1}}. \quad (71)$$

Taking $p = 0$ in (71) gives the surface area of the unit sphere in \mathbf{R}^n . Dividing (71) by this number gives $\int_{\partial U} |\langle \frac{f(s)}{\|f(s)\|}, \theta \rangle|^p d\sigma(\theta)$ and hence (68).

It is not hard to obtain such definite results for integrals with respect to other infinitely divisible random measures than stable ones. It only requires a good estimate for ϕ in (62), which is not hard to obtain when the Lévy measure θ associated with M in (28) has nice properties, such as being regularly varying at zero and infinity.

References

- [1] V.H. de la Peña and E. Giné. *Decoupling. From dependence to independence*. Springer-Verlag, 1999.
- [2] N. Dunford and J.T. Schwartz. *Linear Operators, Part I: General Theory*. Wiley, 1958.
- [3] I.I. Gihman and A.V. Skorohod. *The Theory of Stochastic Processes, Vol. 2*, Springer, Berlin, 1975.
- [4] M.J. Klass. Precision bounds for the relative error in the approximation of $E|S_n|$ and extensions. *Ann. Probab.*, 8: 350–367, 1980.
- [5] S. Kwapien and W.A. Woyczyński. *Random Series and Stochastic Integrals: Single and Multiple. Probability and its Applications* Birkhauser, 1992.
- [6] M. B. Marcus and J. Rosiński. Continuity of stochastic integrals with respect to infinitely divisible random measures. Preprint.
- [7] K. R. Parthasarathy. *Probability measures on metric spaces*, Academic Press, New York, 1967.