

Revista Mexicana de Astronomía y Astrofísica

Revista Mexicana de Astronomía y Astrofísica
Universidad Nacional Autónoma de México
rmaa@astroscu.unam.mx
ISSN (Versión impresa): 0185-1101
MÉXICO

2002
S. Durand / D. Nürnberger / J. Köppen
PN G291.4-00.3: A NEW TYPE I PLANETARY NEBULA
Revista Mexicana de Astronomía y Astrofísica, volumen 012
Universidad Nacional Autónoma de México
Distrito Federal, México
p. 162

Red de Revistas Científicas de América Latina y el Caribe, España y Portugal

Universidad Autónoma del Estado de México

reDalyC
LA BIBLIOTECA CIENTÍFICA EN LÍNEA
<http://redalyc.uaemex.mx>

PN G291.4–00.3: A NEW TYPE I PLANETARY NEBULA¹

S. Durand,² D. Nürnberger,³ and J. Köppen⁴

In the vicinity of the southern hemisphere giant H II region NGC 3603 we discovered a new planetary nebula: PN G291.4–00.3 located at $RA_{J2000.0} = 11^{\text{h}}14^{\text{m}}32^{\text{s}}.1 \pm 0^{\text{s}}.3$ and $DEC_{J2000.0} = -61^{\circ}00'02'' \pm 1''$. Monochromatic images reveal a central ring-like structure with onsets of arc-like filaments which might outline a bipolar outflow. Optical imaging and spectroscopy confirm that the emission line object found on WFI images is a genuine planetary nebula, and that it is a high-excitation bipolar Type I object.

PN G291.4–00.3 (Fig. 1) was discovered serendipitously on wide field CCD exposures centered on the giant H II region NGC 3603. These images were obtained with the Wide Field Imager mounted at the ESO/MPG 2.2-m telescope on La Silla under moderate seeing conditions.

Its appearance consists of a roughly north-south elongated ring-like structure with short onsets of spiral-like arms at the northern and southern tips. The central ring appears to be tilted by 50° to 55° against the plane of sky. Its diameter is about $5''.4$ along the major axis. The arc- or spiral-like shaped filaments might be indicative for a wide-angle hourglass-shaped bipolar outflow (P.A. $\sim 80^{\circ}$).

Spectroscopic observations were performed using EFOSC 2 at the ESO 3.6-m telescope. The spectrum




Fig. 1. $H\alpha$ narrow band image obtained with the WFI mounted at the ESO/MPG 2.2-m telescope.

exhibits a very strong He II $\lambda 4686$ line, but also [S II] lines. Plasma diagnostics shows helium and nitrogen are enriched, while oxygen is depleted. Thus PN G291.4–00.3 is a high-excitation bipolar planetary nebula of Type I. The measured extinction and the radial velocity obtained from the $H\alpha$ and [N II] lines of $+5 \pm 1.5 \text{ km s}^{-1}$ indicate a distance of about 5.5 kpc.

S. Durand thanks PRONEX/Finep for financial assistance. This work is described in greater detail in Nürnberger et al. 2001.

REFERENCES

Nürnberger, D., Durand, S., Köppen, J., Stanke Th., Sterzik, M. & Els, S., 2001, A&A, 377, 241

¹Based on observations obtained at the European Southern Observatory, La Silla, Chile.

²IAG/USP, Departamento de Astronomia, Av. Miguel Stéfano 4200, 04301-904 São Paulo, Brasil (durand@iagusp.usp.br).

³Institut für Theoretische Physik und Astrophysik, Am Hubland, 97074 Würzburg, Germany (nurnberg@astro.uni-wuerzburg.de).

⁴Observatoire Astronomique de Strasbourg, 11 rue de l'Université, 67000 Strasbourg, France (koppen@astro.u-strasbg.fr).