

Revista Mexicana de Astronomía y Astrofísica

Revista Mexicana de Astronomía y Astrofísica
Universidad Nacional Autónoma de México
rmaa@astroscu.unam.mx
ISSN (Versión impresa): 0185-1101
MÉXICO

2002
A. Bullejos / M. Rosado
KINEMATICS AROUND THE NON-THERMAL SUPERBUBBLE IN IC 10
Revista Mexicana de Astronomía y Astrofísica, volumen 012
Universidad Nacional Autónoma de México
Distrito Federal, México
p. 254

Red de Revistas Científicas de América Latina y el Caribe, España y Portugal

Universidad Autónoma del Estado de México

reDalyC
LA BIBLIOTECA CIENTÍFICA EN LÍNEA
<http://redalyc.uaemex.mx>

KINEMATICS AROUND THE NON-THERMAL SUPERBUBBLE IN IC 10

A. Bullejos¹ and M. Rosado¹

We study the kinematics around a non-thermal superbubble found by Yang & Skillman. Considering the H II and [S II] profiles, we find that between 3 and 6 supernovae are required to form this superbubble.

IC 10 may well be the nearest example of a blue compact dwarf galaxy (BCD). Among the Local Group star-forming dwarf galaxies, IC 10 is conspicuous for having the highest surface density of WR stars (11.2 kpc^{-2} , Richer et al. 2000). The presence of so many WR stars and the high H α luminosity emphasize that IC10 is currently undergoing a burst of star formation that began at least 10 Myr ago. Observations of 21-cm emission from neutral hydrogen reveal that IC 10 consists of an inner disk embedded in an extended, complex, counter-rotating envelope (Shostak & Skillman 1989). This leads Wilcots & Miller (1998) to conclude that IC 10 is still in its formative stage. These H I observations make clear the youth of the current star formation episode for there is a notable lack of interstellar medium structures attributable to supernovae (Wilcots & Miller 1998). The galaxy size in the light of H α is 5×3 arcmin while in H I it is 18×15 arcmin. All optical studies of IC 10 are hampered by the large foreground reddening due to its position near the plane of our galaxy ($l = 119^\circ$, $b = -3^\circ$).

Our observations of the non-thermal superbubble, discovered by Yang & Skillman (1993), were obtained in the light of H α and [S II] with the scanning Fabry-Perot Interferometer, PUMA. We suppose that the energy in the ionized interstellar medium comes from the energy injected by stellar winds and supernovae. Considering both the H α and [S II] profiles, we find a typical expansion velocity of 50 to 70 km s^{-1} . Using a distance of 661 kpc (Sakai et al. 1999) and a diameter of $41.5''$ inferred from Yang & Skillman (1993), we calculate a radius of 66 pc for the superbubble. Adopting a density of $1.1 \text{ atoms cm}^{-3}$ (Yang & Skillman 1993), the mass swept out by the superbubble is $4.87 \times 10^{37} \text{ g}$ (2.45×10^4 solar masses).

¹Instituto de Astronomía, UNAM, Apdo Postal 70-264, Ciudad Universitaria, México D.F., México (almudena,maragarit@astroscu.unam.mx).




Fig. 1. A close-up of the region of the non-thermal superbubble in H α with the Yang & Skillman, (1993) iso-contours of the 49-cm flux superposed. This image was formed by summing up the velocity channels containing emission from IC 10. The heliocentric velocity of this galaxy is -344 km s^{-1} (de Vaucouleurs et al. 1991).

With these values, the kinetic energy obtained is 6.09×10^{50} to 11.9×10^{50} erg. Supposing that 20% of a supernova's thermal energy is converted into kinetic energy and that each of them injects 10^{51} erg, then from 3 to 6 supernova are required to form this superbubble.

REFERENCES

- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H. G., et al. 1991, in *The Third Reference Catalog of Bright Galaxies* (New York: Springer)
- Richer, M. G. et al. 2001, *A&A*, 370, 34
- Sakai, S., Madore, B.F., & Freedman, W. L. 1999, *ApJ*, 511, 671
- Shostak, G. S., & Skillman, E. D. 1989, *A&A*, 214, 33
- Wilcots, E. M., Miller, B. W. 1998, *AJ*, 116, 2362
- Yang, H., & Skillman, E. D. 1993, *AJ*, 106, 1448