

Revista Mexicana de Astronomía y Astrofísica

Revista Mexicana de Astronomía y Astrofísica
Universidad Nacional Autónoma de México
rmaa@astroscu.unam.mx
ISSN (Versión impresa): 0185-1101
MÉXICO

2003

M. Relaño / J. E. Beckman / M. Rozas / A. Zurita
OBSERVATIONS OF MASSIVE SUPERSONIC OUTFLOWS IN HIGHLY LUMINOUS
H II REGIONS

Revista Mexicana de Astronomía y Astrofísica, número 015
Universidad Nacional Autónoma de México
Distrito Federal, México
p. 205

Red de Revistas Científicas de América Latina y el Caribe, España y Portugal

Universidad Autónoma del Estado de México

reDalyC
LA BIBLIOTECA CIENTÍFICA EN LÍNEA
<http://redalyc.uaemex.mx>

OBSERVATIONS OF MASSIVE SUPERSONIC OUTFLOWS IN HIGHLY LUMINOUS H II REGIONS

M. Relaño,¹ J. E. Beckman,^{1,2} M. Rozas,³ and A. Zurita⁴

Using H α emission line profiles we detect outflowing ionized gas at velocities of order 50 km s^{-1} from representative luminous H II regions in two sample disk galaxies, NGC 3359 and NGC 1530, and calculate the physical parameters of the outflowing material.

The study is based on two types of observations: *continuum-subtracted H α images*, whose calibration and analysis yields a complete catalog of H α luminosities for the H II regions of a disk galaxy (Rozas et al. 2000), and *Fabry-Perot data cubes* of relative intensity, position and wavelength obtained in H α for the same galaxies. The cube yields spectral emission profiles for each pixel in the galaxy image. We have selected a sample of spherical and isolated H II regions and integrated the profiles over the emission zone of the region, yielding an integrated profile which has been analyzed in terms of Gaussian components.

We found a central peak plus red and blue high-velocity, low-intensity Gaussian components. In Figure 1 we show the plot of the non-thermal component of the line width of the central emission peaks (derived by subtracting off in quadrature from the observed width the thermal, instrumental, and natural widths) against H α luminosity of the region producing the line. The phenomenology of the red and blue observed components suggests a shell in expansion, with velocities of order $\sim 50 \text{ km s}^{-1}$ (see Figure 2). The physical parameters of the outflowing material show that the wings can be accounted for either in terms of the interaction of the combined winds of the OB stars (Dyson, Williams, & Redman 1995) with the surrounding gas or by acceleration of the gas via stellar radiation (Elitzur & Ivezić 2001). For further details see Relaño, Beckman, & Dyson (2003).




Fig. 1. The non-thermal component of the H α emission line widths from the selected isolated H II in NGC 3359 and NGC 1530 vs. the H α luminosity of the H II regions.




Fig. 2. Mean expansion velocity of the H II region shells vs. H α luminosity for the same selected H II regions as in Figure 1.

REFERENCES

- Dyson, J. E., Williams, R. J. R., & Redman, M. P. 1995, MNRAS, 277, 700
 Elitzur, M., & Ivezić, Ž. 2001, MNRAS, 327, 403
 Relaño, M., Beckman, J. E., & Dyson, J. E. 2003, RevMexAA(SC), 15, 281 (this volume)
 Rozas, M., Zurita, A., Beckman, J. E., & Pérez, D. 2000, A&AS, 142, 259

¹Instituto de Astrofísica de Canarias, C/Vía Láctea s/n, 38200-Tenerife, Spain (mpastor@iac.es).

²Consejo Superior de Investigaciones Científicas, CSIC, Spain (jeb@iac.es).

³Observatorio Astronómico Nacional, UNAM, Ensenada, B. C., Mexico (maite@astrosen.unam.mx).

⁴Isaac Newton Group of Telescopes, UK (azurita@ing.iac.es).