

Revista Mexicana de Astronomía y Astrofísica
Universidad Nacional Autónoma de México
rmaa@astroscu.unam.mx
ISSN (Versión impresa): 0185-1101
MÉXICO

2004

T. Di Salvo / R. Iaria / M. Méndez / L. Burderi / N. R. Robba / L. Stella / M. Van der Klis
A BROAD IRON LINE IN THE CHANDRA/HETG SPECTRUM OF 4U 1705{44
Revista Mexicana de Astronomía y Astrofísica, número 020
Universidad Nacional Autónoma de México
Distrito Federal, México 8

Red de Revistas Científicas de América Latina y el Caribe, España y Portugal

Universidad Autónoma del Estado de México

<http://redalyc.uaemex.mx>



A BROAD IRON LINE IN THE CHANDRA/HETG SPECTRUM OF 4U 1705–44

T. Di Salvo,¹ R. Iaria,² M. Méndez,³ L. Burderi,⁴ N.R. Robba,² L. Stella,⁴ and M. van der Klis¹

We present preliminary results from a Chandra 30 ks observation of the atoll source 4U 1705–44. In particular we concentrate on the study of the iron $K\alpha$ line, using the HEG spectrometer. The iron $K\alpha$ line at ~ 6.6 keV is found to be intrinsically broad ($FWHM \sim 1.7$ keV); its width can be explained by reflection from a cold accretion disk extending down to ~ 17 km from the neutron star center or by Compton broadening in the hotter ($\sim 3 - 4$ keV) corona.

Broad emission lines (FWHM up to ~ 1 keV) at energies in the range 6.4 – 6.7 keV are often observed in the spectra of low-mass X-ray binaries hosting an old accreting neutron star. These lines are identified with the $K\alpha$ radiative transitions of iron at different ionization states. These features are powerful tools to investigate the structure of the accretion flow close to the central source; in particular, important information can be obtained from the detailed spectroscopy of the iron $K\alpha$ emission line and absorption edge, since they are determined by the ionization state, geometry and velocity field of the reprocessing plasma. 4U 1705–44 is an atoll source (see Hasinger & van der Klis 1989) which also shows type-I X-ray bursts. We selected this source, in which a broad (1.1 keV FWHM) iron emission line at 6.5 keV has been reported (White et al. 1986; Barret & Olive 2002), for a Chandra observation in order to study the iron line profile and to discriminate among the various models that have been proposed to explain the large line width.

Our 30 ks observation of 4U 1705–44 was performed using the High-Energy Transmission Grating Spectrometer (HETGS) on board of Chandra starting on 2001 July 1. The HETGS consists of two transmission gratings, the Medium-Energy Grating (MEG) and the High-Energy Grating (HEG). It allows high-resolution spectroscopy from 1.2 to 31 Å

(0.4–10 keV) with a peak spectral resolution at 12 Å of $\Delta\lambda/\lambda \sim 1000$ for HEG first order. In our analysis, we utilize the HEG first-order spectrum, which is less affected by pile-up (less than 10%) with respect to the MEG first-order spectrum.

We fit the HEG spectrum of 4U 1705–44 to a continuum model. The best fit model consists of a blackbody plus the Comptonization model *comptt* (Titarchuk 1994), modified by absorption from neutral matter, which gives a $\chi^2_{\text{red}}(d.o.f.)$ of 1.64(846). The addition of a broad ($\sigma \sim 0.7$ keV) Gaussian line centered at 6.57 keV proves necessary, giving $\chi^2_{\text{red}}(d.o.f.) = 1.25(843)$. The high energy resolution of the HEG shows that the line is intrinsically broad ($FWHM \sim 1.7$ keV). Therefore the most probable origin of this line is an accretion disk (in this case the large width of the line would be due to Doppler and relativistic smearing effects) or from a hot corona (in this case the large width of the line would be due to Compton broadening).

In the case the line is produced by reflection in an accretion disk we estimate that the required inner radius of the disk is $8 R_g$ or ~ 17 km for a $1.4 M_\odot$ neutron star. The inclination angle is $\sim 80^\circ$, which seems to be quite high for this source. Alternatively, Comptonization in the corona could explain the large width of the line. Detailed calculations give $\sigma_{\text{Fe}} = 0.019 E_{\text{Fe}} \tau_T (1 + 0.78 kT_e)$, where τ_T is the Thomson optical depth and kT_e is in keV (Kallman & White 1989, see also Brandt & Matt 1994). Assuming an electron temperature $kT_e = 3.8$ keV, as derived from the fit of our data to the Comptonization model, we can explain the width of the iron line for a Thomson optical depth of $\tau_T \sim 1.4$. Therefore, it is possible that the line is produced in the outer region of the Comptonizing corona, where the optical depth decreases (assuming that the temperature remains constant).

REFERENCES

- Barret, D., & Olive, J.F., 2002, ApJ, 576, 391
 Brandt, W.M., & Matt, G., 1994, MNRAS, 268, 1051
 Hasinger, G., & van der Klis, M., 1989, A&A, 225, 79
 Kallman, T., & White, N.E., 1989, ApJ, 341, 955
 Titarchuk, L., 1994, ApJ, 434, 570
 White, N.E., et al., 1986, MNRAS, 218, 129

¹Astronomical Institute "Anton Pannekoek," University of Amsterdam and Center for High-Energy Astrophysics, Kruislaan 403, NL 1098 SJ Amsterdam, the Netherlands.

²Dipartimento di Scienze Fisiche ed Astronomiche, Università di Palermo, Italy.

³National Institute for Space Research, SRON, Utrecht, the Netherlands.

⁴Osservatorio Astronomico di Roma, Italy.