

Revista Mexicana de Astronomía y Astrofísica
Universidad Nacional Autónoma de México
maa@astroscu.unam.mx
ISSN (Versión impresa): 0185-1101
MÉXICO

2008
J. P. Lenain
MULTIWAVELENGTH MODELING OF TEV AGN OBSERVED BY H.E.S.S.
Revista Mexicana de Astronomía y Astrofísica, abril, volumen 032
Universidad Nacional Autónoma de México
Distrito Federal, México
pp. 28-30

Red de Revistas Científicas de América Latina y el Caribe, España y Portugal

Universidad Autónoma del Estado de México

<http://redalyc.uaemex.mx>



MULTIWAVELENGTH MODELING OF TEV AGN OBSERVED BY H.E.S.S.

J.-P. Lenain¹

RESUMEN

El experimento “Sistema Estereoscópico de Alta Energía” (High Energy Stereoscopic System, H.E.S.S.) consiste de un arreglo de telescopios Čerenkov de rayos γ terrestres situado en Namibia que ha detectado ya muchos objetos extragalácticos, cuyos corrimientos al rojo varían de $z=0.00183$ a $z=0.2$, probablemente mayores. Dado el creciente rendimiento de los telescopios Čerenkov, ahora es posible investigar estos objetos en escalas de tiempo muy pequeñas en rayos γ , permitiendo el estudio de regiones que se piensa están muy cercanas a los agujeros negros supermasivos centrales. Además, H.E.S.S. ha confirmado la emisión de rayos γ en M 87, por lo que se convierte entonces en la primera fuente extragaláctica observada en TeV que no es un blazar. Entre los blazares, los TeV BL Lacs son los objetos más estimulantes para evaluar los modelos de emisión del jet y examinar los procesos de aceleración de partículas. La investigación de los blazares con H.E.S.S. revela también distintos comportamientos temporales entre ellos. Unos objetos manifiestan un flujo muy variable en rayos-X con pequeñas variaciones en los rayos γ , mientras que otros muestran el comportamiento inverso. La interpretación de esas diferencias es desconcertante. Las observaciones a muy altas energías proveen también de medidas indirectas del fondo infrarrojo extragaláctico (EBL). La interpretación de la emisión en rayos γ de radiogalaxias como M 87 en términos de blazares no-alineados, y la comprensión de las propiedades del EBL representan nuevos desafíos encontrados por H.E.S.S. al efectuar observaciones de fuentes extragalácticas.

ABSTRACT

The High Energy Stereoscopic System (H.E.S.S.) experiment, a ground-based γ -ray Čerenkov telescope array located in Namibia, has now detected many extragalactic objects, which redshifts range from $z=0.00183$ up to $z=0.2$, possibly more. With the increasing performances of Čerenkov telescopes, it now becomes possible to probe these objects at small timescales in γ -ray, allowing the study of regions thought to be very close to the central supermassive black holes. Furthermore, H.E.S.S. has confirmed a γ -ray emission from M 87, which is thus the first extragalactic source seen at the TeV range that is not a blazar. Among blazars, TeV BL Lacs are the most challenging objects to test the jet emission models and to shed light on particle acceleration mechanisms. The study of blazars with H.E.S.S. also revealed various temporal behaviors among them. Some objects present a highly variable X-ray flux with small variation of the γ -ray, while others show the inverse behavior. The interpretation of such differences is puzzling. Observations at very high energies also bring indirect measurements of the infrared extragalactic background light (EBL). The interpretation of γ -ray emission of radiogalaxies such as M 87 in terms of misaligned blazars and the understanding of the properties of the EBL represent new challenges brought by H.E.S.S. observations of extragalactic sources.

Key Words: galaxies: active — galaxies: general — galaxies: individual (M 87, PKS 2155–304)

1. DETECTING VERY HIGH ENERGY γ -RAY FROM THE GROUND WITH H.E.S.S.

The High Energy Stereoscopic System (H.E.S.S.) experiment² consists of four imaging atmospheric Čerenkov telescopes (see Figure 1), fully operating since 2004, located in the Khomas highlands in Namibia. It operates in the very high energy (VHE) domain, in the GeV–TeV range. The analysis of the Čerenkov light of a single air shower allows the de-




Fig. 1. The H.E.S.S. array located in Namibia.

termination of the energy of the incident particle. Furthermore, the stereoscopy technique allows a precise determination of the arrival direction, as well as a good discrimination between hadrons and photons (Aharonian et al. 2004).

¹LUTH, Observatoire de Paris, CNRS, Université Paris Diderot; 5 Place Jules Janssen, 92190 Meudon, France (jean-philippe.lenain@obspm.fr).

²<http://www.mpi-hd.mpg.de/hfm/HESS/>.




Fig. 2. Spectral energy distribution of PKS 2155–304. Only simultaneous observations are labeled. The solid line is the hadronic model, the dotted and dashed lines are the same leptonic models with different assumptions, from Aharonian et al. (2005).

2. TEV BLAZARS

Blazars are active galactic nuclei (AGN) with the jet closely aligned with the line of sight, thus having their intensity strongly amplified by relativistic effects. Their spectral energy distribution usually shows two bumps, one in the UV–X range thought to be due to synchrotron emission, and the other one in the X– γ range ascribed to inverse Compton effects in leptonic models.

Historically, Fossati et al. (1998) emphasized the existence of two classes of objects: low-peaked BL Lac (LBL) and high-peaked BL Lac (HBL) for which the synchrotron bump peaks at low ($\sim 10^{13}$ Hz), respectively high ($\sim 10^{16}$ Hz), frequency. In this framework, the first TeV sources were searched among the HBL (Costamante & Ghisellini 2002), hence all (except one) AGN detected by H.E.S.S. belong to this class. M 87 (FRI) and BL Lac (LBL) itself are the only non-HBL objects detected in the VHE range up to now by H.E.S.S. and MAGIC, respectively (Aharonian et al. 2006c).

Among these sources, different X/ γ spectral variability patterns can be found. For instance, the X-ray peak observed in H2356–309 strongly constrains leptonic models. H2356–309 presents low flux variations in X-ray, with rather constant spectral shape, with high flux variations in γ -ray. On the contrary, PKS 2005–489 has a quasi constant γ -ray emission along with huge flux variations in X-ray coming with highly varying spectral shape.

PKS 2155–304, a well known variable object, was observed in a long quasi-quiet state (Aharonian et al. 2005) during multiwavelength campaigns in 2003, 2004 and 2005 (see Figure 2). Observations




Fig. 3. Lightcurve of PKS 2155–304 on July 28, 2006. Variability is seen down to 2 minutes scale, from Aharonian et al. (2007).

in July 2006 revealed a very active state over two weeks, with in particular a big flare in VHE up to 15 times the Crab flux with variability timescale of the order of 2 minutes (Aharonian et al. 2007). This is the fastest variability ever seen in PKS 2155–304 at any wavelength and in any blazar (see Figure 3).

3. M 87 AS A MISALIGNED BLAZAR?

Between 2003 and 2006, H.E.S.S. observed the FRI radiogalaxy M 87 (Aharonian et al. 2006c), thus confirming the detection of the HEGRA collaboration. Flux variability was observed, without significant spectral variability, with timescales down to 2 days, thus excluding the radio lobes or the extended kiloparsec jet as VHE emitters. It is still unclear whether the emission comes from the nucleus or the knot *HST-1*.

We assume that the VHE emitting zone is located in the broadened jet formation region, predicted by magnetohydrodynamics models and observed in VLBI in the case of M 87 (see, e.g. Junor et al. 1999). It is then possible to explain the γ -ray radiation in terms of synchrotron self-Compton (SSC) processes, as for blazars and with similar electron energy distribution, with rather moderate Lorentz factor, confirming the thought that M 87 could be a misaligned BL Lac object.

The radiation model presented here relies on the previous work by Katarzyński et al. (2001). Here, we assumed that a cap of seven blobs of plasma is located at ~ 50 gravitational radii from the supermassive black hole, beyond the Alfvén surface predicted by magnetohydrodynamic (MHD) models, see e.g. (McKinney 2006). The population of non-thermal electrons of the blobs, while crossing the Alfvén surface, is accelerated and then radiates through synchrotron emission and inverse Compton scattering of electrons on the synchrotron photons. While ra-




Fig. 4. Spectral energy distribution of M87. The black line in the radio shows a model of extended inner jet, while the two lines from IR to VHE show a SSC model to account for the synchrotron and inverse Compton bumps, from Lenain et al. (2008).

diating in the broadened zone of the jet, some blobs are moving close to the line of sight, with differential Doppler boosting from one blob to the other, allowing to derive parameters similar to those of TeV blazars. Figure 4 shows the spectral energy distribution deduced from such a model of M87 accounting for the recent VHE H.E.S.S. observations. For more details, see (Lenain et al. 2008).

4. COSMOLOGICAL IMPLICATIONS: THE EBL AND THE γ HORIZON

The infrared extragalactic background light (EBL) consists of the sum of the starlight and dust emission of the galaxies through the history of the Universe, including starburst and active galaxies. It could also have an important contribution from the first stars, the so-called population III, which may have formed from primordial gas.

While traveling to the earth, VHE photons interact with infrared photons from the extragalactic background light (EBL). The EBL thus absorbs a part of the observed VHE flux. Relying on observations of nearby blazars, it was found that their intrinsic photon index is $\Gamma > 1.5$. It is then possible to indirectly derive measurements of this background




Fig. 5. Spectral energy distribution of the EBL, with the H.E.S.S. upper limit reported (*bold black line*), from Aharonian et al. (2006b).

through VHE observations of remote ($z \sim 0.2-0.3$) blazars (Aharonian et al. 2006b), see Figure 5.

It was found that the EBL flux seems to be lower than expected, thus allowing more distant VHE sources to be potentially detected. Furthermore the EBL seems to be dominated by galaxy population, mainly excluding stellar population III contribution.

REFERENCES

- Aharonian, F., et al. 2004, *Astropart. Phys.*, 22, 109
 ————. 2005, *A&A*, 442, 895
 ————. 2006a, *A&A*, 457, 899
 ————. 2006b, *Nature*, 440, 1018
 ————. 2006c, *Science*, 314, 1424
 ————. 2007, *ApJ*, 664, L71
 Albert, J., et al. 2007, *ApJ*, 666, L17
 Costamante, L., & Ghisellini, G. 2002, *A&A*, 384, 56
 Fossati, G., Maraschi, L., Celotti, A., Comastri, A., & Ghisellini, G. 1998, *MNRAS*, 299, 433
 Junor, W., Biretta, J. A., & Livio, M. 1999, *Nature*, 401, 891
 Katarzyński, K., Sol, H., & Kus, A. 2001, *A&A*, 367, 809
 Lenain, J.-P., Boisson, C., Sol, H., & Katarzyński, K. 2008, *A&A*, 478, 111
 McKinney, J. C. 2006, *MNRAS*, 368, 1561