

Simple Analytical Expression for Vector Hypernuclear Asymmetry in Nonmesonic Decay of ${}^5_{\Lambda}\text{He}$ and ${}^{12}_{\Lambda}\text{C}$

César Barbero,

*Departamento de Física, Facultad de Ciencias Exactas,
Universidad Nacional de La Plata, C.C. 67, 1900 La Plata, Argentina.*

Francisco Krmpotić,

*Instituto de Física, Universidade de São Paulo,
C. P. 66318, 05315-970 São Paulo, SP, Brazil,
Instituto de Física La Plata,
CONICET, 1900 La Plata, Argentina.*

and Alfredo P. Galeão

*Instituto de Física Teórica, Universidade Estadual Paulista,
Rua Pamplona 145, 01405-900 São Paulo, SP, Brazil.*

Received on 18 March, 2006

We present general explicit expressions for a shell-model calculation of the vector hypernuclear parameter in nonmesonic weak decay. We use a widely accepted effective coupling Hamiltonian involving the exchange of the complete pseudoscalar and vector meson octets ($\pi, \eta, K, \rho, \omega, K^*$). In contrast to the approximated formula widely used in the literature, we correctly treat the contribution of transitions originated from single-proton states beyond the s -shell. Exact and simple analytical expressions are obtained for the particular cases of ${}^5_{\Lambda}\text{He}$ and ${}^{12}_{\Lambda}\text{C}$, within the one-pion-exchange model. Numerical computations of the asymmetry parameter, a_{Λ} , are presented. Our results show a qualitative agreement with other theoretical estimates but also a contradiction with recent experimental determinations. Our simple analytical formulas provide a guide in searching the origin of such discrepancies, and they will be useful for helping to solve the hypernuclear weak decay puzzle.

Keywords: Hypernuclear decay; Asymmetry parameter; One-meson-exchange model

The free decay of a Λ hyperon occurs almost exclusively through the mesonic mode, $\Lambda \rightarrow \pi N$, emerging the nucleon with a momentum of about 100 MeV/c. Inside the nuclear medium ($p_F \sim 270$ MeV/c) this mode is Pauli blocked and, for all but the lightest Λ hypernuclei ($A \geq 5$), the weak decay is dominated by the nonmesonic channel, $\Lambda N \rightarrow NN$, with enough kinetic energy to put the two emitted nucleons above the Fermi surface. The nonmesonic hypernuclear weak decay (NMHWD) offers us a very unique opportunity to investigate strangeness-changing weak interaction between hadrons. Assuming that the NMHWD is saturated by the $\Lambda N \rightarrow NN$ mode, the transitions receive contributions either from neutrons ($\Lambda n \rightarrow nn$) and protons ($\Lambda p \rightarrow np$), with rates Γ_n and Γ_p , respectively ($\Gamma_{NM} = \Gamma_n + \Gamma_p$, $\Gamma_{n/p} = \Gamma_n/\Gamma_p$). Over the last three decades amount of theoretical and experimental effort has been invested in solving an interesting puzzle: the theoretical models cannot reproduce simultaneously the experimental values of $\Gamma_{n/p}$ and a_{Λ} .

From the experimental side there is actually an intense activity, as can be seen in the light of the experiments under way and/or planned at (KEK) [1], (FINUDA) [2] and Brookhaven National Laboratory (BNL) [3]. The preliminary results give a $\Gamma_{n/p}$ ratio value very close to 0.5 [4–6] and the measurements of a_{Λ} favor a negative value for ${}^{12}_{\Lambda}\text{C}$ and a positive value for ${}^5_{\Lambda}\text{He}$ [5–9]. On the other hand, from the pioneering work of Block and Dalitz [10] there have been many theoretical attempts dedicated to solving the puzzle. The earlier studies were based on the simplest model of the virtual pion exchange

[11]. This model naturally explains the long range part of the two body interaction, and it reproduces reasonably well the total decay rate but fails badly in reproducing the other observables. In order to achieve a better description, some improvements have been introduced: (i) models that include the exchange of different combinations of other heavier mesons, like η, K, ρ, ω and K^* [12]–[19], to reproduce the short range part of the interaction; (ii) analysis of the two nucleon (2N) stimulated process $\Lambda NN \rightarrow NNN$ [20, 21]; (iii) inclusion of interaction terms that violate the isospin $\Delta T = 1/2$ rule [22]–[24]; (iv) description of the short range baryon-baryon interaction in terms of quark degrees of freedom [25, 26]; (v) correlated (in the form of σ and ρ mesons) and uncorrelated two-pion exchanges [27]–[31]. We emphasize that none of these models give a fully satisfactory description of all the NMHWD observables simultaneously, even though a consistent (though not sufficient) increase of $\Gamma_{n/p}$ has been found.

All such models reproduce quite well Γ_{nm} , but seem to strongly underestimate the experimental n/p branching ratio, Γ_n/Γ_p . Besides, the measurements favor a negative value of a_{Λ} for ${}^{12}_{\Lambda}\text{C}$ and a positive one for ${}^5_{\Lambda}\text{He}$, meanwhile all existing calculations based on strict one-meson-exchange (OME) models [14–16, 18, 19, 26, 32, 33] find values between -0.73 and -0.19 for ${}^5_{\Lambda}\text{He}$ [34, 35] and, when results are available in the same model, very similar values for ${}^{12}_{\Lambda}\text{C}$.

Most calculations of the asymmetry parameter make use of an approximate formula (Eq. (18), below) which, however, is valid only for s -shell hypernuclei. Since an essential aspect




FIG. 1: Coordinate system for the calculation of decay strengths.

in the asymmetry puzzle presented above concerns the comparison of its values for ${}^5_{\Lambda}\text{He}$ and ${}^{12}_{\Lambda}\text{C}$, it would be of great interest to have an expression that is applicable to both cases. Besides, simple formulas containing kinematical factors and nuclear matrix elements separately, will provide a manageable and useful tool. They will be a guide in searching the

origin of discrepancies between theoretical calculations and experimental data.

Thus, we present in this paper a general formalism for the vector asymmetry parameter in nonmesonic decay and derive exact and simple analytical expressions, within the simplest one-pion exchange model (OPEM), for the particular cases of ${}^5_{\Lambda}\text{He}$ and ${}^{12}_{\Lambda}\text{C}$ hypernuclei.

Single- Λ hypernuclei produced in a (π^+, K^+) reaction end up with considerable vector polarization along the direction normal to the reaction plane, $\hat{n} = (\mathbf{p}_{\pi^+} \times \mathbf{p}_{K^+}) / |\mathbf{p}_{\pi^+} \times \mathbf{p}_{K^+}|$. The angular distribution of protons emitted in $\Lambda p \rightarrow np$ decay is given by [34]

$$\frac{d\Gamma(J_I M_I \rightarrow \hat{p}_2 t_p)}{d\Omega_{p_2}} = \frac{\Gamma_p}{4\pi} (1 + P_V A_V \hat{p}_2 \cdot \hat{n}), \quad (1)$$

where Γ_p is the full proton-induced decay rate, and

$$A_V = 3 \sqrt{\frac{J_I}{J_I + 1}} \frac{\sigma_1(J_I)}{\sigma_0(J_I)}, \quad (2)$$

is the *vector hypernuclear asymmetry*, with

$$\begin{aligned} \sigma_0(J_I) &= \sum_{M_I} \sigma(J_I M_I), \\ \sigma_1(J_I) &= \frac{1}{\sqrt{J_I(J_I + 1)}} \sum_{M_I} M_I \sigma(J_I M_I), \end{aligned} \quad (3)$$

being

$$\sigma(J_I M_I) = \int d\Omega_{p_1} \int dF \sum_{s_1 s_2 M_F} |\langle \mathbf{p}_1 s_1 t_n \mathbf{p}_2 s_2 t_p J_F M_F | V | J_I M_I \rangle_{\mathcal{A}}|^2. \quad (4)$$

Here V is the OME potential involving the complete pseudoscalar and vector meson octets (π , η , K , ρ , ω , K^*) which has been extensively discussed in Refs. [17, 18], $\mathbf{p}_1 s_1$ ($\mathbf{p}_2 s_2$) are the momenta and spin projections of the emitted neutron (proton) and

$$\int dF \dots = 2\pi \sum_F \int \frac{p_2^2 dp_2}{(2\pi)^3} \int \frac{p_1^2 dp_1}{(2\pi)^3} \delta\left(\frac{p_1^2}{2M} + \frac{p_2^2}{2M} + \frac{|\mathbf{p}_1 + \mathbf{p}_2|^2}{2M_F} - \Delta_{J_F}\right) \dots, \quad (5)$$

with M and M_F being the nucleon and residual nucleus masses and Δ_{J_F} the liberated energy.

We assume the hyperon to stay in the $j_{\Lambda} = 1s_{1/2}$ single-particle state, and the initial hypernuclear state is built as $|J_I\rangle \equiv |(j_{\Lambda} J_C) J_I\rangle$, where $|J_C\rangle$ is the $A-1$ Z ground state. Thus, the intrinsic asymmetry parameter is defined as

$$a_{\Lambda} = \begin{cases} A_V & \text{for } J_I = J_C + 1/2, \\ -\frac{J_I + 1}{J_I} A_V & \text{for } J_I = J_C - 1/2. \end{cases} \quad (6)$$

Rewriting the transition amplitude in (4) in the total spin (S, M_S) and isospin (T, M_T) basis we get

$$\sigma(J_I M_I) = \int d\Omega_{p_1} \int dF \sum_{S M_S M_F} \left| \sum_T (-)^T \langle \mathbf{p} P S M_S T J_F M_F | V | J_I M_I \rangle_{\mathcal{A}} \right|^2, \quad (7)$$

where $\mathbf{p} = \frac{1}{2}(\mathbf{p}_2 - \mathbf{p}_1)$, $\mathbf{P} = \mathbf{p}_1 + \mathbf{p}_2$ and

$$\begin{aligned} &\langle \mathbf{p} P S M_S T J_F M_F | V | J_I M_I \rangle_{\mathcal{A}} \\ &= \frac{1}{\sqrt{2}} [\langle \mathbf{p} P S M_S T J_F M_F | V | J_I M_I \rangle - (-)^{S+T} \langle -\mathbf{p} P S M_S T J_F M_F | V | J_I M_I \rangle]. \end{aligned} \quad (8)$$

For the integration in (7) we take the z -axis as indicated in Fig. 1 and choose p_2 , θ_{p_1} and ϕ_{p_1} as independent variables. Thus, expanding the final state in terms of relative and center-of-mass partial waves of the emitted nucleons [17], after integration on ϕ_{p_1} and recoupling of angular momentum, we get

$$\begin{aligned} \sigma(J_I M_I) &= \frac{1}{2}(4\pi)^4 \int d\cos\theta_{p_1} \int dF \sum_{STT'} (-)^{T+T'} \\ &\times \sum_{lL\lambda J} \sum_{l'L'\lambda'J'} i^{-l'-L'-l-L} (-)^{\lambda+S+J+J'+J_I+J_F} \hat{l}' \hat{L}' \hat{\lambda}' \hat{J}' \hat{J} \\ &\times \sum_{kK\kappa} \hat{\kappa} \hat{J}_I (J_I M_I \kappa 0 | J_I M_I) (l 0 l' 0 | k 0) (L 0 L' 0 | K 0) [Y_k(\theta_p, \pi) \otimes Y_K(\theta_p, 0)]_{\kappa 0} \\ &\times \left\{ \begin{matrix} J_I & \kappa & J_I \\ J & J_F & J' \end{matrix} \right\} \left\{ \begin{matrix} \kappa & J' & J \\ S & \lambda & \lambda' \end{matrix} \right\} \left\{ \begin{matrix} l & l' & k \\ L & L' & K \\ \lambda & \lambda' & \kappa \end{matrix} \right\} \\ &\times (p l P L \lambda S J T J_F; J_I | V | J_I) (p l' P L' \lambda' S J' T' J_F; J_I | V | J_I)^* \end{aligned} \tag{9}$$

where $(p l P L \lambda S J T J_F; J_I | V | J_I) = \frac{1}{\sqrt{2}} [1 - (-)^{l+S+T}] \langle p l P L \lambda S J T J_F; J_I | V | J_I \rangle$.

From [17] we have

$$\begin{aligned} &\langle p l P L \lambda S J T J_F; J_I | V | J_I \rangle \\ &= (-)^{J_F+J-J_I} \hat{J}_I^{-1} \sum_{j_p} \langle J_I || (a_{j_\Lambda}^\dagger a_{j_p}^\dagger)_J || J_F \rangle \mathcal{M}(p l P L \lambda S J T; j_\Lambda j_p), \end{aligned} \tag{10}$$

where

$$\mathcal{M}(p l P L \lambda S J T; j_\Lambda j_p) = \frac{1}{\sqrt{2}} [1 - (-)^{l+S+T}] (p l P L \lambda S J T | V | j_\Lambda j_p J). \tag{11}$$

The two-particle spectroscopic amplitudes are cast as [17]

$$\langle J_I || (a_{j_\Lambda}^\dagger a_{j_p}^\dagger)_J || J_F \rangle = (-)^{J+J_I+J_F} \hat{J}_I \left\{ \begin{matrix} J_C & J_I & j_\Lambda \\ J & j_p & J_F \end{matrix} \right\} \langle J_C || a_{j_p}^\dagger || J_F \rangle. \tag{12}$$

To continue, we will adopt the extreme shell model and restrict our attention to cases where the single-proton states are completely filled in $|J_C\rangle$. This is so, for the cores of both $^5_\Lambda\text{He}$ ($J_I = 1/2, J_C = 0$), and $^{12}_\Lambda\text{C}$ ($J_I = 1, J_C = 3/2$) [36].

In such cases, the final nuclear states take the form $|J_F\rangle \equiv |(j_p^{-1} J_C) J_F\rangle$. This leads to

$$\langle J_C || a_{j_p}^\dagger || J_F \rangle = (-)^{J_F+J_C+j_p} \hat{J}_F. \tag{13}$$

Putting all this together and performing the summation on J_F , we obtain

$$\begin{aligned} \sigma_\kappa(J_I) &= \frac{4}{\pi} \hat{J}_I^3 \hat{\kappa}^{-1} \sum_{j_p} \int d\cos\theta_{p_1} \int dF_{j_p} \sum_{STT'} (-)^{T+T'} \\ &\times \sum_{lL\lambda J} \sum_{l'L'\lambda'J'} i^{-l'-L'-l-L} (-)^{\lambda+S+J_I+J_C-j_p+\kappa} \hat{l}' \hat{L}' \hat{\lambda}' \hat{J}' \hat{J}^2 \hat{J}^2 \\ &\times \sum_{kK} (l 0 l' 0 | k 0) (L 0 L' 0 | K 0) [Y_k(\theta_p, \pi) \otimes Y_K(\theta_p, 0)]_{\kappa 0} \\ &\times \left\{ \begin{matrix} j_\Lambda & J_I & J_C \\ J_I & j_\Lambda & \kappa \end{matrix} \right\} \left\{ \begin{matrix} \kappa & j_\Lambda & j_\Lambda \\ j_p & J & J' \end{matrix} \right\} \left\{ \begin{matrix} \kappa & J' & J \\ S & \lambda & \lambda' \end{matrix} \right\} \left\{ \begin{matrix} l & l' & k \\ L & L' & K \\ \lambda & \lambda' & \kappa \end{matrix} \right\} \\ &\times \mathcal{M}(p l P L \lambda S J T; j_\Lambda j_p) \mathcal{M}^*(p l' P L' \lambda' S J' T'; j_\Lambda j_p), \end{aligned} \tag{14}$$

where

$$\int dF_{j_p} \dots = \int p_2^2 dp_2 \int p_1^2 dp_1 \delta \left(\frac{p_1^2}{2M} + \frac{p_2^2}{2M} + \frac{|\mathbf{p}_1 + \mathbf{p}_2|^2}{2M_F} - \Delta_{j_p} \right) \dots \quad (15)$$

The intrinsic asymmetry parameter can finally be written as [31]

$$a_\Lambda = \frac{\omega_1}{\omega_0}, \quad (16)$$

where

$$\begin{aligned} \omega_\kappa &= (-)^\kappa \frac{8}{\sqrt{2\pi}} \hat{\kappa}^{-1} \sum_{j_p} \int d \cos \theta_{p_1} \int dF_{j_p} Y_{\kappa 0}(\theta_{p_1}, 0) \\ &\times \sum_{TT'} (-)^{T+T'} \sum_{LS} \sum_{l\lambda} \sum_{l'\lambda'} i^{l-l'} (-)^{\lambda+\lambda'+S+L+j_p+\frac{1}{2}} \\ &\times \hat{l}' \hat{\lambda}' \hat{j}'^2 (10l'0|\kappa 0) \\ &\times \left\{ \begin{matrix} \kappa & 1/2 & 1/2 \\ j_p & J & J' \end{matrix} \right\} \left\{ \begin{matrix} \kappa & J' & J \\ S & \lambda & \lambda' \end{matrix} \right\} \left\{ \begin{matrix} l' & l & \kappa \\ \lambda & \lambda' & L \end{matrix} \right\} \\ &\times \mathcal{M}(p l P L \lambda S J T; j_\Lambda j_p) \mathcal{M}^*(p l' P L \lambda' S J' T'; j_\Lambda j_p), \quad (17) \end{aligned}$$

with $L = 0$ for the $1s_{1/2}$ state, and $L = 0$ and 1 for the $1p_{3/2}$ state. It has been verified that $\omega_0 = \Gamma_p$, in such way that the new information is carried by ω_1 .

The presence of the Clebsch-Gordan coefficient in Eq. (17), for $\kappa = 1$, ensures that l and l' have opposite parities. Since the initial state in the two matrix elements has a definite parity, this implies that all contributions to ω_1 come from interference terms between the parity-conserving and the parity-violating parts of the transition potential. Furthermore, the antisymmetrization factor in Eq. (11) shows that the two final states have $T \neq T'$. These are general properties of the asymmetry parameter.

We have used these general expressions for the numerical computation of a_Λ within several OME models. The values of ω_0 and ω_1 are shown in Tables I and II, for ${}^5_\Lambda\text{He}$ and ${}^{12}_\Lambda\text{C}$ respectively, in units of the free Λ decay constant, $\Gamma_\Lambda^{(0)} = 2.50 \times 10^{-6}$ eV. In the case of ${}^5_\Lambda\text{He}$, we have also included, between parentheses, the values obtained with the approximate formula usually adopted in the literature [34], which is strictly valid only for s -shell hypernuclei:

$$A_V({}^5_\Lambda\text{He}) \approx \frac{2\Re \left[\sqrt{3} a e^* - b(c^* - \sqrt{2} d^*) + \sqrt{3} f(\sqrt{2} c^* + d^*) \right]}{|a|^2 + |b|^2 + 3(|c|^2 + |d|^2 + |e|^2 + |f|^2)}, \quad (18)$$

where

$$\begin{aligned} a &= \langle np, {}^1S_0 | V | \Lambda p, {}^1S_0 \rangle, & b &= i \langle np, {}^3P_0 | V | \Lambda p, {}^1S_0 \rangle, & c &= \langle np, {}^3S_1 | V | \Lambda p, {}^3S_1 \rangle, \\ d &= -\langle np, {}^3D_1 | V | \Lambda p, {}^3S_1 \rangle, & e &= i \langle np, {}^1P_1 | V | \Lambda p, {}^3S_1 \rangle, & f &= -i \langle np, {}^3P_1 | V | \Lambda p, {}^3S_1 \rangle. \end{aligned} \quad (19)$$

We find values ranging from -0.62 to -0.29 , in qualitative agreement with other theoretical estimates [14–16, 18, 19, 26, 32–35] but in contradiction with some recent experimental determinations [5–9].

We remark here that the approximate formula ignores the fact that the final state of nonmesonic decay is a three-body one and does not include the full contribution of the transitions coming from proton states beyond the s -shell, being therefore of only limited validity, and should not be used for p -shell hypernuclei such as ${}^{12}_\Lambda\text{C}$, or, even worse, for heavier ones. This being said, comparison of the corresponding values for a_Λ in Table I shows that the formula works well within its range of validity.

Also, from Table II we observe that the p -shell contribu-

tions to ω_0 and ω_1 for ${}^{12}_\Lambda\text{C}$ are by no means negligible. However, they are also in approximately the same ratio, so that the effect on a_Λ is much smaller.

In order to obtain simple and manageable analytical expressions for the asymmetry parameter, we adopt now the simplest OME model, which takes into account only the contribution of one pion exchange. As can be seen from our numerical results, the main characteristics of the nonmesonic hypernuclear weak decay asymmetry are well described within this OPEM. Then, as a starting point, this model will be enough to obtain such analytical formula[37].

After a straightforward analytical evaluation of the summation terms indicated in (17), together with the calculation of the nuclear matrix elements indicated in Eqs. (11)-(13) within

TABLE I: Results for the asymmetry parameter, a_Λ , for nonmesonic decay of ${}^5_\Lambda\text{He}$.

Model/Calculations	$\omega_0(1s_{1/2})$	$\omega_1(1s_{1/2})$	a_Λ
π	0.6492	-0.2913	-0.4487 (-0.4456)
(π, η, K)	0.3920	-0.2412	-0.6153 (-0.6384)
$\pi + \rho$	0.5937	-0.1776	-0.2991 (-0.3155)
$(\pi, \eta, K) + (\rho, \omega, K^*)$	0.5526	-0.2974	-0.5382 (-0.5388)
Experiment	KEK-PS E278 [9]		0.24 ± 0.22
$a_\Lambda = A_V({}^5_\Lambda\text{He})$	KEK-PS E462 [7] (preliminary)		$0.11 \pm 0.08 \pm 0.04$

TABLE II: Results for the asymmetry parameter, a_Λ , for nonmesonic decay of ${}^{12}_\Lambda\text{C}$.

Model/Calculations	$\omega_0(1s_{1/2})$	$\omega_0(1p_{3/2})$	$\omega_1(1s_{1/2})$	$\omega_1(1p_{3/2})$	a_Λ
π	0.5206	0.5954	-0.2400	-0.2596	-0.4477
(π, η, K)	0.3336	0.3817	-0.2057	-0.2217	-0.5975
$\pi + \rho$	0.4922	0.5514	-0.1461	-0.1596	-0.2930
$(\pi, \eta, K) + (\rho, \omega, K^*)$	0.4619	0.5083	-0.2546	-0.2599	-0.5303
Experiment	KEK-PS E160 [8]				-0.9 ± 0.3
$a_\Lambda = -2A_V({}^{12}_\Lambda\text{C})$	KEK-PS E508 [7] (preliminary)				-0.44 ± 0.32

the strict OPEM, we can write

$$\begin{aligned}\omega_0 &= \frac{24}{\pi} \int d\cos\theta_{p_1} \int dF_{j_p} \left[3(S_\pi^0)^2 + 18(T_\pi^{20})^2 + (P_\pi^{10})^2 \right], \\ \omega_1 &= \frac{32\sqrt{3}}{\sqrt{\pi}} \int d\cos\theta_{p_1} \int dF_{j_p} Y_{10}(\theta_p, 0) (S_\pi^0 - 2T_\pi^{20}) P_\pi^{10},\end{aligned}\quad (20)$$

for ${}^5_\Lambda\text{He}$, and

$$\begin{aligned}\omega_0 &= \frac{24}{\pi} \int d\cos\theta_{p_1} \int dF_{j_p} \left\{ (1+R) [3(S_\pi^0)^2 + 18(T_\pi^{20})^2 + (P_\pi^{10})^2] \right. \\ &\quad \left. + 7(S_\pi^1)^2 + \frac{1}{5}(T_\pi^{11})^2 + \frac{9}{5}(T_\pi^{31})^2 + \frac{7}{3}(P_\pi^{21})^2 \right\}, \\ \omega_1 &= \frac{32\sqrt{3}}{\sqrt{\pi}} \int d\cos\theta_{p_1} \int dF_{j_p} Y_{10}(\theta_p, 0) \left[(1+R) (S_\pi^0 - 2T_\pi^{20}) P_\pi^{10} \right. \\ &\quad \left. + \left(S_\pi^1 + \frac{1}{5}T_\pi^{11} - \frac{9}{5}T_\pi^{31} \right) P_\pi^{21} \right],\end{aligned}\quad (21)$$

for ${}^{12}_\Lambda\text{C}$. Here S_π^l , T_π^l and P_π^l , and $R = (bP)^2/3$ are, respectively, the nuclear moments and the ratio between the $1p_{3/2}$ and $1s_{1/2}$ contributions defined in Section IV C from Ref. [17]. These formulas clearly show that the s -part of ${}^{12}_\Lambda\text{C}$ results agrees with the ${}^5_\Lambda\text{He}$ ones. This has also been tested numerically, as can be seen from Tables I and II. There we observe that this conclusion is also valid within the complete OME model. Simultaneously, our numerical computation shows that the contribution of the p -shell of ${}^{12}_\Lambda\text{C}$ has approximately the same magnitude that the s -one, both for ω_0 and ω_1 (this has been extensively discussed in Ref. [17] concerning the decay rate).

On the other side, it is also well known that the main con-

tribution comes from the lower quantum numbers and, as it has been discussed in Ref. [17], the tensorial T_π^{20} term is the dominant one in the hypernuclear weak decay case. Thus, approximating the p -shell contribution of ${}^{12}_\Lambda\text{C}$ by the s -one, and retaining only the terms containing the tensorial moment T_π^{20} in our formula, we arrive to a very simple expression for a_Λ :

$$a_\Lambda \simeq -\frac{4\sqrt{3}\pi}{27} \frac{\int d\cos\theta_{p_1} \int dF_{j_p} Y_{10}(\theta_p, 0) T_\pi^{20} P_\pi^{10}}{\int d\cos\theta_{p_1} \int dF_{j_p} (T_\pi^{20})^2}. \quad (22)$$

This expression is valid both for ${}^5_\Lambda\text{He}$ and ${}^{12}_\Lambda\text{C}$ hypernuclear decays, within the OPEM. Our results presented in Eqs. (20), (21) and (22) suggest that, in order to obtain a stronger theo-

retical dependence of a_Λ on the particular hypernucleus considered (as indicated by experimental data) future improvements of the model are required.

We believe that a different decay mechanism should be introduced for s - and p -shell hypernuclei, beyond the Born approximation. X

Summarizing, we have derived simple formulas for the evaluation of the asymmetry parameter, which exactly include the effects of three-body kinematics in the final states and correctly treat the contribution of transitions originated from proton states beyond the s -shell.

Besides, we have deduced exact analytical expressions for the particular cases of $^5_\Lambda\text{He}$ and $^{12}_\Lambda\text{C}$ written, within the OPEM, in terms of nuclear moments.

The numerical values of a_Λ , calculated within different OME models, variate from -0.62 to -0.29 .

The negative value systematically obtained for a_Λ for the two hypernuclei indicates that it will be hard to get a positive or zero value for it in the $^5_\Lambda\text{He}$ case, at least within strict OME models. The puzzle posed by the experimental results for a_Λ in s - and p -shell hypernuclei remains unexplained.

However, our results suggest that additional changes in the nuclear model and/or the decay mechanism should be introduced for s - and p -shell hypernuclei, in such way that the improvements lead to a stronger theoretical dependence of a_Λ on the hypernucleus considered, which could reverse the asymmetry sign in the case of $^5_\Lambda\text{He}$, as required to reproduce the more recent experimental data on this observable.

-
- [1] H. Outa *et al.*, KEK Report No. KEK-PS E462, 2000.
- [2] A. Feliciello, Nucl. Phys. **A691**, 170c (2001); P. Gionatti, Nucl. Phys. **A691**, 483c (2001).
- [3] R.L. Gill, Nucl. Phys. **A691**, 180c (2001).
- [4] G. Garbarino, A. Parreño, and A. Ramos, Phys. Rev. C **69**, 054603 (2004); *ibid*, Nucl. Phys. **A754**, 137c (2005).
- [5] H. Bhang *et al.*, Nucl. Phys. **A754**, 144c (2005).
- [6] H. Outa *et al.*, Nucl. Phys. **A754**, 157c (2005).
- [7] T. Maruta *et al.*, nucl-ex/0402017; *ibid*, Nucl. Phys. **A754**, 168c (2005).
- [8] S. Ajimura *et al.*, Phys. Lett. **B282**, 293 (1992).
- [9] S. Ajimura *et al.*, Phys. Rev. Lett. **84**, 4052 (2000).
- [10] M.M. Block and Dalitz, Phys. Rev. Lett. **11**, 96 (1963).
- [11] J. B. Adams, Phys. Rev. **156**, 832 (1967) (*Cf.* correction pointed out in Ref. [12].)
- [12] B. H. J. McKellar and B. F. Gibson, Phys. Rev. C **30**, (1984) 322.
- [13] K. Takeuchi, H. Takaki, and H. Bandō, Prog. Theor. Phys. **73**, 841 (1985).
- [14] J. F. Dubach, G. B. Feldman, B. R. Holstein, and L. de la Torre, Ann. Phys. (N.Y.) **249**, 146 (1996).
- [15] A. Parreño, A. Ramos, and C. Bennhold, Phys. Rev. C **56**, 339 (1997) and references therein.
- [16] A. Parreño and A. Ramos, Phys. Rev. C **65**, 015204 (2001).
- [17] C. Barbero, D. Horvat, F. Krmpotić, T. T. S. Kuo, Z. Narančić, and D. Tadić, Phys. Rev. C **66**, 055209 (2002).
- [18] C. Barbero, C. de Conti, A.P. Galeão, and F. Krmpotić, Nucl. Phys. **A726**, 267 (2003).
- [19] C. Barbero, A.P. Galeão, and F. Krmpotić, Phys. Rev. C **72**, 035210 (2005).
- [20] A. Ramos, E. Oset, and L. L. Salcedo, Phys. Rev. C **50**, 2314 (1995).
- [21] A. Ramos, M.J. Vicente-Vacas, and E. Oset, Phys. Rev. C **55**, 735 (1997), Erratum-*ibid* Phys. Rev. C **66**, 039903 (2002).
- [22] A. Parreño, A. Ramos, C. Bennhold, and K. Maltman, Phys. Lett. B **435**, 1 (1998).
- [23] W. M. Alberico and G. Garbarino, Phys. Lett. B **486**, 362 (2000).
- [24] J-H. Jun, Phys. Rev. C **63**, 044012 (2001).
- [25] T. Inoue, M. Oka, T. Motoba, and K. Itonaga, Nucl. Phys. **A633**, 312 (1998).
- [26] K. Sasaki, T. Inoue, and M. Oka, Nucl. Phys. **A669**, 331 (2000), Erratum-*ibid* **A678**, 455 (2000).
- [27] M. Shmatikov, Phys. Lett. B **322**, 311 (1994); *ibid*, Nucl. Phys. **A580**, 538 (1994).
- [28] K. Itonaga, T. Ueda, and T. Motoba, Nucl. Phys. **A577**, 301c; (1994) *ibid*, Nucl. Phys. **A585**, 331c (1995); *ibid*, Nucl. Phys. **A639**, 329c (1998).
- [29] E. Jido, E. Oset, and J.A. Palomar, Nucl. Phys. **A694**, 525 (2001).
- [30] K. Sasaki, M. Izaki, and M. Oka, Phys. Rev. C **71**, 035502 (2005).
- [31] C. Barbero and A. Mariano, Phys. Rev. C **73**, 024309 (2006).
- [32] K. Itonaga, T. Ueda, and T. Motoba, Phys. Rev. C **65**, 034617 (2002); K. Itonaga, T. Motoba, and T. Ueda, *Electrophoto Production of Strangeness on Nuand Nuclei* (Sendai03), K. Maeda, H. Tamura, S.N. Nakamura, and O. Hashimoto eds., World Scientific (2004) pp. 397-402.
- [33] K. Sasaki, T. Inoue, and M. Oka, Nucl. Phys. **A702**, 477 (2002).
- [34] W. M. Alberico and G. Garbarino, Phys. Rep. **369**, 1 (2002).
- [35] W. M. Alberico, G. Garbarino, A. Parreño, and A. Ramos, Phys. Rev. Lett. **94**, 1 (2005).
- [36] The decay of the $^{11}_\Lambda\text{B}$ hypernuclei will be analyzed in a forthcoming paper.
- [37] An analytical expression including the contribution of the complete octets meson will be deduced and analyzed in a forthcoming paper.