

BPS Monopoles & Open Spin Chains

Anastasia Doikou* and Theodora Ioannidou†

*Department of Engineering Sciences, University of Patras, GR-26500 Patras, Greece

E-mail: adoikou@upatras.gr

†Department of Mathematics, Physics and Computational Sciences, Faculty of Engineering,
Aristotle University of Thessaloniki, GR-54124 Thessaloniki, Greece

E-mail: ti3@auth.gr

Abstract

We construct $SU(n+1)$ BPS monopoles with minimal symmetry breaking by solving the full Weyl equation. In this context, we explore and discuss the existence of open spin chain-like part within the Weyl equation. For instance, in the $SU(3)$ case the relevant spin chain is the 2-site spin $1/2$ XXX chain with open boundary conditions. We exploit the existence of such a spin chain part in order to solve the full Weyl equation.

1 Introduction

The integrability of the self-dual Yang-Mills equation was first realized by Ward [2], who demonstrated that the twistor transform of Penrose could be used to provide a correspondence between instantons and certain holomorphic vector bundles over the twistor space CP^3 [3]. Since then many alternative methods (see [4] and References therein) have been proposed in order to find instanton and monopole solutions of the self-dual Yang-Mills equation.

A powerful approach introduced by Nahm [5] is the so-called Atiyah-Driinfeld-Hitchin-Manin-Nahm (ADHMN) construction. The Atiyah-Driinfeld-Hitchin-Manin (ADHM) approach [6], allows the construction of instantons in terms of linear algebras in a vector space, which dimension is related to the instanton number. Since monopoles correspond to infinite action instantons, an adaption of the ADHM construction involving an infinite dimensional vector space might also be possible. Nahm was able to formulate such an adaption in the ADHMN construction. To perform this construction, a nonlinear ordinary differential equation (i.e. the Nahm equation) must be solved and its solutions (i.e. the Nahm data) are used to define the Weyl equation.

The Nahm equations provide a system of non-linear ordinary differential equations for three $n \times n$ anti-hermitian matrices T_i (the so-called Nahm data) functions of the variable s :

$$\frac{dT_i}{ds} = \frac{1}{2} \varepsilon_{ijk} [T_j, T_k] \quad (1)$$

where n is the magnetic charge of the BPS monopole configuration and ε_{ijk} is the totally antisymmetric tensor.

In the ADHMN approach, the construction of $SU(n+1)$ monopole solutions of the Bogomolny equation with topological charge n is translated to the following problem, known as the *inverse Nahm transform* [5]. Given the Nahm data for a n -monopole the one-dimensional Weyl equation

$$\left(\mathbb{I}_{2n} \frac{d}{ds} - \mathbb{I}_n \otimes x_j \sigma_j + iT_j \otimes \sigma_j \right) \mathbf{v}(\mathbf{x}, s) = 0 \quad (2)$$

for the complex $2n$ -vector $\mathbf{v}(\mathbf{x}, s)$, must be solved. \mathbb{I}_n denotes the $n \times n$ identity matrix, $\mathbf{x} = (x_1, x_2, x_3)$ is the position in space at which the monopole fields are to be calculated and $(\sigma_1, \sigma_2, \sigma_3)$ are the familiar 2×2 Pauli matrices. In the minimal symmetry breaking,

the Nahm data T'_i s can be cast as

$$T_i = -\frac{i}{2} f_i \tau_i, \quad i = 1, 2, 3, \quad (3)$$

where τ_i 's form the n -dimensional representation of $su(2)$ and satisfy

$$[\tau_i, \tau_j] = 2i\varepsilon_{ijk}\tau_k. \quad (4)$$

Let us choose an orthonormal basis for these solutions, satisfying

$$\int \hat{\mathbf{v}}^\dagger \hat{\mathbf{v}} ds = \mathbb{I}. \quad (5)$$

Given $\hat{\mathbf{v}}(\mathbf{x}, s)$, the normalized vector computed from (2) satisfying (5), the Higgs field Φ and gauge potential A_k are given by

$$\begin{aligned} \Phi &= -i \int s \hat{\mathbf{v}}^\dagger \hat{\mathbf{v}} ds \\ A_k &= \int \hat{\mathbf{v}}^\dagger \partial_k \hat{\mathbf{v}} ds, \end{aligned} \quad (6)$$

where the integrations are to be performed over the range spanned by the minimum and maximum eigenvalues of the asymptotic form of Φ . Then the corresponding Higgs field and gauge potentials satisfy the self-dual Yang-Mills equations, and they are smooth functions of \mathbf{x} .

Recently in [1], $SU(n+1)$ (for generic values of n) *spherically symmetric* BPS monopoles with minimal symmetry breaking located at $(0, 0, r)$ were obtained by solving the corresponding Weyl equation. However, their azimuthal dependence can not be implemented in a straightforward manner. To do so either the full Weyl equation should be solved or, a suitable similarity transformation should be introduced in order to derive the corresponding solutions in terms of the simple ones of [1]. That way the gauge fields can be recovered, possessing an off-diagonal form as opposed to the diagonal one of the spherical symmetric monopoles.

As will be transparent in the next section, solving the full Weyl equation for the generic case is a rather technically difficult task. However, the crucial observation is the existence of a spin chain-like part within the full Weyl equation. This observation allows us to bypass the

existing technical difficulties. More precisely, we present a new approach for constructing the monopole solutions of the self-dual Yang-Mills equation by connecting the corresponding full Weyl equation with a one-dimensional quantum integrable model, that is the XXX spin chain (also known as, *isotropic Heisenberg model*). To our knowledge this is the first attempt to make such a connection; although, the integrability of the Weyl equation was also discussed in [7].

In fact, it is worth noting that the effective Yang-Mills dynamics in several special limits is described by completely integrable systems related to Heisenberg spin chain and its generalization (see, e.g. [8] and References within). It was shown for instance in [9], that the one loop mixing matrix for anomalous dimensions in $N = 4$ super-symmetric Yang-Mills theory can be identified with the conventional isotropic Heisenberg spin 1/2 chain. Also, it was shown in [10, 11] that the XXX Heisenberg chain of noncompact spin $s = 0$, describes the high energy scattering of hadrons in multi-color Quantum Chromo-Dynamics [10].

2 BPS Monopoles

2.1 $SU(2)$ BPS Monopoles

Recall the full Weyl equation (2) takes the form

$$\left(\frac{d}{ds} + iT_j \otimes \sigma_j - x_j \otimes \sigma_j \right) \mathbf{v}(\mathbf{x}, s) = 0, \quad (7)$$

where $(x_1, x_2, x_3) = (x, y, z)$ and the vector $\mathbf{v}(\mathbf{x}, s)$ is of the form $\mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}$.

The spin zero representation of the $su(2)$ case is trivial (that is, $T_i = 0$). Therefore, the corresponding full Weyl equation (7) for the $SU(2)$ BPS monopole reduces to:

$$\frac{d}{ds} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} - \begin{pmatrix} ru & r\psi \\ r\bar{\psi} & -ru \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} = 0, \quad (8)$$

where we define $u = \cos \theta$, $\psi = \sin \theta e^{-i\phi}$ and its complex conjugate $\bar{\psi}$ and thus,

$$z = ru, \quad x - iy = r\psi, \quad x + iy = r\bar{\psi}. \quad (9)$$

Notice that $u^2 + |\psi|^2 = 1$. Then, equation (8) reduces to the following set of differential equations

$$\begin{aligned} \dot{v}_1 - ru v_1 - r\psi v_2 &= 0 \\ \dot{v}_2 + ru v_2 - r\bar{\psi} v_1 &= 0, \end{aligned} \quad (10)$$

and its solution is given by

$$\begin{aligned} v_1 &= \kappa_1 e^{rs} + \kappa_2 e^{-rs} \\ v_2 &= \frac{\kappa_1}{\psi} (1 - u) e^{rs} - \frac{\kappa_2}{\psi} (1 + u) e^{-rs}. \end{aligned} \quad (11)$$

Next choose a vector $\tilde{\mathbf{v}}$ to be of the form (11) with constants $\tilde{\kappa}_1, \tilde{\kappa}_2$, such that the set $\{\mathbf{v}, \tilde{\mathbf{v}}\}$ defines an orthonormal system. Thus, introducing the scalar product

$$\langle \mathbf{v}, \tilde{\mathbf{v}} \rangle = \int_{-1}^1 \mathbf{v}^\dagger(s) \tilde{\mathbf{v}}(s) ds, \quad (12)$$

the following relations must be satisfied

$$\langle \mathbf{v}, \mathbf{v} \rangle = \langle \tilde{\mathbf{v}}, \tilde{\mathbf{v}} \rangle = 1, \quad \langle \mathbf{v}, \tilde{\mathbf{v}} \rangle = 0. \quad (13)$$

Then, due to the constraints (13), the solutions (11) lead to the following choice of constants

$$\begin{aligned} \kappa_1 &= \sqrt{\frac{r}{4 \sinh 2r}} (1 + u), & \kappa_2 &= \sqrt{\frac{r}{4 \sinh 2r}} (1 - u), \\ \tilde{\kappa}_1 &= \sqrt{\frac{r}{4 \sinh 2r}} \psi, & \tilde{\kappa}_2 &= -\sqrt{\frac{r}{4 \sinh 2r}} \psi. \end{aligned} \quad (14)$$

Therefore, we end up with two orthonormal vectors of the form

$$\begin{aligned} \mathbf{v} &= \sqrt{\frac{r}{4 \sinh 2r}} \begin{pmatrix} (1 + u) e^{rs} + (1 - u) e^{-rs} \\ \bar{\psi} e^{rs} - \bar{\psi} e^{-rs} \end{pmatrix}, \\ \tilde{\mathbf{v}} &= \sqrt{\frac{r}{4 \sinh 2r}} \begin{pmatrix} \psi e^{rs} - \psi e^{-rs} \\ (1 - u) e^{rs} + (1 + u) e^{-rs} \end{pmatrix}. \end{aligned} \quad (15)$$

Any other choice of constants is gauge equivalent to the above ones (due to gauge freedom).

Let $\mathbf{v} = \mathbf{v}_1$ and $\tilde{\mathbf{v}} = \mathbf{v}_2$. Then, the associated Higgs field and gauge potentials are 2×2 matrices with elements

$$\begin{aligned}\Phi_{ij} &= -i \int_{-1}^1 s \mathbf{v}_i^\dagger \mathbf{v}_j ds \\ A_{kij} &= \int_{-1}^1 \mathbf{v}_i^\dagger \partial_k \mathbf{v}_j ds.\end{aligned}\tag{16}$$

2.2 $SU(n+1)$ BPS Monopoles

The n -dimensional representation of $su(2)$ is of the form

$$\tau_1 = \sum_{k=1}^{n-1} C_k \left(e_{kk+1}^{(n)} + e_{k+1k}^{(n)} \right), \quad \tau_2 = i \sum_{k=1}^{n-1} C_k \left(e_{k+1k}^{(n)} - e_{kk+1}^{(n)} \right), \quad \tau_3 = \sum_{k=1}^n a_k e_{kk}^{(n)}\tag{17}$$

where $e_{ij}^{(n)}$ are $n \times n$ matrices defined by: $\left(e_{ij}^{(n)} \right)_{kl} = \delta_{ik} \delta_{jl}$ and

$$a_k = n + 1 - 2k, \quad C_k = \sqrt{k(n-k)}.\tag{18}$$

The Nahm data for the $SU(n+1)$ spherically symmetric monopoles of charge n are given by (3) where $f_i = f = -\frac{1}{s}$. Assume that the vector $\mathbf{v}(\mathbf{x}, s)$ is of the form

$$\mathbf{v}(\mathbf{x}, s) = \sum_{l=1}^n h_l(r, s) \hat{e}_l^{(n)} \otimes \left(g_1(r, s) \hat{e}_1^{(2)} + g_2(r, s) \hat{e}_2^{(2)} \right)\tag{19}$$

where $\hat{e}_k^{(n)}$ is the n -dimensional column vector with one at the position $k \in \mathbb{Z}^+$ and zero elsewhere, i.e. the standard basis of \mathbb{R}^n .

Then, the Weyl equation (2) takes the form

$$\begin{aligned}\left[\frac{d}{ds} + \frac{f}{2} \sum_{k=1}^{n-1} C_k \left(e_{kk+1}^{(n)} + e_{k+1k}^{(n)} \right) \otimes \left(e_{12}^{(2)} + e_{21}^{(2)} \right) - \frac{f}{2} \sum_{k=1}^{n-1} C_k \left(e_{k+1k}^{(n)} - e_{kk+1}^{(n)} \right) \otimes \left(e_{21}^{(2)} - e_{12}^{(2)} \right) \right. \\ \left. + \frac{f}{2} \sum_{k=1}^n a_k e_{kk}^{(n)} \otimes \left(e_{11}^{(2)} - e_{22}^{(2)} \right) - z \mathbb{I} \otimes \left(e_{11}^{(2)} - e_{22}^{(2)} \right) - x \mathbb{I} \otimes \left(e_{12}^{(2)} + e_{21}^{(2)} \right) - iy \mathbb{I} \otimes \left(e_{21}^{(2)} - e_{12}^{(2)} \right) \right] \\ \sum_{l=1}^n h_l \hat{e}_l^{(n)} \otimes \left(g_1 \hat{e}_1^{(2)} + g_2 \hat{e}_2^{(2)} \right) = 0.\end{aligned}\tag{20}$$

To proceed with our computation we exploit the following properties

$$e_{ij}^{(n)} e_{kl}^{(n)} = \delta_{kj} e_{il}^{(n)}, \quad e_{ij}^{(n)} \hat{e}_k^{(n)} = \delta_{jk} \hat{e}_i^{(n)}.\tag{21}$$

With the use of the latter identities, and after setting

$$v_l(r, s) = h_l(r, s) g_1(r, s), \quad w_l(r, s) = h_l(r, s) g_2(r, s), \quad (22)$$

equation (20) is equivalent to the following first-order system of differential equations

$$\begin{aligned} \dot{v}_1 - \left(\frac{1}{2s} a_1 + z \right) v_1 - (x - iy) w_1 &= 0, \\ \dot{v}_{k+1} - \frac{1}{s} C_k w_k - \left(\frac{1}{2s} a_{k+1} + z \right) v_{k+1} - (x - iy) w_{k+1} &= 0, \\ \dot{w}_k - \frac{1}{s} C_k v_{k+1} + \left(\frac{1}{2s} a_k + z \right) w_k - (x + iy) v_k &= 0, \\ \dot{w}_n + \left(\frac{1}{2s} a_n + z \right) w_n - (x + iy) v_n &= 0, \end{aligned} \quad (23)$$

where $k = 1, 2, \dots, n - 1$. Here, \dot{v}_i and \dot{w}_i for $i = 1, \dots, n$ are the total derivatives of the functions $v_i(r, s)$ and $w_i(r, s)$ with respect to the argument s .

Solving the aforementioned system is the first step in reconstructing the solution of the self-dual Yang-Mills equation from Nahm data. Then the problem of recovering the Higgs field and gauge potentials given by (6) is linear. However, solving the system (23) is a rather technically difficult task. In the next section, we show how the existing technical difficulties can be overcome by connecting the corresponding full Weyl equation (2) with the isotropic Heisenberg model.

3 Weyl Equation & Open Spin Chains

The Weyl equation (2) can be identified as a Hamiltonian system. In particular, one can observe that (2) can be described by a Hamiltonian containing some bulk *spin-spin* interaction and a *boundary term*.

For simplicity, let us focus on the $SU(3)$ case. The corresponding results can be extended in all other cases; however, not in a straightforward manner. After implementing the simple Nahm data of the minimal symmetry breaking (3), the corresponding full Weyl equation (2) can be written in the form

$$\left(\frac{d}{ds} - \mathcal{H} \right) \mathbf{v} = 0 \quad (24)$$

where

$$\mathcal{H} = \frac{1}{2s} (\sigma_x \otimes \sigma_x + \sigma_y \otimes \sigma_y + \sigma_z \otimes \sigma_z) + x \mathbb{I} \otimes \sigma_x + y \mathbb{I} \otimes \sigma_y + z \mathbb{I} \otimes \sigma_z \quad (25)$$

which is nothing else but the open XXX (Heisenberg) spin chain. The first term in (25) corresponds to the spin-spin interaction; while the rest describes the boundary interaction. This is a quantum integrable model and the corresponding Hamiltonian is immediately obtained from the so-called *open transfer matrix*. We shall not provide the details of this construction in the present investigation, but we refer the interested reader to the original work discussed in [12].

The boundary part of the XXX Hamiltonian (25) can be explicitly expressed as

$$x\sigma_x + y\sigma_y + z\sigma_z = z \begin{pmatrix} 1 & \tan \theta e^{-i\phi} \\ \tan \theta e^{i\phi} & -1 \end{pmatrix}. \quad (26)$$

It is clear that there is an one to one correspondence between the coordinates and the boundary parameters of the open XXX model.

The key point in solving the generic problem described by (24) and (25) is the following observation: A similarity transformation U (see, e.g. [13]) exist, which diagonalizes the boundary contribution in (25), but leaves the bulk spin-spin interaction invariant. More precisely, let

$$U = \begin{pmatrix} 1 + \cos \theta & \sin \theta e^{-i\phi} \\ (1 - \cos \theta) e^{i\phi} & -\sin \theta \end{pmatrix}. \quad (27)$$

Then it is straightforward to show that, the boundary term is transformed as

$$U \begin{pmatrix} 1 & \tan \theta e^{-i\phi} \\ \tan \theta e^{i\phi} & -1 \end{pmatrix} U^{-1} = \frac{1}{\cos \theta} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (28)$$

while the bulk spin-spin interaction term remains unaffected:

$$U \otimes U (\sigma_x \otimes \sigma_x + \sigma_y \otimes \sigma_y + \sigma_z \otimes \sigma_z) U^{-1} \otimes U^{-1} = \sigma_x \otimes \sigma_x + \sigma_y \otimes \sigma_y + \sigma_z \otimes \sigma_z. \quad (29)$$

When the transformation (27) acts to the full Weyl equation (24) and (25), one gets:

$$U \otimes U \left(\frac{d}{ds} - \mathcal{H} \right) \mathbf{v} = 0 \Rightarrow \left(\frac{d}{ds} - \frac{1}{2s} \sigma_i \otimes \sigma_i - r \mathbb{I} \otimes \sigma_z \right) (U \otimes U) \mathbf{v} = 0 \quad (30)$$

which is nothing else but the Weyl equation with only *diagonal boundary* terms. In [1], solutions of the aforementioned equation have been obtained. Therefore, the inverse similarity transformation can be performed in order to solve explicitly the full *non-diagonal* problem.

Let \mathbf{v}_0 to be the solution of the *diagonal problem* [1]. Then the solutions \mathbf{v} of the full problem can be obtained from \mathbf{v}_0 due to the relation

$$\mathbf{v} = (U^{-1} \otimes U^{-1}) \mathbf{v}_0. \quad (31)$$

Let us also briefly consider the simplest case, i.e. the $SU(2)$ BPS monopole. In this case, the corresponding Hamiltonian is just the boundary term (26). This Hamiltonian emerges from the open transfer matrix as an 1-site open XXX chain. After implementing the similarity transformation (27) one gets:

$$U \left(\frac{d}{ds} - \mathcal{H} \right) \mathbf{v} = 0 \Rightarrow \left(\frac{d}{ds} - r \sigma_z \right) U \mathbf{v} = 0. \quad (32)$$

Therefore, the vector \mathbf{v} in terms of the solution of the diagonal problem is of the form

$$\mathbf{v} = U^{-1} \mathbf{v}_0. \quad (33)$$

In general, for the $SU(n+1)$ case the corresponding Hamiltonian has the form:

$$\mathcal{H} = \frac{1}{2s} \tau_i \otimes \sigma_i + x_i \mathbb{I} \otimes \sigma_i. \quad (34)$$

Assume that there exist a transformation \mathcal{U} such that the “bulk” interaction term is

$$\mathcal{U} \otimes U (\tau_i \otimes \sigma_i) \mathcal{U}^{-1} \otimes U^{-1} = (\tau_i \otimes \sigma_i). \quad (35)$$

Then, the aforementioned procedure can be applied in order to obtain all the solutions of the general $SU(n+1)$ case. The generic solution of the Weyl equation will then be

$$\mathbf{v} = (\mathcal{U}^{-1} \otimes U^{-1}) \mathbf{v}_0, \quad (36)$$

where \mathbf{v}_0 the solution of the *diagonal* case found in [1]. Hence, the main question to be investigated is the derivation of the transformation \mathcal{U} . This is an open problem that merits further investigation.

Finally, one needs to implement the orthonormality conditions to the solutions $\mathbf{v}(\mathbf{x}, s)$. The idea is to pick $n + 1$ distinct solutions and require that they satisfy the orthonormality relations:

$$\langle \mathbf{v}_i, \mathbf{v}_j \rangle = \delta_{ij}. \quad (37)$$

This way one may eventually identify the Higgs field and gauge potentials for the generic situation via (6). In the next section two specific examples, displaying explicitly how the proposed methodology works, are discussed in detail.

4 Explicit Examples

4.1 $SU(2)$ Case

Let us first consider the simplest case within the frame we described above, i.e. the $SU(2)$ gauge. Recall that, the solution of the *diagonal* Weyl equation is of the form

$$\begin{aligned} \mathbf{v}_0 &= \begin{pmatrix} g_1(r, s) \\ g_2(r, s) \end{pmatrix} \\ &= \begin{pmatrix} a(r) e^{rs} \\ b(r) e^{-rs} \end{pmatrix}. \end{aligned} \quad (38)$$

Then, the corresponding solution of the full Weyl equation, that is after performing the inverse similarity transformation (33), is given by

$$\mathbf{v} = -\frac{1}{2 \sin \theta} \begin{pmatrix} -[a(r) e^{rs} + b(r) e^{-rs} e^{-i\phi}] \sin \theta \\ a(r) e^{rs} (\cos \theta - 1) e^{i\phi} + b(r) e^{-rs} (\cos \theta + 1) \end{pmatrix}. \quad (39)$$

Next, chose two vectors \mathbf{v}_i for $i = 1, 2$ to be of the form (39) with constants a_i and b_i , such that the set $\{\mathbf{v}_1, \mathbf{v}_2\}$ defines an orthonormal system. One may readily extract that

$$\begin{aligned} \langle \mathbf{v}_i, \mathbf{v}_j \rangle &= \frac{1}{2 \sin^2 \theta} \int_{-1}^1 [\bar{a}_i a_j (1 - \cos \theta) e^{2rs} + \bar{b}_i b_j (\cos \theta + 1) e^{-2rs}] ds \\ &= \frac{\sinh(2r)}{2r} \left[\frac{\bar{a}_i a_j}{(1 + u)} + \frac{\bar{b}_i b_j}{(1 - u)} \right] \end{aligned} \quad (40)$$

using the variables (9). Then, from the orthonormality conditions (37) the form of the constants a_i and b_i which are now functions of the variables (r, u, ψ) (up to a gauge choice) can be obtained. In fact, (39) becomes (15) by setting

$$\begin{aligned} a_1 &= \sqrt{\frac{r}{\sinh 2r}} \psi, & b_1 &= -\sqrt{\frac{r}{\sinh 2r}} \sqrt{|\psi|^2}, \\ a_2 &= \sqrt{\frac{r}{\sinh 2r}} (1+u), & b_2 &= \sqrt{\frac{r}{\sinh 2r}} \sqrt{\frac{\bar{\psi}}{\psi}} (1-u). \end{aligned} \quad (41)$$

Finally, the associated Higgs field and gauge potentials are given by (16). For example, the elements of the Higgs field are of the form

$$\begin{aligned} \Phi_{ij} &= -\frac{i}{2} \int_{-1}^1 s \left[\frac{\bar{a}_i a_j}{(1+u)} e^{2rs} + \frac{\bar{b}_i b_j}{(1-u)} e^{-2rs} \right] ds \\ &= -\frac{i}{2} \left(\frac{-\sinh 2r + 2r \cosh 2r}{2r^2} \right) \left[\frac{\bar{a}_i a_j}{(1+u)} - \frac{\bar{b}_i b_j}{(1-u)} \right]. \end{aligned} \quad (42)$$

4.2 $SU(3)$ Case

The next non-trivial case, which can be easily treated is the $SU(3)$ gauge. Recall that the solution of the diagonal case has the form

$$\mathbf{v}_0 = \begin{pmatrix} h_1(r, s) \\ h_2(r, s) \end{pmatrix} \otimes \begin{pmatrix} g_1(r, s) \\ g_2(r, s) \end{pmatrix} = \begin{pmatrix} v_1(r, s) \\ w_1(r, s) \\ v_2(r, s) \\ w_2(r, s) \end{pmatrix}, \quad (43)$$

where $v_i = g_1 h_i$ and $w_i = g_2 h_i$ for $i = 1, 2$. The entries u_i and w_i are known explicitly due to [1]:

$$\begin{aligned} v_1 &= a(r) \sqrt{s} e^{rs}, & w_1 &= \frac{W}{s}, \\ v_2 &= \dot{W} + \left(r - \frac{1}{2s} \right) W, & w_2 &= b(r) \sqrt{s} e^{-rs} \end{aligned} \quad (44)$$

where

$$W = c(r) M \left(\frac{1}{2}, 1; 2rs \right) = c(r) \left(\frac{\sinh(rs) - rs e^{-rs}}{\sqrt{rs}} \right). \quad (45)$$

$M(k, m; z)$ is the Whittaker function of first type (see also [1]) and \dot{W} denotes the derivative of the function W with respect to s .

After performing the inverse similarity transformation the solution of the full Weyl equation becomes

$$\mathbf{v} = \frac{1}{4 \sin^2 \theta} \begin{pmatrix} -(h_1 + h_2 e^{-i\phi}) \sin \theta \\ h_1 (\cos \theta - 1) e^{i\phi} + h_2 (\cos \theta + 1) \end{pmatrix} \otimes \begin{pmatrix} -(g_1 + g_2 e^{-i\phi}) \sin \theta \\ g_1 (\cos \theta - 1) e^{i\phi} + g_2 (\cos \theta + 1) \end{pmatrix}. \quad (46)$$

As before, we choose three distinct vectors \mathbf{v}_j for $j = 1, 2, 3$ of the form (46). Then, the associated solutions $v_i^{(j)}, w_i^{(j)}$ are given in terms of the constants a_j, b_j and c_j by (44) and (45). Again, one may readily extract that their norm is equal to

$$\begin{aligned} \langle \mathbf{v}_i, \mathbf{v}_j \rangle &= \frac{1}{4 \sin^2 \theta} \int_0^3 \left[\frac{(\cos \theta - 1)^2}{\sin^2 \theta} v_1^{*(i)} v_1^{(j)} + \frac{(\cos \theta + 1)^2}{\sin^2 \theta} w_2^{*(i)} w_2^{(j)} + v_2^{*(i)} v_2^{(j)} + w_1^{*(i)} w_1^{(j)} \right] ds \\ &= \frac{1}{16} \left[\bar{a}_i a_j \frac{q_0}{r^2 (1+u)^2} + \bar{b}_i b_j \frac{q_1}{r^2 (1-u)^2} + \bar{c}_i c_j \frac{q_0 q_1}{9r |\psi|^2} \right], \end{aligned} \quad (47)$$

where $q_0 = 1 + (-1 + 6r) e^{-6r}$, $q_1 = 1 - (1 + 6r) e^{6r}$ and (r, u, ψ) are the parameters defined in (9). Note that, for convenience the limits of integration have been shifted in this case to 0 and 3.

Requiring the orthonormality conditions (37) one may fix, not uniquely, the constants which are now functions of (r, u, ψ) and determine the vectors \mathbf{v}_j . It is straightforward to show that the constants a_j, b_j , and c_j are of the form:

$$\begin{aligned} a_1 &= \frac{4r}{\sqrt{3} q_0} (u+1) e^{i\alpha}, & b_1 &= \frac{4r}{\sqrt{3} q_1} (1-u) e^{i\beta}, & c_1 &= \frac{4\sqrt{r}}{\sqrt{3} q_0 q_1} \psi e^{i\gamma}, \\ a_2 &= \frac{4r}{\sqrt{3} q_0} (u+1) e^{i\alpha - \frac{2\pi i}{3}}, & b_2 &= \frac{4r}{\sqrt{3} q_1} (1-u) e^{i\beta + \frac{2\pi i}{3}}, & c_2 &= \frac{4\sqrt{r}}{\sqrt{3} q_0 q_1} \psi e^{i\gamma}, \\ a_3 &= \frac{4r}{\sqrt{3} q_0} (u+1) e^{i\alpha - \frac{\pi i}{3}}, & b_3 &= \frac{4r}{\sqrt{3} q_1} (1-u) e^{i\beta + \frac{\pi i}{3}}, & c_3 &= \frac{4\sqrt{r}}{\sqrt{3} q_0 q_1} \psi e^{i\gamma - \pi i}. \end{aligned} \quad (48)$$

where α, β , and γ are real free parameters, which for simplicity may be set equal to zero.

Then, the Higgs field and gauge potentials can be recovered from (6). In particular, the elements of the Higgs field are equal to

$$\Phi_{ij} = -\frac{i}{4|\psi|^2} \int_0^3 (s-2) \left[\frac{(u-1)^2}{|\psi|^2} v_1^{*(i)} v_1^{(j)} + \frac{(u+1)^2}{|\psi|^2} w_2^{*(i)} w_2^{(j)} + v_2^{*(i)} v_2^{(j)} + w_1^{*(i)} w_1^{(j)} \right] ds$$

$$\begin{aligned}
&= -\frac{i}{4} \left[\frac{\bar{a}_i a_j}{(1+u)^2} \left(\frac{-1-2r+(1-4r+6r^2)e^{6r}}{4r^3} \right) - \frac{\bar{b}_i b_j}{(1-u)^2} \left(\frac{-1+2r+(1+4r+6r^2)e^{-6r}}{4r^3} \right) \right. \\
&\quad \left. - \frac{\bar{c}_i c_j}{(1-u^2)} \left(\frac{-36r^2-4+(2-3r)e^{6r}+(2+3r)e^{-6r}}{18r} \right) \right]. \tag{49}
\end{aligned}$$

5 Conclusions

$SU(n+1)$ monopole solutions of the full Weyl equation in the case of minimally symmetry breaking were obtained. This was done by implementing azimuthal dependence to the spherically symmetric solutions found in [1] via a suitable similarity transformation. More precisely, the existence of a spin chain-like (XXX chain) part with a bulk spin-spin interaction and a boundary term within the Weyl equation was exploited. In this context, a similarity transformation exists that turns the diagonal boundary terms to generic non-diagonal ones but, leaves the bulk interaction invariant [13]. This is precisely the transformation one utilizes to implement the azimuthal dependence to the spherical symmetric solutions.

In this paper, a particular case that involves the simplest Nahm data is considered and thus, the isotropic Heisenberg model (XXX chain) is involved. For generic Nahm data the situation becomes more complicated and thus, the anisotropic Heisenberg model (XXZ or XYZ chain) might be involved. In the general situation, such a *boundary similarity transformation* that leaves the bulk part unaffected does not exist; thus, more sophisticated methods need to be employed (see, for example, References [14, 15]). This is an intriguing issue, that we hope to address in future investigations.

References

- [1] A. Doikou and T. Ioannidou, JHEP 1008, (2010) 105.
- [2] R. S. Ward, Phys. Lett. A 61, (1977) 81.

- [3] R. S. Ward and R.O. Wells, *Twistor Geometry and Field Theory*, Cambridge University Press (1990).
- [4] N.S. Manton and P.M. Sutcliffe, *Topological Solitons*, Cambridge Monographs on Mathematical Physics, Cambridge University Press (2004).
- [5] W. Nahm, *The construction of all self-dual multimonopoles by the ADHM method*, in *Monopoles in Quantum Field Theory*, eds N.S. Craigie, P. Goddard and W. Nahm (World Scientific, Singapore, 1982).
- [6] M.F. Atiyah, N.J. Hitchin, V.G. Drinfeld and Yu. I. Manin, *Phys. Lett. A* 65, (1978) 185.
- [7] A. Tongas, D. Tsoubelis and P. Xenitidis, *J. Math. Phys.* 42, (2001) 5762.
- [8] A.V. Belitsky, V.M. Braun, A.S. Gorsky and G.P. Korchemsky, *Int. J. Mod. Phys. A* 19, (2004) 4715.
- [9] J.A. Minahan and K. Zarembo, *JHEP* 0303 (2003) 013.
- [10] L.N. Lipatov, *JETP Lett.* 59, (1994) 596.
- [11] L.D. Faddeev and G.P. Korchemsky, *Phys. Lett. B* 342, (1995) 311.
- [12] E.K. Sklyanin, *J. Phys. A* 21, (1988) 2375.
- [13] D. Arnaudon, J. Avan, N. Crampe, A. Doikou, L. Frappat and E. Ragoucy, *J. Stat. Mech.* 0408, (2004) P005.
- [14] J. Cao, H.-Q. Lin, K.-J. Shi and Y. Wang, *Nucl. Phys. B* 663, (2003) 487.
- [15] A. Doikou, *J. Stat. Mech.* 0605, (2006) P010;
A. Doikou, *Phys. Lett. A* 366, (2007) 556.