

Segment representation of a subclass of co-planar graphs

Mathew C. Francis, Jan Kratochvíl, and Tomáš Vyskočil

Department of Applied Mathematics, Charles University,
Malostranské náměstí 25, 11800 Praha 1, Czech Republic.
{francis,honza,whisky}@kam.mff.cuni.cz

Abstract. A graph is said to be a segment graph if its vertices can be mapped to line segments in the plane such that two vertices have an edge between them if and only if their corresponding line segments intersect. Kratochvíl and Kuběna [2] asked the question of whether the complements of planar graphs are segment graphs. We show here that the complements of all partial 2-trees are segment graphs.

1 Introduction

Given a family of sets \mathcal{F} , a simple, undirected graph $G(V, E)$ is said to be an “intersection graph of sets from \mathcal{F} ” if there exists a function $f : V(G) \rightarrow \mathcal{F}$ such that for $u, v \in V(G)$ such that $u \neq v$, $uv \in E(G) \Leftrightarrow f(u) \cap f(v) \neq \emptyset$. We let $IG(\mathcal{F}) = \{G \mid G \text{ is an intersection graph of sets from } \mathcal{F}\}$. When \mathcal{F} is a collection of geometric objects, $IG(\mathcal{F})$ is said to be a class of “geometric intersection graphs”. Some well-known classes of geometric intersection graphs are:

$\text{INT} = IG(\{\text{all intervals on the real line}\})$ (“Interval graphs”)

$\text{STRING} = IG(\{\text{all simple curves in the plane}\})$ (“String graphs”)

$\text{CONV} = IG(\{\text{all convex arc-connected regions in the plane}\})$

$\text{SEG} = IG(\{\text{all straight line segments in the plane}\})$ (“Segment graphs”)

Clearly, $\text{SEG} \subseteq \text{CONV} \subseteq \text{STRING}$. The last inclusion follows from the fact that string graphs are exactly the intersection graphs of arc-connected regions in the plane. It was shown in [2] that the complement of every planar graph is in CONV . In the same paper, the authors pose the question of whether the complement of every planar graph is in SEG . A positive answer to this question would imply that the MAXCLIQUE problem for segment graphs is NP-complete, thus resolving a long standing open problem raised by Kratochvíl and Nešetřil in 1990 [3]. It is worth noting that the question of whether every planar graph is in SEG , known as Scheinerman’s conjecture, was resolved by Chalopin and Gonçalves [1] who showed that every planar graph is indeed the intersection graph of line segments in the plane.

Partial 2-trees are a subclass of planar graphs that includes series-parallel graphs and outer-planar graphs as proper subclasses. In this paper, we show that the complement of every partial 2-tree is in SEG .

2 Definitions

All the graphs that we consider shall be finite, simple and undirected. We denote the vertex set of a graph G by $V(G)$ and its edge set by $E(G)$. Given a graph G and $X \subseteq V(G)$, we denote the subgraph induced by $V(G) \setminus X$ in G as $G - X$. The complement of a graph G , denoted as \overline{G} , is the graph with $V(\overline{G}) = V(G)$ and $E(\overline{G}) = \{uv \mid u \neq v \text{ and } uv \notin E(G)\}$.

2.1 Partial 2-trees

A *2-tree* is defined as follows:

1. A single edge is a 2-tree.
2. If G is a 2-tree, then the graph G' with $V(G') = V(G) \cup \{v\}$ and $E(G') = E(G) \cup \{vx, vy\}$ where $xy \in E(G)$ is a 2-tree.

A *partial 2-tree* is any spanning subgraph of a 2-tree.

2.2 Segments, rays and compatible segment representations

Let $a, b \in \mathbb{R}^2$ be two points in the plane. Then:

- A *segment* with end-points a and b , denoted as ab , is the set $\{a + \rho(b - a) \mid \rho \in [0, 1]\}$. Any point on a segment that is not one of its end-points is said to be an *interior* point of the segment. For the purposes of this paper, we shall assume that every segment has a non-zero length—i.e., the end-points of a segment may not coincide.
- A *ray* starting at a point a and passing through a point b is the set $\{a + \rho(b - a) \mid \rho \in [0, \infty)\}$. A ray has a single end-point, which is its starting point.

Given l_1 and l_2 , where each could be a segment or a ray, they are said to *cross* each other if $l_1 \cap l_2$ consists exactly of a single point that is not an end-point of either l_1 or l_2 . If $l_1 \cap l_2 = \emptyset$, then they are said to be *disjoint*.

Given a segment graph G , there exists a function $f : V(G) \rightarrow \mathcal{R}$ such that $\forall u, v \in V(G), u \neq v, f(u) \cap f(v) \neq \emptyset \Leftrightarrow uv \in E(G)$ where \mathcal{R} is a collection of segments. We say that \mathcal{R} is a “segment representation” of G .

Definition 1. Let G be a partial 2-tree and let G_T be a 2-tree of which G is a spanning subgraph. Let \mathcal{R} be a segment representation of \overline{G} with segments $\{s_u \mid u \in V(G)\}$. \mathcal{R} is said to be a segment representation of \overline{G} that is “compatible with G_T ” if for each $uv \in E(G_T)$, a ray r_{uv} can be drawn in \mathcal{R} such that the collection of these rays satisfies the following properties:

1. r_{uv} starts from an interior point on one of s_u or s_v and passes through an end-point of the other and meets no other points of either s_u or s_v .
2. r_{uv} crosses every segment other than s_u and s_v , and
3. r_{uv} crosses every ray r_{xy} where $xy \in E(G_T) \setminus \{uv\}$.

The rays r_{uv} where $uv \in E(G_T)$ shall be called “special rays”.

3 The construction

Theorem 1. *If G is a partial 2-tree which is a spanning subgraph of a 2-tree G_T , then \overline{G} has a segment representation that is compatible with G_T .*

Proof. We shall prove this by induction on $|V(G)|$. There is nothing to prove for $|V(G)| < 2$. If $|V(G)| = 2$, then clearly, \overline{G} has a segment representation compatible with G_T as shown in Figure 1. Consider a partial 2-tree G with $|V(G)| > 2$. By definition of a 2-tree, there exists a degree 2




Fig. 1. Segment representations that are compatible with G_T for \overline{G} when $|V(G)| = 2$.

vertex v in G_T with neighbours x and y such that $xy \in E(G_T)$ and $G_T - \{v\}$ is also a 2-tree. Let G' be $G - \{v\}$. G' is also a partial 2-tree as it is a spanning subgraph of the 2-tree $G_T - \{v\}$. For ease of notation, we shall denote the 2-tree $G_T - \{v\}$ by G'_T . By our induction hypothesis, there is a segment representation for $\overline{G'}$ that is compatible with G'_T . We shall show how we can extend this to a segment representation for \overline{G} that is compatible with G_T , thereby completing the proof.

Let \mathcal{R}' be a segment representation of $\overline{G'}$ which is compatible with G'_T and let s_u denote the segment corresponding to a vertex $u \in V(\overline{G'})$. We shall add a new segment s_v to \mathcal{R}' so that we get the required segment representation \mathcal{R} for \overline{G} .

Since \mathcal{R}' is compatible with G'_T , there is a special ray r_{xy} in \mathcal{R}' which we shall assume without loss of generality starts from an interior point of s_x and passes through an end-point of s_y . Let p be the starting point of the ray r_{xy} on s_x (refer Figure 2). l_1 and l_2 are two parallel rays starting from points p_1 and p_2 on s_x on either side of p and parallel to r_{xy} such that they cross every segment and special ray that r_{xy} crosses. By our definition of crossing, the ray r_{xy} meets every segment and special ray that it crosses at a point that is an end-point of neither of them. Therefore, we can always choose rays l_1 and l_2 distinct from r_{xy} as long as s_x is not a point or parallel to r_{xy} . Note that by our definition of r_{xy} , s_x cannot lie along r_{xy} and s_x can also not be a point which ensures that the two rays l_1 and l_2 distinct from r_{xy} can be obtained. Of all the crossing points of segments and special rays on l_1 , let q_1 be the farthest from p_1 . Let q_2 be the similarly defined point on l_2 . Clearly, any segment or special ray that meets the segment p_1p_2 and crosses q_1q_2 crosses every segment and special ray that r_{xy} crosses.

The segment s_v is placed in the following way in \mathcal{R}' to obtain \mathcal{R} :

Case 1. $yv \in E(G)$. We place s_v as shown in Figure 3. As $yv \notin E(\overline{G})$, the segment s_v is disjoint from the segment s_y . Let us first consider the case when $xv \notin E(G)$. In \overline{G} , v is adjacent to all




Fig. 2. Starting point of construction




Fig. 3. The case when $yv \in E(G)$




Fig. 4. The case when $yv \notin E(G)$

the vertices in $V(G) \setminus \{v, y\}$. This requirement is satisfied, as since s_v meets p_1p_2 and crosses q_1q_2 , it crosses all the segments that cross r_{xy} and meets s_x too. Moreover, s_v also crosses all the special rays in \mathcal{R}' including r_{xy} . We will show that we can now draw two rays r_{vy} and r_{xv} , so that they, together with the special rays in \mathcal{R}' , form the collection of special rays in \mathcal{R} which make it compatible with G_T . The rays r_{vy} and r_{xv} can be drawn as shown in the figure so that they cross s_x and s_y respectively. In addition, both of them cross each other and r_{xy} . Since they also meet p_1p_2 and cross q_1q_2 , each of them crosses all the special rays and segments that cross r_{xy} . Thus \mathcal{R} is a segment representation of \overline{G} that is compatible with G_T . Note that we can draw the segment s_v and the rays r_{vy} and r_{xv} in this way as long as the segments s_x and s_y have non-zero length and s_y does not lie along the ray r_{xy} . But our definitions of segments and special rays ensure that these pathological situations do not occur. Now, if $xv \in E(G)$, then s_v can be slightly shortened at the end p_3 so that it becomes disjoint from s_x without affecting any of the other arguments so that we still obtain the segment representation \mathcal{R} for \overline{G} .

Case 2. $yv \notin E(G)$. We place s_v , r_{xv} and r_{vy} as shown in Figure 4. The rest of the argument is similar to that of Case 1.

We thus have a segment representation of \overline{G} which is compatible with G_T . This completes the proof. ■

Corollary 1. *If G is any partial 2-tree, then \overline{G} is a segment graph.*

Corollary 2. *The complements of series-parallel graphs and outerplanar graphs are segment graphs.*

Proof. Series-parallel graphs and outerplanar graphs are subclasses of partial 2-trees. □

References

1. Jérémie Chalopin and Daniel Gonçalves. Every planar graph is the intersection graph of segments in the plane: extended abstract. In *STOC '09: Proceedings of the 41st annual ACM symposium on Theory of computing*, pages 631–638, 2009.
2. J. Kratochvíl and Aleš Kuběna. On intersection representations of co-planar graphs. *Discrete Mathematics*, 178(1–3):251–255, 1998.
3. J. Kratochvíl and J. Nešetřil. INDEPENDENT SET and CLIQUE problems in intersection-defined classes of graphs. *Comment. Math. Univ. Carolinae*, 31(1):85–93, 1990.