

FREE RELATIONS FOR MATRIX INVARIANTS IN MODULAR CASE

A.A. LOPATIN

ABSTRACT. A classical linear group $G < GL(n)$ acts on d -tuples of $n \times n$ matrices by simultaneous conjugation. Working over an infinite field of characteristic different from two we establish that the ideal of free relations, i.e. relations valid for matrices of any order, between generators for matrix $O(n)$ - and $Sp(n)$ -invariants is zero. We also prove similar result for invariants of mixed representations of quivers.

As a consequence, we complete the description of relations between generators for $O(n)$ -invariants as well as the description of relations for invariants of mixed representations of quivers. We also obtain an independent proof of the result that the ideal of free relations for $GL(n)$ -invariants is zero, which was proved by Donkin in [Math. Proc. Cambridge Philos. Soc. 113 (1993), 23–43].

Keywords: invariant theory, polynomial invariants, classical linear groups, polynomial identities.

2010 MSC: 13A50; 16R30; 16G20.

1. INTRODUCTION

We assume that \mathbb{F} is an infinite field of arbitrary characteristic $p = \text{char } \mathbb{F}$. All vector spaces, algebras and modules are over \mathbb{F} and all algebras are associative with unity unless otherwise stated.

1.1. Matrix invariants. Consider a group G from the list $GL(n)$, $O(n) = \{A \in \mathbb{F}^{n \times n} \mid AA^T = E\}$, $Sp(n) = \{A \in \mathbb{F}^{n \times n} \mid AA^* = E\}$, where we assume that $p \neq 2$ in case G is $O(n)$ and n is even in the case of $Sp(n)$. Here $\mathbb{F}^{n \times n}$ is the space of $n \times n$ matrices over \mathbb{F} and $A^* = -JA^T J$ is the symplectic transpose of A , where $J = \begin{pmatrix} 0 & E \\ -E & 0 \end{pmatrix}$ is the matrix of the skew-symmetric bilinear form. The group G acts on $V = (\mathbb{F}^{n \times n})^{\oplus d}$ by the diagonal conjugation:

$$g \cdot (A_1, \dots, A_d) = (gA_1g^{-1}, \dots, gA_dg^{-1})$$

for $g \in G$ and A_1, \dots, A_d in $\mathbb{F}^{n \times n}$. The coordinate algebra of V is the polynomial ring

$$R = \mathbb{F}[x_{ij}(k) \mid 1 \leq i, j \leq n, 1 \leq k \leq d]$$

in n^2d variables. The ring R is generated by the entries of *generic matrices* $X_k = (x_{ij}(k))_{1 \leq i, j \leq n}$ ($1 \leq k \leq d$). The action of G on V induces the action on R as follows:

$$g \cdot x_{ij}(k) = (i, j)^{\text{th}} \text{ entry of } g^{-1}X_k g.$$

Denote by R^G the algebra of *matrix G -invariants*, where

$$f \in R^G \text{ iff } g \cdot f = f \text{ for all } g \in G.$$

Consider an arbitrary $n \times n$ matrix X . Denote coefficients in the characteristic polynomial of X by $\sigma_t(X)$, i.e.,

$$\det(X + \lambda E) = \sum_{t=0}^n \lambda^{n-t} \sigma_t(X).$$

So, $\sigma_0(X) = 1$, $\sigma_1(X) = \text{tr}(X)$ and $\sigma_n(X) = \det(X)$.

The following definitions were given in [10]. Let \mathcal{M} be the monoid (without unity) freely generated by *letters*

- x_1, \dots, x_d , if $G = GL(n)$;
- $x_1, \dots, x_d, x_1^T, \dots, x_d^T$, otherwise.

Assume that $a = a_1 \cdots a_r$ and b are elements of \mathcal{M} , where a_1, \dots, a_r are letters.

- Introduce an involution T on \mathcal{M} as follows. If $G = GL(n)$, then $a^T = a$. Otherwise, define $b^{TT} = b$ for a letter b and $a^T = a_r^T \cdots a_1^T \in \mathcal{M}$.
- We say that a and b are *cyclic equivalent* and write $a \stackrel{\circ}{\sim} b$ if there exists a cyclic permutation $\pi \in S_r$ such that $a_{\pi(1)} \cdots a_{\pi(r)} = b$. If $a \stackrel{\circ}{\sim} b$ or $a \stackrel{\circ}{\sim} b^T$, then we say that a and b are *equivalent* and write $a \sim b$.

An element from \mathcal{M} is called *primitive* if it is not equal to the power of a shorter monomial.

- Let $\mathcal{N} \subset \mathcal{M}$ be the subset of primitive elements.
- Let \mathcal{N}_σ be a ring with unity of (commutative) polynomials over \mathbb{F} freely generated by “symbolic” elements $\sigma_t(a)$, where $t > 0$ and $a \in \mathcal{N}$ ranges over \sim -equivalence classes.

We will use the following conventions: $\sigma_0(a) = 1$ and $\sigma_1(a) = \text{tr}(a)$, where $a \in \mathcal{N}$. For a letter $b \in \mathcal{M}$ define

$$X_b = \begin{cases} X_k, & \text{if } b = x_k \\ X_k^T, & \text{if } b = x_k^T \text{ and } G = O(n) \\ X_k^*, & \text{if } b = x_k^T \text{ and } G = Sp(n) \end{cases}.$$

Given $a = a_1 \cdots a_r \in \mathcal{M}$, where a_i is a letter, we set $X_a = X_{a_1} \cdots X_{a_r}$. It is known that the algebra of matrix G -invariants $R^G \subset R$ is generated over \mathbb{F} by $\sigma_t(X_a)$, where $1 \leq t \leq n$ and $a \in \mathcal{N}$. These results were established in [15], [12] in characteristic zero case and in [3], [17] in the general case. Note that if in the case of $p = 0$ we drop the restriction that a is primitive, then it is enough to take $\text{tr}(X_a)$ instead of $\sigma_t(X_a)$, $1 \leq t \leq n$, in the description of generators for R^G . Relations between the mentioned generators were established by Razmyslov [13], Procesi [12] in case $p = 0$ and Zubkov [16] in case $G = GL(n)$ and $p > 0$.

Consider a surjective homomorphism

$$\Psi_n : \mathcal{N}_\sigma \rightarrow R^G$$

defined by $\sigma_t(a) \rightarrow \sigma_t(X_a)$, if $t \leq n$, and $\sigma_t(a) \rightarrow 0$ otherwise. Note that for any $n \times n$ matrices A, B over R and $1 \leq t \leq n$ we have $\sigma_t(A^\delta) = \sigma_t(A)$, $(A^\delta)^\delta = A$, and $(AB)^\delta = B^\delta A^\delta$, where δ stands for the transposition or symplectic transposition. Hence, the map Ψ_n is well defined. Its kernel K_n is the ideal of *relations* for R^G . Elements of

$$K_\infty = \bigcap_{i>0} K_i$$

are called *free relations*. In other words, a relation between the above mentioned generators for R^G is called free if it is valid for $n \times n$ generic matrices for an arbitrary

$n > 0$. In characteristic zero case all free relations are zero (for example, see [12]). We generalize this result to the case of arbitrary characteristic different from two:

Theorem 1.1. *If G is $O(n)$ or $Sp(n)$ and $p \neq 2$, then the ideal K_∞ of free relations for R^G is zero.*

This theorem is proven at the end of Section 5. As a consequence, we obtain an independent proof of the result by Donkin [4] that for an arbitrary p there is no free relations for $R^{GL(n)}$ (see Remark 5.5). The following conjecture is discussed in Remark 5.6.

Conjecture 1.2. *If $p = 2$, then the ideal K_∞ of free relations for $R^{Sp(n)}$ is generated by $\sigma_t(a)$ for $a \in \mathcal{N}$ satisfying $a \stackrel{\sim}{\sim} a^T$ and odd $t \in \{1, \dots, n\}$.*

Applying Theorem 1.1 to Theorem 1.1 from [10], which was proved using an approach from [20], we complete the description of relations between generators for $R^{O(n)}$:

Theorem 1.3. *If $G = O(n)$, then the ideal of relations K_n for $R^{O(n)} \simeq \mathcal{N}_\sigma / K_n$ is generated by $\sigma_{t,r}(a, b, c)$, where $t + 2r > n$ ($t, r \geq 0$) and a, b, c are linear combinations of elements from \mathcal{M} .*

The exact definition of $\sigma_{t,r}(a, b, c) \in \mathcal{N}_\sigma$ can be found in Section 3 of [10]. Note that the function $\sigma_{t,r}$ was introduced by Zubkov in [20] and it relates to the *determinant-pfaffian* from [6] in the same way as σ_t relates to the determinant. More details on the different approaches to the definition of $\sigma_{t,r}$ can be found in Section 1.3 of [10]. Let us remark that the definition of $\sigma_{t,r}$ from [10] is slightly different from the original definition from [20] (see Lemma 7.14 of [10] for details).

The description of relations for $R^{O(n)}$ was applied to the special case of $n = 3$ in [8] and [9].

1.2. Mixed representations of quivers. The notion of *supermixed* representations of a quiver was introduced by Zubkov [18] and [19]. It is equivalent to the notion of representations of a *signed* quiver considered by Shmelkin in [14]. *Orthogonal* and *symplectic* representations of *symmetric* quivers studied by Derksen and Weyman in [2] as well as *mixed* representations of quivers are partial cases of supermixed representations. More details can be found in Section 2.1 of [7].

Section 6 is dedicated to the algebra of invariants $I(\mathcal{Q}, \mathbf{n}, 1)$ of mixed representations of a quiver \mathcal{Q} . As example, a special case of $I(\mathcal{Q}, \mathbf{n}, 1)$ is the algebra of invariants of bilinear forms on vector spaces $V = \mathbb{F}^n$ and V^* under the action of $GL(n)$ as base change (see Example 6.3).

Zubkov established generators for $I(\mathcal{Q}, \mathbf{n}, 1)$ in [19] and described relations between generators modulo free relations in [20]. In Lemma 6.1 we show that there are no non-zero free relations for $I(\mathcal{Q}, \mathbf{n}, 1)$ and in Theorem 6.2 we complete the description of relations for $I(\mathcal{Q}, \mathbf{n}, 1)$. Applying Theorem 6.2, in upcoming paper [11] we will explicitly describe a minimal generating set for invariants of bilinear forms in dimension two case.

2. AUXILIARIES

Denote the degree of $a \in \mathcal{M}$ by $\deg a$, the degree of a in a letter b (i.e., the number of appearances of the letters b and b^T in a) by $\deg_b a$. For $t > 0$ we set $\deg \sigma_t(a) = t \deg a$ and define $\deg f$ for a monomial $f \in \mathcal{N}_\sigma$ in the natural way.

Assume that m is a positive integer. Given $i \in \mathbb{Z}$, we write $|i|_m$ for $1 \leq j \leq m$ such that $i \equiv j \pmod{m}$.

Definition 2.1 (of l -subword). Assume that $a = a_1 \cdots a_r$, $b = b_1 \cdots b_s \in \mathcal{M}$, and $l > 0$, where a_i, b_j are letters for all i, j . We say that

- a is an l -subword of b , if $a_i = b_{|l+i-1|_s}$ for all $1 \leq i \leq r$;
- a is an l^T -subword of b , if $a_i = b_{|l-i+1|_s}^T$ for all $1 \leq i \leq r$.

As an example, for $a = x_1 x_2 x_3^T x_4 \in \mathcal{M}$ we have that $x_3^T x_4 x_1$ is a 3-subword of a and $x_1^T x_4^T$ is an 1^T -subword of a .

Remark 2.2. If $a \sim b$ for $a, b \in \mathcal{M}$, then a is an l -subword of b or l^T -subword of b for some $l > 0$.

Lemma 2.3. Let $b, c \in \mathcal{M}$. Then

- a) if $bc = cb$, then there is an $e \in \mathcal{M}$ such that $b = e^i$ and $c = e^j$ for some $i, j > 0$.
- b) if $b = b^T$, then $b = cc^T$ for a $c \in \mathcal{M}$.

Proof. a) The proof is by induction on $\deg b + \deg c > 1$. If $\deg b + \deg c = 2$, then $b = c$ is a letter and the statement is valid.

If $\deg b = \deg c$, then $b = c$ and the required is proven. Otherwise, without loss of generality we can assume that $\deg b > \deg c$. Then $b = cb_1$ for a $b_1 \in \mathcal{M}$. Thus, $b_1 c = cb_1$. Induction hypothesis completes the proof.

b) The proof is by induction on $\deg b > 0$. If b is a letter, then $b \neq b^T$. Otherwise, $b = yb_1$ for a letter y and $b_1 \in \mathcal{M}$. Since $yb_1 = b_1^T y^T$, we have $b_1 = b_2 y^T$ for a $b_2 \in \mathcal{M}_1$. Then $b_2 = b_2^T$ and the induction hypothesis implies $b_2 = cc^T$ for a $c \in \mathcal{M}_1$. Therefore, $b = yc(yc)^T$. \square

Lemma 2.4. Assume that $a \in \mathcal{N}$ with $\deg a = r$ and $1 \leq l \leq r$. Then

- a) a is an l -subword of a if and only if $l = 1$;
- b) if $a \not\stackrel{c}{\sim} a^T$, then a is not an l^T -subword of a ;
- c) $a \stackrel{c}{\sim} a^T$ if and only if there exists a $b \in \mathcal{N}$ satisfying $a \stackrel{c}{\sim} b$ and $b = b^T$;
- d) if $a = a^T$, then a is an l^T -subword of a if and only if $l = r$.

Proof. a) If a is an l -subword of a for $1 < l \leq m$, then there are $a_1, a_2 \in \mathcal{M}$ such that $a = a_1 a_2$ and $a_1 a_2 = a_2 a_1$. Part a) of Lemma 2.3 implies a contradiction.

b) If a is an l^T -subword of a , then $a = a_1 a_2$ for $a_1, a_2 \in \mathcal{M}_1$ satisfying $a_1 a_2 = a_1^T a_2^T$. Thus $a_i = a_i^T$ and, by part b) of Lemma 2.3, $a_i = c_i c_i^T$ for $c_1, c_2 \in \mathcal{M}_1$ ($i = 1, 2$). Therefore, $a = c_1 c_1^T c_2 c_2^T \stackrel{c}{\sim} a^T$; a contradiction.

c) Since $a \stackrel{c}{\sim} a^T$, we have $a = a_1 a_2$ for $a_1, a_2 \in \mathcal{M}_1$ satisfying $a_1 a_2 = a_1^T a_2^T$. As in the proof of part b), we obtain $a = c_1 c_1^T c_2 c_2^T$, where $c_1, c_2 \in \mathcal{M}_1$. Then $b = c_1^T c_2 (c_1^T c_2)^T$ satisfies the required condition.

d) Let $a = a^T$ and a be an l^T -subword of a for $1 \leq l < m$. As in the proof of part b), we obtain $a = c_1 c_1^T c_2 c_2^T$, where $c_1, c_2 \in \mathcal{M}$. Then for $b_i = c_i c_i^T$ ($i = 1, 2$) we have $b_1 b_2 = b_2 b_1$. Part a) of Lemma 2.3 implies that $a = b_1 b_2$ is not primitive; a contradiction. \square

3. DERIVATIONS

In this section we assume that G is $O(n)$ or $Sp(n)$. Given $q > 0$, we set

$$\widehat{R} = R \otimes \mathbb{F}[y_{ij}(k, q) \mid 1 \leq i, j \leq n, 1 \leq k \leq d, q > 0] \text{ and } Y_{k,q} = (y_{ij}(k, q))_{1 \leq i, j \leq n}.$$

Let G act on \widehat{R} by the same way as on R :

$$g \cdot y_{ij}(k, q) = (i, j)^{\text{th}} \text{ entry of } g^{-1} Y_{k,q} g.$$

Define a linear map $\partial_q : \widehat{R} \rightarrow \widehat{R}$ as follows: given an $f \in \widehat{R}$, we have

$$\partial_q(f) = \sum_{k=1}^d \sum_{1 \leq i, j \leq n} \frac{\partial f}{\partial x_{ij}(k)} y_{ij}(k, q),$$

where $\frac{\partial f}{\partial x_{ij}(k)}$ stands for the partial derivation. As an example, if $f = x_{11}(1) y_{22}(2, r)$, then

$$\partial_q(f) = y_{11}(1, q) y_{22}(2, r) \text{ and } \partial_2 \partial_1(f) = 0.$$

For an $n \times n$ matrix $A = (f_{ij})_{1 \leq i, j \leq n}$ over \widehat{R} we set $\partial_q(A) = (\partial_q(f_{ij}))_{1 \leq i, j \leq n}$. Obviously, $\partial_q(X_k) = Y_{k,q}$, $\partial_q(Y_{k,r}) = 0$ and $\partial_q(A^T) = \partial_q(A)^T$ for $q, r > 0$.

Remark 3.1. The linear map ∂_q has the usual properties of the derivation. Namely, if $f, h \in \widehat{R}$ and $m > 0$, then

$$\partial_q(fh) = \partial_q(f)h + f\partial_q(h) \text{ and } \partial_q(f^m) = m f^{m-1} \partial_q(f).$$

Lemma 3.2. For any $n \times n$ matrices A, B over \widehat{R} and $q > 0$ the following properties hold:

- a) $\partial_q(AB) = \partial_q(A)B + A\partial_q(B)$;
- b) $\partial_q(\sigma_t(A)) = \sum_{i=0}^{t-1} (-1)^i \text{tr}(A^i \partial_q(A)) \sigma_{t-i-1}(A)$ for all $1 \leq t \leq n$.

Proof. For short, we write ∂ for ∂_q . Let $A = (f_{ij})_{1 \leq i, j \leq n}$ and $B = (h_{ij})_{1 \leq i, j \leq n}$. Remark 3.1 implies that $(i, j)^{\text{th}}$ entry of $\partial(AB)$ is equal to

$$\sum_{k=1}^n (\partial(f_{ik}) h_{kj} + f_{ik} \partial(h_{kj})).$$

Hence, part a) is proven. Since

$$(1) \quad \sigma_t(A) = \sum_{1 \leq i_1 < \dots < i_t \leq n} \sum_{\tau \in S_t} \text{sgn}(\tau) f_{i_1, i_{\tau(1)}} \cdots f_{i_t, i_{\tau(t)}},$$

Remark 3.1 implies that $\partial(\sigma_t(A))$ is the coefficient of $\lambda^{t-1} \mu$ in the polynomial $\sigma_t(\lambda A + \mu \partial(A))$ in λ, μ . Amitsur's formula from [1] completes the proof of part b). \square

Example 3.3. Applying Lemma 3.2, we obtain the next equalities.

- Let $f = \sigma_2(X_1)$. Then $\partial_q(f) = -\text{tr}(Y_{1q} X_1) + \text{tr}(X_1) \text{tr}(Y_{1q})$.

- Let $f = \text{tr}(X_1)^2 \text{tr}(X_1 X_2)$ and $p = 2$. Then $\partial_1(f) = \text{tr}(X_1)^2 \text{tr}(Y_{11} X_2) + \text{tr}(X_1)^2 \text{tr}(X_1 Y_{21})$ and $\partial_2 \partial_1(f) = \text{tr}(X_1)^2 \text{tr}(Y_{11} Y_{22}) + \text{tr}(X_1)^2 \text{tr}(Y_{12} Y_{21})$. Note that $\partial_1 \partial_1(f) = 0$.

Similarly to $\mathcal{M}, \mathcal{N}, \mathcal{N}_\sigma$, we introduce the following notions.

- For $q \geq 0$ we denote by $\widehat{\mathcal{M}}(q)$ the monoid (without unity) freely generated by letters $x_k, x_k^T, y_{ks}, y_{ks}^T$, where $1 \leq k \leq d$ and $1 \leq s \leq q$, and set $\widehat{\mathcal{M}} = \cup_{q>0} \widehat{\mathcal{M}}(q)$. Note that $\widehat{\mathcal{M}}(0) = \mathcal{M}$.
- Define the involution T and the equivalences \sim and \simeq on $\widehat{\mathcal{M}}$ in the same way as they were defined on \mathcal{M} .
- Let $\widehat{\mathcal{M}}_1 = \widehat{\mathcal{M}} \sqcup \{1\}$ and $\widehat{\mathcal{M}}_{\mathbb{F}}$ be the vector space with the basis $\widehat{\mathcal{M}}$.
- We denote by $\widehat{\mathcal{N}}(q) \subset \widehat{\mathcal{M}}(q)$ and $\widehat{\mathcal{N}} \subset \widehat{\mathcal{M}}$ subsets of primitive elements and define $\widehat{\mathcal{N}}_\sigma(q), \widehat{\mathcal{N}}_\sigma$ similarly to \mathcal{N}_σ .

Define the notion of degree for $\widehat{\mathcal{M}}$ and $\widehat{\mathcal{N}}_\sigma$ in the same way as for \mathcal{M} and \mathcal{N}_σ . For a letter $b \in \widehat{\mathcal{M}}$ define the $n \times n$ matrix X_b as follows:

$$X_b = \begin{cases} Y_{ks}, & \text{if } b = y_{ks} \\ Y_{ks}^T, & \text{if } b = y_{ks}^T \text{ and } G = O(n) \\ Y_{ks}^*, & \text{if } b = y_{ks}^T \text{ and } G = Sp(n) \\ X_b, & \text{if } b \in \mathcal{M} \end{cases}.$$

Given letters $a_1, \dots, a_r \in \widehat{\mathcal{M}}$, we set $X_{a_1 \dots a_r} = X_{a_1} \dots X_{a_r}$. A homomorphism $\widehat{\Psi}_n : \mathcal{N}_\sigma \rightarrow R^G$ is defined by $\sigma_t(a) \rightarrow \sigma_t(X_a)$, if $t \leq n$, and $\sigma_t(a) \rightarrow 0$ otherwise.

Assume that $q > 0$. We define a linear map $\partial_q : \widehat{\mathcal{M}}_{\mathbb{F}} \rightarrow \widehat{\mathcal{M}}_{\mathbb{F}}$ as follows:

- $\partial_q(x_k) = y_{kq}, \partial_q(x_k^T) = y_{kq}^T$, and $\partial_q(y_{ks}) = \partial_q(y_{ks}^T) = 0$ for all $1 \leq k \leq d$ and $s > 0$;
- $\partial_q(a_1 \dots a_r) = \sum_{i=1}^r a_1 \dots a_{i-1} \partial_q(a_i) a_{i+1} \dots a_r$ for letters $a_1, \dots, a_r \in \widehat{\mathcal{M}}$.

Define a linear map $\partial_q : \widehat{\mathcal{N}}_\sigma(q-1) \rightarrow \widehat{\mathcal{N}}_\sigma(q)$ as follows: for $a, a_1, \dots, a_r \in \widehat{\mathcal{M}}(q-1)$ and $t, t_1, \dots, t_r > 0$, we set

- $\partial_q(\alpha) = 0$ for $\alpha \in \mathbb{F}$;
- $\partial_q(\sigma_t(a)) = \sum_{i=0}^{t-1} (-1)^i \text{tr}(a^i \partial_q(a)) \sigma_{t-i-1}(a)$, where we use the convention that

$$\text{tr}(a_1 + \dots + a_r) = \text{tr}(a_1) + \dots + \text{tr}(a_r);$$

- $\partial_q(\sigma_{t_1}(a_1) \dots \sigma_{t_r}(a_r)) =$

$$\sum_{i=1}^r \sigma_{t_1}(a_1) \dots \sigma_{t_{i-1}}(a_{i-1}) \partial_q(\sigma_{t_i}(a_i)) \sigma_{t_{i+1}}(a_{i+1}) \dots \sigma_{t_r}(a_r).$$

Since an element $(a_1 \dots a_r)^i a_1 \dots a_{j-1} \partial_q(a_j) a_{j+1} \dots a_r$ is either primitive or zero, where $a_1, \dots, a_r \in \widehat{\mathcal{M}}(q-1)$ are letters, the following remark implies that the map $\partial_q : \widehat{\mathcal{N}}_\sigma(q-1) \rightarrow \widehat{\mathcal{N}}_\sigma(q)$ is well defined.

Remark 3.4. For $a, b \in \widehat{\mathcal{M}}$ and $q > 0$ we have

- $\partial_q(a^T) = \partial_q(a)^T$;
- if $a \sim b$, then $\text{tr}(a^m \partial_q(a)) = \text{tr}(b^m \partial_q(b))$ for all $m > 0$.

Note that by abuse of notation we denote three different linear maps by one and the same symbol ∂_q .

Lemma 3.5. *For $q > 0$ the following diagram is commutative:*

$$\begin{array}{ccc} \widehat{\mathcal{N}}_\sigma(q-1) & \xrightarrow{\widehat{\Psi}_n} & \widehat{R}^G \\ \partial_q \downarrow & & \downarrow \partial_q \\ \widehat{\mathcal{N}}_\sigma(q) & \xrightarrow{\widehat{\Psi}_n} & \widehat{R}^G \end{array}$$

In particular, if $f \in \widehat{\mathcal{N}}_\sigma(q-1)$ is a free relation (i.e., $\widehat{\Psi}_n(f) = 0$ for all $n > 0$), then $\partial_q(f) \in \widehat{\mathcal{N}}_\sigma$ is also a free relation.

Proof. Obviously, the statement of the lemma holds for $\text{tr}(X_a)$, where $a \in \widehat{\mathcal{M}}$ is a letter. Lemma 3.2 and the definition of $\partial_q : \widehat{\mathcal{N}}_\sigma(q-1) \rightarrow \widehat{\mathcal{N}}_\sigma(q)$ complete the proof. \square

In the proof of the next lemma we use statements from Section 2, which obviously hold for elements from $\widehat{\mathcal{M}}$.

Lemma 3.6. *Assume that $q, s > 0$ and $a \in \widehat{\mathcal{N}}(q-1)$ satisfies*

- a) $\deg_{x_k}(a) \neq 0$ for some $1 \leq k \leq d$;
- b) $a \not\stackrel{c}{=} a^T$.

Let monomials

$$f_i = \prod_{j=1}^{r_i} \sigma_{t_{ij}}^{m_{ij}}(a) \in \widehat{\mathcal{N}}_\sigma,$$

be pairwise different, where $r_i > 0$, $t_{i1} > \dots > t_{ir_i} \geq 1$ and $m_{i1}, \dots, m_{ir_i} > 0$ are not divided by p ($1 \leq i \leq s$). Then $\partial_q(f_1), \dots, \partial_q(f_s)$ are linear independent over \mathbb{F} .

Proof. Let $\alpha_1 \partial_q(f_1) + \dots + \alpha_s \partial_q(f_s) = 0$, where $\alpha_i \in \mathbb{F}$, be a non-trivial linear combination. Then without loss of generality we can assume that $\alpha_i \neq 0$ for all i . Moreover, without loss of generality we can assume that

$$t_{11} = \max_{i,j} \{t_{ij}\}.$$

For short, we denote $t = t_{11}$. We have

$$\partial_q(a) = b_1 + \dots + b_r$$

for pairwise different $b_1, \dots, b_r \in \widehat{\mathcal{N}}$ and $r > 0$. By the definition of ∂_q ,

$$\partial_q(f_i) = \sum_{w \in \Omega_i} \beta_{i,w} f_{i,w},$$

where Ω_i is equal to the set of pairs

$$\{(u, v, k) \mid 1 \leq u \leq r_i, 0 \leq v < t_{iu}, 1 \leq k \leq r\},$$

$\beta_{i,(u,v,k)} = (-1)^v m_{iu}$ is non-zero, and

$$f_{i,(u,v,k)} = \sigma_{t_{iu}}(a)^{m_{iu}-1} \text{tr}(a^v b_k) \sigma_{t_{iu}-v-1}(a) \prod_{1 \leq j \leq r_i, j \neq u} \sigma_{t_{ij}}^{m_{ij}}(a).$$

We claim that for $i_0 = 1$ and $w_0 = (1, t-1, 1)$ the following statement holds:

If $f_{i,w} = f_{i_0,w_0}$, then $i = i_0$ and $w = w_0$.

Assume that i and $w = (u, v, k)$ satisfy $f_{i,w} = f_{i_0,w_0}$. There exists a unique $1 \leq k' \leq d$ such that $\deg_z(b_1) = 1$ for $z = y_{k',q}$. The only multiplier of f_{i_0,w_0} that contains z or z^T is $\text{tr}(a^{t-1}b_1)$ and the only multiplier of $f_{i,w}$ that can contain z or z^T is $\text{tr}(a^v b_k)$. Therefore, $a^{t-1}b_1 \sim a^v b_k$. The last equivalence implies that $v = t - 1$ and one of the following cases holds:

1. $a^{t-1}b_1 \stackrel{\sim}{\sim} a^{t-1}b_k$;
2. $a^{t-1}b_1 \stackrel{\sim}{\sim} (a^{t-1}b_k)^T$.

Note that the result of the substitutions $z \rightarrow x_{k'}$, $z^T \rightarrow x_{k'}^T$ in b_1 as well as in b_k is a . Thus, making these substitutions in the above equivalences, we obtain that a is the l -subword of a in case 1 and a is the l^T -subword of a in case 2, where $1 \leq l \leq \deg a$. Part b) of Lemma 2.4 implies a contradiction in case 2. Part a) of Lemma 2.4 implies that $l = 1$ in case 1. Thus, $a^{t-1}b_1 = a^{t-1}b_k$ and $k = 1$. Since $v < t_{iu} \leq t$, we obtain $t_{iu} = t$. The inequalities $t_{i,1} > \dots > t_{i,r_i}$ imply $u = 1$. Therefore, $w = w_0$.

It is not difficult to see that in the quotient field of $\widehat{\mathcal{N}}_\sigma$ we have

$$f_{i,w} = \frac{f_i}{\sigma_i(a)} \text{tr}(a^{t-1}b_1) \text{ and } f_{i_0,w_0} = \frac{f_1}{\sigma_t(a)} \text{tr}(a^{t-1}b_1).$$

Thus $i = 1$ and the claim is proven. Obviously, the claim implies a contradiction to the fact that $\alpha_{i_0} \neq 0$. The lemma is proven. \square

Lemma 3.7. *Assume that in the formulation of Lemma 3.6 we have $a \stackrel{\sim}{\sim} a^T$ instead of condition b). Then*

- if $p \neq 2$, then $\partial_q(f_1), \dots, \partial_q(f_s)$ are linear independent over \mathbb{F} ;
- if $p = 2$, then $\partial_q(f_i) = 0$ for all i .

Proof. We use notations from the formulation of Lemma 3.6. Without loss of generality we can assume that

$$t_{11} = \max_{i,j} \{t_{ij}\}.$$

For short, we denote $t = t_{11}$. By part b) of Lemma 2.3 and part c) of Lemma 2.4, without loss of generality we can assume that $a = cc^T$ for some $c \in \widehat{\mathcal{M}}$. We have $\partial_q(c) = b_1 + \dots + b_r$, where $b_1, \dots, b_r \in \widehat{\mathcal{M}}$ are pairwise different and $r > 0$. By Remark 3.4,

$$\partial_q(a) = b_1 c^T + \dots + b_r c^T + c b_1^T + \dots + c b_r^T.$$

Since $\text{tr}(a^v b_k c^T) = \text{tr}(a^v c b_k^T)$ for all $1 \leq k \leq r$ and $v > 0$, we obtain that

$$\partial_q(f_i) = \sum_{w \in \Omega_i} \beta_{i,w} f_{i,w},$$

where Ω_i is equal to the set of pairs

$$\{(u, v, k) \mid 1 \leq u \leq r_i, 0 \leq v < t_{iu}, 1 \leq k \leq r\},$$

$\beta_{i,(u,v,k)} = 2(-1)^v m_{iu}$, and

$$f_{i,(u,v,k)} = \sigma_{t_{iu}}(a)^{m_{iu}-1} \text{tr}(a^v b_k c^T) \sigma_{t_{iu}-v-1}(a) \prod_{1 \leq j \leq r_i, j \neq u} \sigma_{t_{ij}}^{m_{ij}}(a).$$

Therefore, if $p = 2$, then $\partial_q(f_i) = 0$ for all i and the required is proven.

Let $p \neq 2$. We claim that if $f_{i,w} = f_{i_0,w_0}$, then $i = i_0$ and $w = w_0$, where $i_0 = 1$ and $w_0 = (1, t-1, 1)$.

Let i and $w = (u, v, k)$ satisfy $f_{i,w} = f_{i_0,w_0}$. There exists a unique $1 \leq k' \leq d$ such that $\deg_z(b_1) = 1$ for $z = y_{k',q}$. The only multiplier of f_{i_0,w_0} that contains z or z^T is $\text{tr}(a^{t-1}b_1c^T)$ and the only multiplier of $f_{i,w}$ that can contain z or z^T is $\text{tr}(a^v b_k c^T)$. Therefore, $a^{t-1}b_1c^T \sim a^v b_k c^T$. The last equivalence implies that $v = t-1$ and one of the following cases holds:

1. $a^{t-1}b_1c^T \stackrel{c}{\sim} a^{t-1}b_k c^T$;
2. $a^{t-1}b_1c^T \stackrel{c}{\sim} a^{t-1}c b_k^T$.

Note that the result of the substitutions $z \rightarrow x_{k'}$, $z^T \rightarrow x_{k'}^T$ in b_1 as well as in b_k is c . Making these substitutions, we obtain that a is the l -subword of a in both cases, where $1 \leq l \leq \deg a$. Part a) of Lemma 2.4 implies that $l = 1$. Since $\deg_z(a) = 0$, we obtain a contradiction in case 2 and the equality $a^{t-1}b_1 = a^{t-1}b_k$ in case 1.

So, we proved that $k = 1$. The rest of the proof of the claim is the same as in the proof of Lemma 3.6 and the required follows from the claim. \square

4. p -MULTILINEAR FREE RELATIONS

We assume that G is $O(n)$ or $Sp(n)$. Let $f \in \widehat{\mathcal{N}}_\sigma$ be a monomial. If $p > 0$, then we write $f = f^+ f^-$ for

$$(2) \quad f^+ = \sigma_{t_1}^p(a_1) \cdots \sigma_{t_r}^p(a_r) \text{ and } f^- = \sigma_{l_1}^{q_1}(b_1) \cdots \sigma_{l_s}^{q_s}(b_s),$$

where $a_1, \dots, a_r, b_1, \dots, b_s \in \widehat{\mathcal{N}}$, $1 \leq q_1, \dots, q_s < p$, and $\sigma_{l_1}(b_1), \dots, \sigma_{l_s}(b_s)$ are pairwise different elements of $\widehat{\mathcal{N}}_\sigma$. If $p = 0$, then we set $f^+ = 1$ and $f^- = f$.

As example, if $f = \text{tr}^5(x_1)$ and $p = 2$, then $f^+ = \text{tr}^4(x_1)$ and $f^- = \text{tr}(x_1)$.

Definition 4.1. Let $f = \sum_{w \in \Omega} \alpha_w f_w \in \widehat{\mathcal{N}}_\sigma$, where $\alpha_w \in \mathbb{F}$ is non-zero and f_w is a monomial. Then f is called *multilinear* if $\deg_z(f_w) \leq 1$ for every letter $z \in \widehat{\mathcal{M}}$ and $w \in \Omega$.

The element f is called *p -multilinear* if there is a subset $I \subset \{x_k, y_{k,q} \mid 1 \leq k \leq d, q > 0\}$ such that every $w \in \Omega$ satisfies the following conditions:

- $\deg_z(f_w^+) = 0$ and $\deg_z(f_w^-) \leq 1$ for every letter $z \notin I$;
- $\deg_z(f_w^-) = 0$ for every $z \in I$.

In this section we prove that if there is a non-zero free relation, then there exists a non-zero p -multilinear free relation (see Corollary 4.4 below).

For f as in Definition 4.1 we set

$$\deg^+(f) = \max_{w \in \Omega} \{\deg(f_w^+)\} \quad \text{and} \quad \deg^-(f) = \max_{w \in \Omega} \left\{ \sum_{k=1}^d \deg_{x_k}(f_w^-) \right\}.$$

Note that for $q > 0$ and a monomial $f \in \widehat{\mathcal{N}}_\sigma$ we have

$$(3) \quad \partial_q(f) = f^+ \partial_q(f^-).$$

The next remark follows from the definition of ∂_q and the fact that if $b^r = c^s$ for $b, c \in \mathcal{M}$ and $r, s > 0$, then there is an $e \in \mathcal{M}$ such that $b = e^i$ and $c = e^j$ for some $i, j > 0$ (see part a) of Lemma 2.3).

Remark 4.2. Let $q, t, l > 0$, $a, b \in \widehat{\mathcal{N}}(q-1)$, $\partial_q(\sigma_t(a)) = \sum_i \pm f_i$ and $\partial_q(\sigma_l(b)) = \sum_j \pm h_j$ for monomials f_i, h_j . Then there exist i, j with $f_i = h_j$ if and only if $a \sim b$ and $t = l$.

Lemma 4.3. Let $p \neq 2$ and $f \in \mathcal{N}_\sigma$. Then there is a $q \geq 0$ such that $\partial_q \cdots \partial_1(f)$ is a non-zero p -multilinear element of $\widehat{\mathcal{N}}_\sigma$.

Proof. Let $q > 0$. Consider $h = \sum_{w \in \Omega} \alpha_w h_w \in \widehat{\mathcal{N}}_\sigma$ for non-zero elements $\alpha_w \in \mathbb{F}$ and pairwise different monomials $f_w \in \widehat{\mathcal{N}}_\sigma$. Let h be in $\widehat{\mathcal{N}}_\sigma(q-1)$ and

$$(4) \quad \deg_{y_{ks}}(h_w^+) = 0 \text{ and } \deg_{y_{ks}}(h_w^-) \leq 1$$

for all $1 \leq k \leq d$, $s > 0$. Then we claim that one the following possibilities holds:

- h is a p -multilinear;
- $h' = \partial_q(h)$ is non-zero, h' satisfies condition (4) and $\deg^-(h') < \deg^-(h)$.

If $\deg^-(h) = 0$, then h is p -multilinear for $I = \{x_1, \dots, x_d\}$.

We assume that $\deg^-(h) > 0$. Let $\{a_1, \dots, a_s\}$ be a subset of $\widehat{\mathcal{N}}$ such that for every $w \in \Omega$ we have

$$h_w = \prod_{1 \leq i \leq s} h_{w,i},$$

where $h_{w,i}$ is a product of some elements of the set $\{\sigma_t(a_i) \mid t > 0\}$ or $h_{w,i} = 1$. Given $1 \leq i \leq s$, denote by Θ_i the set of $w \in \Omega$ with $\deg_{x_k}(h_{w,i}^-) \neq 0$ for some $1 \leq k \leq d$.

Since $\deg^-(h) > 0$, the set Θ_{i_0} is not empty for some $1 \leq i_0 \leq s$. Using formula (3) and the equality $h_w^- = \prod_{i=1}^s h_{w,i}^-$, we obtain that for every $w \in \Omega$

$$(5) \quad \partial_q(h_w) = h_w^+ \sum h_{w,1}^- \cdots h_{w,i-1}^- \partial_q(h_{w,i}^-) h_{w,i+1}^- \cdots h_{w,s}^-,$$

where the sum ranges over $1 \leq i \leq s$ satisfying $w \in \Theta_i$. Note that if $w \notin \Theta_i$ for all i , then $\partial_q(h_w) = 0$. Applying Lemmas 3.6 and 3.7, we obtain that $\{\partial_q(h_{w,i_0}^-) \mid w \in \Theta_{i_0}\}$ are linear independent over \mathbb{F} . Thus, the definition of $h_{w,i}$ together with Remark 4.2 implies that $h' = \partial_q(h) \neq 0$. It follows from formula (5) that h' satisfies condition (4) and $\deg^-(h') < \deg^-(h)$. Therefore, the claim is proven.

Applying the claim to f , $\partial_1(f)$, $\partial_2(\partial_1(f))$ and so on, we prove the required statement by induction on $\deg^-(f)$. \square

Corollary 4.4. Let G be $O(n)$ or $Sp(n)$ and $p \neq 2$. Assume that $f \in \mathcal{N}_\sigma$ is a non-zero free relation and $d \gg 0$ is large enough. Then there exists a non-zero p -multilinear free relation $h \in \mathcal{N}_\sigma$ with $\deg^+(h) \leq \deg^+(f)$.

Proof. Applying Lemma 4.3, we obtain $q \geq 0$ such that $f' = \partial_q \cdots \partial_1(f)$ is a non-zero p -multilinear element of $\widehat{\mathcal{N}}_\sigma$. By Lemma 3.5, f' is a free relation. Equality (3) implies that $\deg^+(f') \leq \deg^+(f)$.

Let $f' = \sum \alpha_i f_i$ for non-zero $\alpha_i \in \mathbb{F}$ and pairwise different monomials f_i . Since d is large enough, there is an injective map φ from the set of y_{ks} satisfying $\deg_{y_{ks}}(f_i) \neq 0$ for some i ($1 \leq k \leq d$, $s > 0$) to the set of x_j satisfying $\deg_{x_j}(f_i) = 0$ for all

i ($1 \leq j \leq d$). Making substitutions $y_{ks} \rightarrow \varphi(y_{ks})$ and $y_{ks}^T \rightarrow \varphi(y_{ks})^T$ in f' , we obtain the required $h \in \mathcal{N}_\sigma$. \square

5. MULTILINEAR FREE RELATIONS

We assume that G is $O(n)$ or $Sp(n)$. Given an $n \times n$ matrix $A = (f_{ij})_{1 \leq i, j \leq n}$ over R , we denote

$$A^{(p)} = (f_{ij}^p)_{1 \leq i, j \leq n}.$$

Remark 5.1. Let \mathcal{A} be a commutative \mathbb{F} -algebra and $p > 0$. Then for $a_1, \dots, a_r \in \mathcal{A}$ we have $(a_1 + \dots + a_r)^p = a_1^p + \dots + a_r^p$.

Lemma 5.2. For $n \times n$ matrices A and B over R the following properties hold:

- a) $(AB)^{(p)} = A^{(p)}B^{(p)}$;
- b) $\sigma_t(A)^p = \sigma_t(A^{(p)})$ for $1 \leq t \leq n$;
- c) if n is even, then $(A^*)^{(p)} = (A^{(p)})^*$.

Proof. We set $A = (f_{ij})_{1 \leq i, j \leq n}$ and $B = (h_{ij})_{1 \leq i, j \leq n}$. Then $(i, j)^{\text{th}}$ entry of AB is $(\sum_{k=1}^n f_{ik}h_{kj})^p$ and Remark 5.1 completes the proof of part a). Part b) follows from formula (1) and Remark 5.1. Part c) follows from part a). \square

Lemma 5.3. Assume that $p \neq 2$, $f \in \mathcal{N}_\sigma$ is a non-zero p -multilinear free relation and $d \gg 0$ is large enough. Then there exists a non-zero multilinear free relation in \mathcal{N}_σ .

Proof. Without loss of generality we can assume that $p > 0$. Let $f = \sum_{w \in \Omega} \alpha_w f_w \in \mathcal{N}_\sigma$ be not multilinear, where $\alpha_w \in \mathbb{F}$ is non-zero and f_w is a monomial. Note that $f_w^+ = h_w^p$ for some $h_w \in \mathcal{N}_\sigma$. Definition 4.1 implies that there is a set $I \subset \{1, \dots, d\}$ such that for every w the element f_w^+ “depends” only on $\{x_k \mid k \in I\}$ whereas f_w^- “depends” only on $\{x_k \mid k \notin I\}$. Hence $h = \sum_{w \in \Omega} \alpha_w h_w f_w^-$ is a non-zero element of \mathcal{N}_σ satisfying $\deg^+(h) < \deg^+(f)$.

Given $n > 0$, we have $\Psi_n(f) = 0$. By Remark 5.1, $\Psi_n(h_w^p)$ is a polynomial in $x_{ij}^p(k)$, where $k \in I$ and $1 \leq i, j \leq n$. It follows from Lemma 5.2 that the result of substitution $x_{ij}^p(k) \rightarrow x_{ij}(k)$ ($k \in I$, $1 \leq i, j \leq n$) in $\Psi_n(h_w^p)$ is $\Psi_n(h_w)$. Thus, applying the mentioned substitution to $\Psi_n(f) = 0$ we obtain $\Psi_n(h) = 0$. Therefore, h is a free relation.

Applying Corollary 4.4 to h , we obtain a non-zero p -multilinear free relation f' satisfying $\deg^+(f') \leq \deg^+(h)$. Repeating this procedure several times and using the fact that $\deg^+(f)$ decreases at each step by at least one, we finally obtain a non-zero multilinear free relation. \square

Lemma 5.4. There is no a non-zero multilinear free relation in \mathcal{N}_σ for $p \geq 0$.

Proof. We assume that $f = \sum_{w \in \Omega} \alpha_w f_w \in \mathcal{N}_\sigma$ is a non-zero multilinear free relation for non-zero $\alpha_w \in \mathbb{F}$ and pairwise different monomials f_w . Since \mathbb{F} is infinite, without loss of generality we can assume that f is homogeneous with respect to \mathbb{N}^d -grading of \mathcal{N}_σ , i.e., $\deg_{x_1}(f_w) = \dots = \deg_{x_d}(f_w) = 1$ for all w .

We set $n = d$ in case G is the orthogonal group and $n = 2d$ in case G is the symplectic group. Denote by $e_{i,j}$ the $n \times n$ matrix whose $(i, j)^{\text{th}}$ entry is 1 and any other entry is 0. Let $u \in \Omega$ and $f_u = \text{tr}(a_1) \cdots \text{tr}(a_r)$ for some $a_1, \dots, a_r \in \mathcal{N}$.

Given $a_1 = z_1 \cdots z_s$, $a_2 = z_{s+1} \cdots z_l$, and so on, where z_1, \dots, z_l are letters, we set $Z_i = e_{i,i+1}$ for $1 \leq i < s$ and $Z_s = e_{s,1}$. Similarly, we define $Z_i = e_{i,i+1}$ for $s+1 \leq i < l$ and $Z_l = e_{l,s+1}$. Considering a_3, \dots, a_r , we define Z_i for all $l < i \leq d$ as above.

Note that in the symplectic case $e_{ij}^* = e_{j+d,i+d}$ for $1 \leq i, j \leq d$. Hence in both cases the result of substitutions

$$x_{ij}(k) \rightarrow (i, j)^{\text{th}} \text{ entry of } Z_k \quad (1 \leq k \leq d)$$

in $\Psi_n(f_w)$ is zero for $w \neq u$ and one for $w = u$. Since f is a free relation, we have $\Psi_n(f) = 0$. Thus we obtain $\alpha_u = 0$; a contradiction. \square

We now can prove Theorem 1.1:

Proof. Let f be a non-zero free relation. Obviously, without loss of generality we can assume that d is large enough. Then Corollary 4.4 and Lemmas 5.3, 5.4 imply a contradiction. \square

Remark 5.5. In case $G = GL(n)$ we can repeat the proof of Theorem 1.1 without reference to Lemma 3.7, where the restriction $p \neq 2$ is essential. As the result, we obtain that there is no free relations for $R^{GL(n)}$ for an arbitrary p .

Remark 5.6. Let $p = 2$ and $G = Sp(n)$. By straightforward calculations we can see that $\text{tr}(AA^*) = 0$ for every $n \times n$ matrix A over R . By part b) of Lemma 2.3 and part c) of Lemma 2.4, elements $\text{tr}(a) \in \mathcal{N}_\sigma$ with $a \stackrel{\mathcal{L}}{\sim} a^T$ are free relations. On the other hand, it is not difficult to see that $\sigma_2(x_i x_i^T)$ is not a free relation ($1 \leq i \leq d$).

6. INVARIANTS OF MIXED REPRESENTATIONS OF QUIVERS

A *quiver* $\mathcal{Q} = (\mathcal{Q}_0, \mathcal{Q}_1)$ is a finite oriented graph, where \mathcal{Q}_0 (\mathcal{Q}_1 , respectively) stands for the set of vertices (the set of arrows, respectively). For an arrow a , denote by a' its head and by a'' its tail. We say that $a = a_1 \cdots a_r$ is a *path* in \mathcal{Q} (where $a_1, \dots, a_r \in \mathcal{Q}_1$), if $a'_1 = a'_2, \dots, a'_{r-1} = a'_r$. The head of the path a is $a' = a'_1$ and the tail is $a'' = a''_r$. A path a is called *closed* if $a' = a''$.

Given a *dimension vector* $\mathbf{n} = (\mathbf{n}_v \mid v \in \mathcal{Q}_0)$, we consider

- the space $H = \sum_{a \in \mathcal{Q}_1} \mathbb{F}^{\mathbf{n}_{a'} \times \mathbf{n}_{a''}} \simeq \sum_{a \in \mathcal{Q}_1} \text{Hom}(\mathbb{F}^{\mathbf{n}_{a''}}, \mathbb{F}^{\mathbf{n}_{a'}})$;
- the coordinate ring $R = \mathbb{F}[x_{ij}^a \mid a \in \mathcal{Q}_1, 1 \leq i \leq \mathbf{n}_{a'}, 1 \leq j \leq \mathbf{n}_{a''}]$ of H ;
- the $\mathbf{n}_{a'} \times \mathbf{n}_{a''}$ *generic* matrix $X_a = (x_{ij}^a)$ for every $a \in \mathcal{Q}_1$;
- the group $GL(\mathbf{n}) = \sum_{v \in \mathcal{Q}_0} GL(\mathbf{n}_v)$, acting on H as the base change, i.e.,

$$g \cdot (h_a) = (g_{a'} h_a g_{a''}^{-1})$$

for $g = (g_v) \in GL(\mathbf{n})$ and $(h_a) \in H$; this action induces the action of $GL(\mathbf{n})$ on R .

Given a path $a = a_1 \cdots a_r$ with $a_i \in \mathcal{Q}_1$, we write X_a for $X_{a_1} \cdots X_{a_r}$. Donkin [5] proved that the algebra of *invariants* of representations of \mathcal{Q}

$$I(\mathcal{Q}, \mathbf{n}) = R^{GL(\mathbf{n})}$$

is the subalgebra of R generated by $\sigma_t(X_a)$, where a is a closed path in \mathcal{Q} and $1 \leq t \leq \mathbf{n}_{a'}$. Moreover, we can assume that a is *primitive*, i.e., is not equal to the power of a shorter closed path in \mathcal{Q} .

Let $\iota : \mathcal{Q}_0 \rightarrow \mathcal{Q}_0$ be an involution, i.e., ι^2 is the identical map, satisfying $\iota(v) \neq v$ and $\mathbf{n}_{\iota(v)} = \mathbf{n}_v$ for every vertex $v \in \mathcal{Q}_0$. Define

- the group $GL(\mathbf{n}, \mathbf{1}) \subset GL(\mathbf{n})$ by $(g_v) \in GL(\mathbf{n}, \mathbf{1})$ if and only if $g_v g_{\iota(v)}^T = E$ for all v ;
- the double quiver \mathcal{Q}^D by $\mathcal{Q}_0^D = \mathcal{Q}_0$ and $\mathcal{Q}_1^D = \mathcal{Q}_1 \amalg \{a^T \mid a \in \mathcal{Q}_1\}$, where $(a^T)' = \iota(a'')$, $(a^T)'' = \iota(a')$ for all $a \in \mathcal{Q}_1$.

We set $X_{a^T} = X_a^T$ for all $a \in \mathcal{Q}_1$. Zubkov [19] showed that the algebra of *invariants* of mixed representations of \mathcal{Q}

$$I(\mathcal{Q}, \mathbf{n}, \mathbf{1}) = R^{GL(\mathbf{n}, \mathbf{1})}$$

is the subalgebra of R generated by $\sigma_t(X_a)$, where a is a closed path in \mathcal{Q}^D and $1 \leq t \leq \mathbf{n}_{a'}$. As above, we can assume that a is primitive. An example of mixed representations of a quiver is given at the end of the section.

Let $\mathcal{Q}, \mathbf{n}, \mathbf{1}$ be as above. We write $\mathcal{M}(\mathcal{Q}, \mathbf{1})$ for the set of all closed paths in \mathcal{Q}^D and $\mathcal{N}(\mathcal{Q}, \mathbf{1})$ for the subset of primitive paths. Given a path a in \mathcal{Q}^D , we define the path a^T in \mathcal{Q}^D and introduce \sim -equivalence on $\mathcal{M}(\mathcal{Q}, \mathbf{1})$ in the same way as in Section 1. Denote by $\mathcal{M}_{\mathbb{F}}(\mathcal{Q}, \mathbf{1})$ the vector space with the basis $\mathcal{M}(\mathcal{Q}, \mathbf{1})$ and define $\mathcal{N}_{\sigma}(\mathcal{Q}, \mathbf{1})$ in the same way as \mathcal{N}_{σ} have been defined in Section 1. Consider a surjective homomorphism

$$\Upsilon_{\mathbf{n}} : \mathcal{N}_{\sigma}(\mathcal{Q}, \mathbf{1}) \rightarrow I(\mathcal{Q}, \mathbf{n}, \mathbf{1})$$

defined by $\sigma_t(a) \rightarrow \sigma_t(X_a)$, if $t \leq \mathbf{n}_{a'}$, and $\sigma_t(a) \rightarrow 0$ otherwise. Its kernel $K_{\mathbf{n}}(\mathcal{Q}, \mathbf{1})$ is the ideal of relations for $I(\mathcal{Q}, \mathbf{n}, \mathbf{1})$. Elements of $K(\mathcal{Q}, \mathbf{1}) = \bigcap_{m>0} K_m(\mathcal{Q}, \mathbf{1})$ are called *free relations* for $I(\mathcal{Q}, \mathbf{n}, \mathbf{1})$.

Let $u, v \in \mathcal{Q}_0$ be vertices. We say that $a \in \mathcal{M}_{\mathbb{F}}(\mathcal{Q}, \mathbf{1})$ goes from u to v if $a = \sum_i \alpha_i a_i$, where $\alpha_i \in \mathbb{F}$ and $a_i \in \mathcal{M}(\mathcal{Q}, \mathbf{1})$ satisfies $a_i' = u$, $a_i'' = v$. If a goes from u to u , then we say that a is *incident* to u .

Lemma 6.1. *The ideal $K(\mathcal{Q}, \mathbf{1})$ of free relations for $I(\mathcal{Q}, \mathbf{n}, \mathbf{1})$ is zero for an arbitrary p .*

Proof. If f is a free relation, then $f \in K_m(\mathcal{Q}, \mathbf{1})$ for a dimension vector $\mathbf{m} = (m, \dots, m)$ of \mathcal{Q} , where $m > 0$ is arbitrary. By part b) of Lemma 2.3 and part c) of Lemma 2.4, there does not exist a closed path a in \mathcal{Q}^D with $a \stackrel{\sim}{\sim} a^T$. Hence f does not contain a summand with a multiplier $\sigma_t(a)$, where a is a closed path in \mathcal{Q}^D with $a \stackrel{\sim}{\sim} a^T$ and $t > 0$.

Exactly in the same way as we proved Theorem 1.1 for $G = O(m)$, we can show that $f = 0$. Here we do not use the first part of Lemma 3.7, which holds for $p = 2$, but we only need Lemma 3.6, which holds for an arbitrary p . \square

Let us recall that the definition of $\sigma_{t,r}$ can be found in Section 3 of [10].

Theorem 6.2. *The ideal of relations $K_{\mathbf{n}}(\mathcal{Q}, \mathbf{1})$ for $I(\mathcal{Q}, \mathbf{n}, \mathbf{1}) \simeq \mathcal{N}_{\sigma}(\mathcal{Q}, \mathbf{1})/K_{\mathbf{n}}(\mathcal{Q}, \mathbf{1})$ is generated by*

$$\sigma_{t,r}(a, b, c) \in \mathcal{N}_{\sigma}(\mathcal{Q}, \mathbf{1}),$$

where $t+2r > \mathbf{n}_v$ ($t, r \geq 0$), $a, b, c \in \mathcal{M}_{\mathbb{F}}(\mathcal{Q}, \mathbf{1})$, a is incident to some vertex $v \in \mathcal{Q}_0$, b goes from $\iota(v)$ to v , c goes from v to $\iota(v)$.

Proof. As in [20], we denote by $J(\mathcal{Q}, 1)$ the inverse limit of algebras

$$\{I(\mathcal{Q}, \mathbf{n}(1), 1), \varphi_{n(1), n(2)} \mid \mathbf{n}(1) \geq \mathbf{n}(2)\},$$

where $\varphi_{n(1), n(2)} : I(\mathcal{Q}, \mathbf{n}(1), 1) \rightarrow I(\mathcal{Q}, \mathbf{n}(2), 1)$ is the natural epimorphism. It is not difficult to see that $J(\mathcal{Q}, 1) \simeq \mathcal{N}_\sigma(\mathcal{Q}, 1)/K(\mathcal{Q}, 1)$. Lemma 6.1 implies that $J(\mathcal{Q}, 1) \simeq \mathcal{N}_\sigma(\mathcal{Q}, 1)$. By Theorem 2 of [20], the kernel of the natural epimorphism $J(\mathcal{Q}, 1) \rightarrow I(\mathcal{Q}, \mathbf{n}, 1)$ is generated by elements from Theorem 6.2. \square

Example 6.3. Let $(\cdot, \cdot)_1, \dots, (\cdot, \cdot)_r$ be bilinear forms on $V = \mathbb{F}^n$ defined by $n \times n$ matrices A_1, \dots, A_r and $\langle \cdot, \cdot \rangle_1, \dots, \langle \cdot, \cdot \rangle_s$ be bilinear forms on the dual space V^* defined by $n \times n$ matrices B_1, \dots, B_s . Then $G = GL(n)$ acts on the space

$$H = \bigoplus_{k=1}^r \mathbb{F}^{n \times n} \oplus \bigoplus_{l=1}^s \mathbb{F}^{n \times n}$$


of the above mentioned bilinear forms as base change:

$$g \cdot (A_1, \dots, A_r, B_1, \dots, B_s) = (gA_1g^T, \dots, gA_rg^T, g^{-T}B_1g^{-1}, \dots, g^{-T}B_sg^{-1}),$$

where g^{-T} stands for $(g^T)^{-1}$. This action induces the action of $GL(n)$ on the coordinate ring

$$\mathbb{F}[H] = \mathbb{F}[x_{ij}(k), y_{ij}(l), \mid 1 \leq i, j \leq n, 1 \leq k \leq r, 1 \leq l \leq s].$$

Denote generic matrices by $X_k = (x_{ij}(k))$ and $Y_l = (y_{ij}(l))$. Let \mathcal{Q} be the following quiver



where there are r arrows from v to u and s arrows in the opposite direction, $1(u) = v$, and $\mathbf{n} = (n, n)$. Then the algebra of invariants $\mathbb{F}[H]^{GL(n)}$ is isomorphic to $I(\mathcal{Q}, \mathbf{n}, 1)$. By the above mentioned result of Zubkov [19], $\mathbb{F}[H]^{GL(n)}$ is generated by $\sigma_t(Z_1 \cdots Z_m)$, where $1 \leq t \leq n$ and Z_i is one of the following products:

$$X_k Y_l, X_k^T Y_l, X_k Y_l^T, X_k^T Y_l^T \quad (1 \leq k \leq r, 1 \leq l \leq s).$$

Relations between these generators are described by Theorem 6.2.

ACKNOWLEDGEMENTS

The final version of the paper was prepared during author's visit to Max-Planck Institute for Mathematics in Bonn. The author is grateful to MPIM for this support. The research was also supported by RFFI 10-01-00383 and DFG.

REFERENCES

- [1] S.A. Amitsur, *On the characteristic polynomial of a sum of matrices*, Linear and Multilinear Algebra **8** (1980), 177–182.
- [2] H. Derksen, J. Weyman, *Generalized quivers associated to reductive groups*, Colloq. Math. **94** (2002), No. 2., 151–173.
- [3] S. Donkin, *Invariants of several matrices*, Invent. Math. **110** (1992), 389–401.
- [4] S. Donkin, *Invariant functions on matrices*, Math. Proc. Cambridge Philos. Soc. **113** (1993), 23–43.
- [5] S. Donkin, *Polynomial invariants of representations of quivers*, Comment. Math. Helvetici **69** (1994), 137–141.
- [6] A.A. Lopatin, A.N. Zubkov, *Semi-invariants of mixed representations of quivers*, Transform. Groups **12** (2007), N2, 341–369.

- [7] A.A. Lopatin, *Invariants of quivers under the action of classical groups*, J. Algebra **321** (2009), 1079–1106.
- [8] A.A. Lopatin, *On minimal generating systems for matrix $O(3)$ -invariants*, to appear in Linear Mult. Algebra, arXiv: 0902.4270.
- [9] A.A. Lopatin, *Orthogonal invariants of skew-symmetric matrices*, to appear in Linear Mult. Algebra, arXiv: 1004.3082.
- [10] A.A. Lopatin, *Relations between $O(n)$ -invariants of several matrices*, submitted, arXiv: 0902.4266.
- [11] A.A. Lopatin, *Minimal generating set for invariants of bilinear forms in dimension two case*, in preparation.
- [12] C. Procesi, *The invariant theory of $n \times n$ matrices*, Adv. Math. **19** (1976), 306–381.
- [13] Yu.P. Razmyslov, *Trace identities of full matrix algebras over a field of characteristic 0*, Izv. Akad. Nauk SSSR Ser. Mat. **38** (1974), No. 4, 723–756 (Russian).
- [14] D.A. Shmelkin, *Signed quivers, symmetric quivers, and root systems*, J. Lond. Math. Soc., II. Ser. **73** (2006), No. 3, 586–606.
- [15] K.S. Sibirskii, *Algebraic invariants of a system of matrices*, Sibirsk. Mat. Zh. **9** (1968), No. 1, 152–164 (Russian).
- [16] A.N. Zubkov, *A generalization of the Razmyslov–Procesi theorem*, Algebra and Logic **35** (1996), No. 4, 241–254.
- [17] A.N. Zubkov, *Invariants of an adjoint action of classical groups*, Algebra and Logic **38** (1999), No. 5, 299–318.
- [18] A.N. Zubkov, *Mixed representations of quivers and relative problems*, Bielefeld University, SFB **343** (2000), Preprint 00-094.
- [19] A.N. Zubkov, *Invariants of mixed representations of quivers I*, J. Algebra Appl. **4** (2005), No. 3, 245–285.
- [20] A.N. Zubkov, *Invariants of mixed representations of quivers II: Defining relations and applications*, J. Algebra Appl. **4** (2005), No. 3, 287–312.

ARTEM A. LOPATIN
OMSK BRANCH OF
SOBOLEV INSTITUTE OF MATHEMATICS, SB RAS,
PEVTSOVA STREET, 13,
644099, OMSK, RUSSIA
[HTTP://WWW.ITAM.OMSK.NET.RU/~LOPATIN](http://www.itam.omsk.net.ru/~lopatin)
E-mail address: artem_lopatin@yahoo.com