

LARGE DEVIATIONS OF THE EMPIRICAL CURRENTS FOR A BOUNDARY DRIVEN REACTION DIFFUSION MODEL

THIERRY BODINEAU AND MAXIME LAGOUGE

ABSTRACT. We derive a large deviation principle for the empirical currents of lattice gas dynamics which combine a fast stirring mechanism (Symmetric Simple Exclusion Process) and creation/annihilation mechanisms (Glauber dynamics). Previous results on the density large deviations can be recovered from this general large deviation principle. The contribution of external driving forces due to reservoirs at the boundary of the system is also taken into account.

1. INTRODUCTION

A major challenge in non-equilibrium statistical physics is to provide a counterpart to the notion of free energy and to the Gibbs measure which is the cornerstone of the equilibrium statistical physics. Large deviation principles have been proposed as a good alternative to investigate properties of non equilibrium systems [G, D, BDGJL3]. In particular, a lot of attention has been devoted to the case of lattice gas dynamics for which explicit large deviation principles can be derived in the hydrodynamic scaling (we refer the reader to [D, BDGJL3] for recent surveys and further references). Motivated by these recent progress in physics, the original mathematical works [KOV, DV] on the hydrodynamic large deviations for conservative dynamics have been generalized to take into account the contribution of reservoirs at the boundary of the system [BDGJL1, BLM, FLM] and the current of particles flowing in the system [BDGJL2].

An interesting class of models has been introduced in [DFL] to describe reaction diffusion equations by combining the Symmetric Simple Exclusion Process (SSEP) to a Glauber dynamics which models the annihilation and creation of particles. In [DFL], the hydrodynamic limit as well as the fluctuations of the density have been investigated for these models. The density hydrodynamic large deviations have then been proved in [JLV]. In this paper, we generalize this result by deriving the joint large deviations of the density and of the (conservative and non-conservative) currents flowing in the system. We also

Date: September 3, 2010.

2010 Mathematics Subject Classification. 60F10, 82C22.

Key words and phrases. Large deviations, Interacting particle systems.

We are deeply indebted to L. Bertini, B. Derrida, D. Hilhorst, C. Landim, J. Lebowitz for many enlightening discussions and useful suggestions. TB acknowledges the support of the French Ministry of Education through the ANR BLAN07-2184264 grant. This work was partially supported by the NSF Grant DMR-044-2066 and AFOSR Grant AF-FA9550-04 during a stay at Rutgers University and by the *Florence Gould Foundation Endowment* during a stay at the Institute for Advanced Study.

take into account the contribution of reservoirs acting at the boundary of the system. Our results were motivated by the recent research in non-equilibrium statistical physics on dissipative dynamics and in particular on granular media [B, LL, SL]. We refer to [BL] for a more comprehensive discussion on the physical aspects of the large deviations for dissipative systems.

Contrary to the purely conservative dynamics [BDGJL2], one has to introduce two types of currents: the conservative integrated current Q_t which records the particle jumps from the diffusive part of the dynamics (SSEP) and the non-conservative integrated current K_t associated to the creation annihilation process (Glauber). Heuristically, if one denotes by $\dot{Q}_t(r)$ and $\dot{K}_t(r)$ the instantaneous currents at time t and location r , then the density obeys the following equation

$$\partial_t \rho_t(r) = -\partial_r \dot{Q}_t(r) + \dot{K}_t(r).$$

The goal of this paper is to compute the asymptotic cost of observing an atypical trajectory of the currents Q, K and of the density ρ when the number of particles tends to infinity. Even so it is apparently more complicated to consider the joint deviations of the empirical currents and the empirical density, it turns out that the structure of the joint large deviation functional I_0 is more transparent as it splits into two distinct contributions involving either the diffusive part or the Glauber part of the dynamics

$$I_0(\rho, Q, K) = I_1(\rho, Q) + I_2(\rho, K).$$

The precise form of the functional I_0 can be found in (2.12). In section 6, the density large deviation functional derived in [JLV] is recovered by a contraction principle. This provides a natural interpretation of the density large deviation functional as the optimal combination between the two macroscopic currents Q, K in order to create the atypical density trajectory ρ at a minimal cost $I_1 + I_2$.

Our proof relies on the standard machinery developed to study hydrodynamic large deviations [KOV, KL], as well as on more recent tools introduced in [BDGJL1, BDGJL2, BLM, FLM]. Therefore in this paper, we will not detail the aspects of the proof which can be deduced readily from the existing literature and we will focus on the new features occurring from the non-conservative part of the dynamics. The paper is organized as follows. In section 2, we introduce the model and state the main results. A strong form of local equilibrium is stated in section 3. The upper and lower bound of the joint density/current large deviations are derived in sections 4 and 5. Finally the density large deviations are recovered in section 6.

2. NOTATIONS AND RESULTS

2.1. The microscopic dynamics and Notations. We consider the one-dimensional Symmetric Simple Exclusion Process (SSEP) in the domain $\{-N, \dots, N\}$ with creation and annihilation of particles in the bulk and reservoirs at the boundaries. More precisely, the particles perform random walks with an exclusion constraint which imposes at most one particle per site and particles can be removed or created in the bulk according to a rate

which depends on the local configurations. At the boundaries $\pm N$, two reservoirs maintain constant densities. The dynamics can be viewed as a toy model for chemical reactions where the chemicals are injected at the boundaries, then diffuse and react in the system [DFL].

The stochastic dynamics is a Markov process on $\{0, 1\}^{2N+1}$ whose generator is obtained by adding the generators of the different dynamics

$$(2.1) \quad L_N = \frac{N^2}{2}L_{0,N} + \frac{N^2}{2}L_{+,N} + \frac{N^2}{2}L_{-,N} + L_{1,N},$$

with the SSEP generator

$$L_{0,N}f(\eta) = \sum_{x=-N}^{N-1} [f(\eta^{x,x+1}) - f(\eta)],$$

and creation and annihilation generators at the boundaries depending on the parameters β^+ and β^-

$$\begin{aligned} L_{+,N}f(\eta) &= [\eta(N) + \beta_+(1 - \eta(N))] [f(\eta^N) - f(\eta)] \\ L_{-,N}f(\eta) &= [\eta(-N) + \beta_-(1 - \eta(-N))] [f(\eta^{-N}) - f(\eta)] \end{aligned}$$

where

$$\eta^x(z) = \begin{cases} \eta(z) & \text{if } z \neq x \\ 1 - \eta(z) & \text{if } z = x \end{cases}; \quad \eta^{x,y}(z) = \begin{cases} \eta(y) & \text{if } z = x \\ \eta(x) & \text{if } z = y \\ \eta(z) & \text{else} \end{cases}.$$

Finally the creation and annihilation generator in the bulk is given by

$$L_{1,N}f(\eta) = \sum_{x=-N+M+1}^{N-M-1} c(x, \eta) [f(\eta^x) - f(\eta)],$$

where the rate of creation and annihilation $c(x, \cdot)$ is a non negative cylindric function with range M , i.e. there exists a fixed integer M (possibly equal to 0) such that $c(x, \eta) = c(\eta_{x-M}, \dots, \eta_{x+M})$ depends only of the values of η in $\{x - M, \dots, x + M\}$. Remark that, the diffusive part of the process is speeded up by N^2 to obtain a non trivial hydrodynamic evolution.

For a given trajectory $\eta : [0, T] \rightarrow \{0, 1\}^{2N+1}$, let ρ_t^N be the empirical density of particles in $[-1, 1]$ at time $t \in [0, T]$

$$\rho_t^N = \frac{1}{N} \sum_{x=-N}^N \eta_t(x) \delta_{x/N}.$$

We denote by $Q_t^N(x)$ the conservative current through the edge $(x, x + 1)$, i.e. the total number of particles that have jumped from x to $x + 1$ minus the total number of particles that have jumped from $x + 1$ to x between the times 0 and t . The empirical measure

associated to this current is defined as the signed measure on $[-1, 1]$

$$(2.2) \quad Q_t^N = \frac{1}{N^2} \sum_{x=-N}^{N-1} Q_t^N(x) \delta_{x/N}.$$

The renormalization by N^2 takes into account the space renormalization as well as the diffusive scaling of the SSEP dynamics which leads to an extra factor N . We denote by $K_t^N(x)$ the non-conservative current at site x , i.e. the total number of particles created minus the total number of particles annihilated at site x between times 0 and t . The corresponding empirical measure is

$$K_t^N = \frac{1}{N} \sum_{x=-N}^N K_t^N(x) \delta_{x/N}.$$

For any continuous function $\varphi \in C([-1, 1])$, we will use the notation

$$\langle \rho_t^N \varphi \rangle = \frac{1}{N} \sum_{x=-N}^{N-1} \eta_t(x) \varphi\left(\frac{x}{N}\right).$$

The same notation will be used for K_t^N . As the conservative current applies to edges, we will write for $\varphi \in C^1([-1, 1])$

$$\langle Q_t^N \nabla \varphi \rangle = \frac{1}{N} \sum_{x=-N}^{N-1} Q_t^N(x) \left(\varphi\left(\frac{x+1}{N}\right) - \varphi\left(\frac{x}{N}\right) \right).$$

Note that $\varphi\left(\frac{x+1}{N}\right) - \varphi\left(\frac{x}{N}\right)$ is of order $1/N$ so that the scaling is coherent with (2.2). Finally, for any functions $f(s, r)$ in $[0, T] \times [-1, 1]$, we use the shorthand notation

$$\forall s \in [0, T], \quad \langle f_s \rangle = \int_{-1}^1 dr f(s, r).$$

The density profiles bounded away from 0 and 1 will be relevant so that we introduce $C_\epsilon([-1, 1])$ the set of continuous functions f on $[-1, 1]$ for which there exists a constant $\epsilon > 0$ such that $\epsilon < f < 1 - \epsilon$. Given a function $\gamma \in C_\epsilon([-1, 1])$, let ν_γ^N be the Bernoulli product measure on $\{-N, N\}$ with marginals

$$(2.3) \quad \forall k \in \{-N, N\}, \quad \nu_\gamma^N(\eta(k) = 1) = \gamma\left(\frac{k}{N}\right).$$

For $\alpha \in [0, 1]$, the Bernoulli product measure with uniform density α is denoted by ν_α^N . Let \mathbb{P}_γ^N be the probability measure associated to the Markov process (ρ_t^N, Q_t^N, K_t^N) on $[0, T]$ with initial measure ν_γ^N on the particle configurations.

We define \mathcal{M} the set of signed measures on $[-1, 1]$ endowed with the weak topology. We also consider \mathcal{M}_0 the subset of \mathcal{M} of all absolutely continuous measures wrt the Lebesgue measure with positive density bounded by 1

$$\mathcal{M}_0 = \{\rho(x)dx \in \mathcal{M}, \quad 0 \leq \rho(x) \leq 1 \quad a.e.\}.$$

In order to consider the joint large deviations of (ρ_t^N, Q_t^N, K_t^N) during the time interval $[0, T]$, we will work on $\mathcal{E} = D([0, T], \mathcal{M}_0 \times \mathcal{M} \times \mathcal{M})$ the space of cad-lag trajectories with values in $\mathcal{M}_0 \times \mathcal{M} \times \mathcal{M}$ endowed with the Skorohod topology [EK].

2.2. The results. The hydrodynamic behavior of the microscopic dynamics introduced in section 2.1, can be described in terms of a few macroscopic parameters

$$(2.4) \quad \forall \alpha \in [0, 1], \quad C(\alpha) = \nu_\alpha(c(0, \eta)(1 - \eta(0))), \quad A(\alpha) = \nu_\alpha(c(0, \eta)\eta(0)),$$

where C and A represent the average creation and annihilation rates at density α and

$$(2.5) \quad \forall \alpha \in [0, 1], \quad \sigma(\alpha) = \nu_\alpha(\eta_0(1 - \eta_1)) = \alpha(1 - \alpha),$$

which is the conductivity of the SSEP. Finally, we denote by $\bar{\rho}_\pm = \frac{\beta_\pm}{1 + \beta_\pm}$ the densities at the boundaries imposed by the reservoirs.

We are now ready to state the hydrodynamic limit

Theorem 2.1. *Let $\gamma \in C_e([-1, 1])$ with $\gamma(-1) = \bar{\rho}_-$ and $\gamma(1) = \bar{\rho}_+$. For each $T > 0$ and $\varphi \in C([0, T] \times [-1, 1])$,*

$$\forall \delta > 0, \quad \lim_{N \rightarrow \infty} \mathbb{P}_\gamma^N \left[\left| \int_0^T dt \langle \rho_t^N \varphi_t \rangle - \langle \bar{\rho}_t \varphi_t \rangle \right| > \delta \right] = 0,$$

where $\bar{\rho}(t, x)$ is the unique weak solution of

$$(2.6) \quad \begin{cases} \partial_t \bar{\rho}(t, x) &= \frac{1}{2} \Delta \bar{\rho}(t, x) + C(\bar{\rho}(t, x)) - A(\bar{\rho}(t, x)) \\ \bar{\rho}(t, \pm 1) &= \bar{\rho}_\pm \\ \bar{\rho}(0, x) &= \gamma(x) \end{cases}$$

The meaning of weak solution of (2.6) is recalled in the Appendix (with $H = G = 0$).

As a consequence of Theorem 2.1 a law of large numbers holds for the currents.

Theorem 2.2. *For any test function $\varphi \in C^1([-1, 1])$ and $t > 0$*

$$\begin{aligned} \forall \delta > 0, \quad \lim_{N \rightarrow \infty} \mathbb{P}_\gamma^N \left[\left| \langle Q_t^N \varphi \rangle + \frac{1}{2} \int_0^t ds \langle \varphi \nabla \bar{\rho}_s \rangle \right| > \delta \right] &= 0, \\ \lim_{N \rightarrow \infty} \mathbb{P}_\gamma^N \left[\left| \langle K_t^N \varphi \rangle - \int_0^t ds \langle \varphi (C(\bar{\rho}_s) - A(\bar{\rho}_s)) \rangle \right| > \delta \right] &= 0, \end{aligned}$$

where $\bar{\rho}$ is the unique weak solution of (2.6).

Theorems 2.1, 2.2 can be deduced from the methods used to prove the large deviations so that their derivation is omitted (see also [DFL, BDGJL2]).

Before stating the large deviation principle, we need more notation. Let G and H be smooth functions in $[0, T] \times [-1, 1]$. For a given trajectory (ρ, Q, K) in \mathcal{E} , we set

$$(2.7) \quad J_{G,H}(\rho, Q, K) = J_H^1(\rho, Q) + J_G^2(\rho, K)$$

with

(2.8)

$$\begin{aligned} J_H^1(\rho, Q) &= \langle Q_T \nabla H_T \rangle - \int_0^T \left\langle Q_s \frac{d}{ds} \nabla H_s \right\rangle ds - \frac{1}{2} \int_0^T \langle \rho_s \Delta H_s \rangle ds \\ &\quad - \frac{1}{2} \int_0^T \langle \sigma(\rho_s) (\nabla H_s)^2 \rangle ds + \frac{1}{2} \bar{\rho}_+ \int_0^T \nabla H(s, 1) ds - \frac{1}{2} \bar{\rho}_- \int_0^T \nabla H(s, -1) ds, \end{aligned}$$

where $\sigma(u) = u(1 - u)$ is defined in (2.5) and

(2.9)

$$J_G^2(\rho, K) = \langle K_T G_T \rangle - \int_0^T \left\langle K_s \frac{d}{ds} G_s \right\rangle ds - \int_0^T ds \langle C(\rho_s)(e^{G_s} - 1) + A(\rho_s)(e^{-G_s} - 1) \rangle,$$

where A and C were introduced in (2.4).

The first functional is related to the contribution of the conservative currents and the second one to the non-conservative currents (see theorem 2.4). We define

$$J(\rho, Q, K) = \sup_{G, H} J_{G, H}(\rho, Q, K) = \sup_{H \in C^{1,2}} J_H^1(\rho, Q) + \sup_{G \in C^{1,0}} J_G^2(\rho, K),$$

where the supremum is taken on regular functions G and H . Note that the functions G and H can take arbitrary (finite) values at the boundaries.

Define \mathcal{A} as the set of trajectories (ρ, Q, K) satisfying the following two conditions:

- *Conservation law.* For all test function $\varphi \in C^1([-1, 1])$ vanishing at the boundaries

$$(2.10) \quad \langle \rho_t \varphi \rangle - \langle \rho_0 \varphi \rangle = \langle Q_t \partial_x \varphi \rangle + \langle K_t \varphi \rangle, \quad Q_0 = 0, \quad K_0 = 0.$$

- *Energy condition.* The energy $\mathcal{Q}(\rho)$ of the density trajectory is finite with

$$(2.11) \quad \mathcal{Q}(\rho) = \sup_{\varphi} \left\{ \int_0^T dt \int_{-1}^1 dx \rho(t, x) \nabla \varphi(t, x) - \frac{1}{2} \int_0^T dt \int_{-1}^1 dx \varphi(t, x)^2 - \int_0^T dt (\bar{\rho}_+ \varphi(t, 1) - \bar{\rho}_- \varphi(t, -1)) \right\},$$

and the supremum is taken over smooth functions φ in $[0, T] \times [-1, 1]$.

For smooth trajectories the conservation law reduces to

$$\partial_t \rho + \partial_x \partial_t Q - \partial_t K = 0,$$

where $\partial_t Q, \partial_t K$ are the instantaneous currents and the energy condition reads

$$\mathcal{Q}(\rho) = \frac{1}{2} \int_0^T dt \int_{-1}^1 dx (\nabla \rho(t, x))^2 < \infty.$$

The energy condition was introduced in [QRV, BLM, FLM] to control the approximation procedure in the derivation of the large deviation lower bound (see Theorem 2.6).

Finally we define the dynamical rate function

$$(2.12) \quad I_0(\rho, Q, K) = \begin{cases} J(\rho, Q, K) & \text{if } (\rho, Q, K) \in \mathcal{A}, \\ +\infty & \text{otherwise.} \end{cases}$$

To take into account the large deviations of the initial measure ν_γ^N , we introduce for any function $m : [-1, 1] \rightarrow [0, 1]$

$$h_\gamma(m) = \left\langle m \log \frac{m}{\gamma} \right\rangle + \left\langle (1-m) \log \frac{1-m}{1-\gamma} \right\rangle,$$

and

$$I_\gamma(\mu) = \begin{cases} h_\gamma(m) & \text{if } \mu(dx) = m(x) dx, \\ +\infty & \text{otherwise.} \end{cases}$$

The rate function is the sum of the dynamical deviation cost from the initial measure and the deviation cost from the hydrodynamic trajectory

$$I(\rho, Q, K) = I_0(\rho, Q, K) + I_\gamma(\rho_0).$$

From now, the initial density profile γ is a given smooth function in $C_e([-1, 1])$ equal to $\bar{\rho}_\pm$ at the boundaries. We state the large deviation Theorems.

Theorem 2.3. *For all closed set $\mathcal{F} \in \mathcal{E}$*

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N [(\rho_t^N, Q_t^N, K_t^N) \in \mathcal{F}] \leq - \inf_{(\rho, Q, K) \in \mathcal{F}} I(\rho, Q, K).$$

We first state the lower bound for regular trajectories.

Theorem 2.4. *Let (ρ, Q, K) be a regular trajectory. Then the large deviation functional I_0 has an explicit form*

$$(2.13) \quad I_0(\rho, Q, K) = \int_0^T dt \int_0^1 dx \left\{ \frac{(\partial_t Q(x, t) + \partial_x \rho(x, t))^2}{2\sigma(\rho(x, t))} + \Phi(\rho(x, t), \partial_t K(x, t)) \right\},$$

with

$$(2.14) \quad \Phi(\rho, \kappa) = C(\rho) + A(\rho) - \sqrt{\kappa^2 + 4A(\rho)C(\rho)} + \kappa \log \left(\frac{\sqrt{\kappa^2 + 4A(\rho)C(\rho)} + \kappa}{2C(\rho)} \right).$$

If $C(\rho) = 0$, then Φ becomes

$$\Phi(\rho, \kappa) = \begin{cases} A(\rho) + \kappa - \kappa \log \left(\frac{-\kappa}{A(\rho)} \right), & \text{if } \kappa \leq 0, \\ \infty, & \text{if } \kappa > 0. \end{cases}$$

A similar formula holds if $A(\rho) = 0$.

For any open set $\mathcal{O} \in \mathcal{E}$ containing the regular trajectory (ρ, Q, K)

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N [(\rho_t^N, Q_t^N, K_t^N) \in \mathcal{O}] \geq -I(\rho, Q, K).$$

Remark 2.5. *The first contribution to (2.13) comes from the difference between the instantaneous empirical current $\partial_t Q$ and the canonical instantaneous current associated to ρ which is $-\frac{1}{2}\nabla\rho$. This term has already been analyzed in the conservative dynamics [BD, BDGJL2]. The second term in (2.13) should be interpreted as the large deviation functional associated to Poisson processes with parameters $C(\rho_t)$ and $A(\rho_t)$.*

In order to derive the lower bound for general trajectories, we introduce two technical assumptions on the rates (2.4):

Assumption (L1). *The rate A (resp C) is either concave and positive on $]0, 1[$ or uniformly equal to zero.*

Assumption (L2). *The functions A and C are monotonous and*

$$(2.15) \quad \forall z \in [0, 1], \quad A'(z) \geq 0, \quad C'(z) \leq 0.$$

Then

Theorem 2.6. *Assume (L1) and (L2), then for all open set $\mathcal{O} \in \mathcal{E}$,*

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N [(\rho_t^N, Q_t^N, K_t^N) \in \mathcal{O}] \geq - \inf_{(\rho, Q, K) \in \mathcal{O}} I(\rho, Q, K).$$

The concavity assumption (L1) has been introduced in [JLV]. As we shall see in Section 5.2, it simplifies the proof of the lower bound, however it is mainly technical and Theorem 2.6 should be valid without assumption (L1). We refer to [QRV, BLM, FLM] for further results on this generalization in the case of conservative dynamics. Assumption (L2) will be used in the Appendix only to ensure the uniqueness of the weak solutions for singular perturbations of the hydrodynamic equation (2.6).

3. MODIFIED DYNAMICS AND LOCAL EQUILIBRIUM

Local equilibrium lies at the heart of the hydrodynamic limit theory and it states that during the time evolution the local measure remains close to an equilibrium measure with a varying density. In this section, we state, in our framework, a strong form of local equilibrium which will be useful for the derivation of the hydrodynamic large deviations. The proofs are omitted as they follow the scheme introduced in [KOV, KL, BDGJL1].

3.1. The modified dynamics. We first define a modification of the original process (2.1) which will be used to derive the large deviations. For regular functions G and H on $[0, T] \times [-1, 1]$, denote $L_N^{G_t, H_t}$ the time dependent generator given by

$$(3.1) \quad L_N^{G_t, H_t} = \frac{N^2}{2} L_{0,N}^{H_t} + L_{1,N}^{G_t} + \frac{N^2}{2} L_{+,N} + \frac{N^2}{2} L_{-,N},$$

with

$$L_{0,N}^{H_t} f(\eta) = \sum_{x=-N}^{N-1} [f(\eta^{x,x+1}) - f(\eta)] \exp \left[(\eta(x) - \eta(x+1)) \left(H(t, \frac{x+1}{N}) - H(t, \frac{x}{N}) \right) \right],$$

$$L_{1,N}^{G_t} f(\eta) = \sum_{x=-N}^N c(x, \eta) [\eta(x) e^{-G(t,x/N)} + (1 - \eta(x)) e^{G(t,x/N)}] [f(\eta^x) - f(\eta)].$$

The modified dynamics induces a weak drift in the conservative dynamics. For large N , a particle jumps from x to $x \pm 1$ at rate $\frac{1}{2} \left(1 \pm \frac{1}{N} \partial_x H(t, \frac{x}{N}) \right)$. $L_{1,N}^{G_t}$ is the generator of a non-conservative dynamics for which the intensity of creation and annihilation varies in time and space according to G . Finally, let $\mathbb{P}_{\gamma,G,H}^N$ be the probability measure associated to the process with initial measure ν_γ^N and generator $L_N^{G_t, H_t}$. We stress the fact that the reservoir dynamics are unchanged. As in Theorem 2.1, one can show that the modified dynamics follows the hydrodynamic limit equation

$$(3.2) \quad \partial_t \rho(t, x) = \frac{1}{2} \Delta \rho(t, x) - \nabla \cdot (\sigma(\rho(t, x)) \nabla H(t, x)) + C(\rho(t, x)) e^{G(t,x)} - A(\rho(t, x)) e^{-G(t,x)}.$$

3.2. Local equilibrium. We set $\Lambda_l^{(x)} = \{y \in \{-N, \dots, N\} \text{ such that } |y - x| \leq l\}$ and define the local density as

$$(3.3) \quad \bar{\eta}^l(x) = \frac{1}{|\Lambda_l^{(x)}|} \sum_{y \in \Lambda_l^{(x)}} \eta(y),$$

where $|\Lambda_l^{(x)}|$ stands for the number of sites in $\Lambda_l^{(x)}$. Let ψ be a cylindric function with support in $\{-R, \dots, R\}$. Given φ a regular function on $[0, T] \times [-1, 1]$ and $\delta, \varepsilon > 0$, we define the set $B_{\delta, \varepsilon, \varphi}(\psi)$ on the trajectories $\{\eta_s\}_{s \leq T}$

$$(3.4) \quad B_{\delta, \varepsilon, \varphi}(\psi) = \left\{ \{\eta_s\}_{s \leq T}, \quad \left| \int_0^T ds \frac{1}{2(N-R)} \sum_{x=-N+R}^{N-R} \varphi\left(s, \frac{x}{N}\right) [\psi(\tau_x \eta_s) - \nu_{\bar{\eta}_s^{\varepsilon N}(x)}(\psi)] \right| \leq \delta \right\},$$

where $\tau_x \eta$ is the configuration η shifted by x .

The local equilibrium property also holds for the modified dynamics (3.1).

Theorem 3.1. *Given φ, ψ and $\delta > 0$, the trajectories concentrate super-exponentially fast on the set $B_{\delta, \varepsilon, \varphi}(\psi)$*

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_{\gamma, G, H}^N \left(B_{\delta, \varepsilon, \varphi}(\psi)^c \right) = -\infty.$$

Moreover, the reservoirs impose local equilibrium at the boundaries with the densities $\bar{\rho}_+, \bar{\rho}_-$. For any continuous function Φ in $[0, T]$

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_{\gamma, G, H}^N \left[\left| \int_0^T ds \Phi(s) (\eta_s(\pm N) - \bar{\rho}_\pm) \right| \geq \delta \right] = -\infty.$$

The derivation of Theorem 3.1 follows from the bounds on entropy production [KOV] which can be adapted to control the boundary terms as in [BDGJL1].

4. LARGE DEVIATION UPPER BOUND

The derivation of Theorem 2.3 is split into several steps. First an upper bound with the rate function J (2.12) is derived for compact sets, then for closed sets. Finally, we prove that the rate function is infinite for the trajectories which do not belong to the set \mathcal{A} introduced in (2.10), (2.11).

4.1. The upper bound for compact sets. In order to compare the original dynamics starting from the initial profile γ to a modified dynamics with regular drifts G, H (3.1) starting from the initial profile ω , we compute the Radon-Nikodym derivative [KL]

$$(4.1) \quad \frac{d\mathbb{P}_{\omega, G, H}^N}{d\mathbb{P}_{\gamma}^N} = \exp \left[N \left\{ \langle Q_T^N \nabla H_T \rangle - \int_0^T ds \langle Q_s^N \frac{d}{ds} \nabla H_s \rangle + \langle K_T^N G_T \rangle - \int_0^T ds \langle K_s^N \frac{d}{ds} G_s \rangle \right. \right. \\ \left. \left. + O_H\left(\frac{1}{N}\right) \right\} - \int_0^T R_N(s) ds + h_{\gamma, \omega}(\rho_0^N) \right],$$

where $h_{\gamma, \omega}(\rho) = \langle \rho \log(\omega/\gamma) \rangle + \langle (1 - \rho) \log((1 - \omega)/(1 - \gamma)) \rangle$ and

$$(4.2) \quad R_N(s) = \frac{N^2}{2} \left(\sum_{x=-N}^{N-1} \eta_s(x)(1 - \eta_s(x+1)) \left[\exp \left(H \left(s, \frac{x+1}{N} \right) - H \left(s, \frac{x}{N} \right) \right) - 1 \right] \right. \\ \left. + \eta_s(x+1)(1 - \eta_s(x)) \left[\exp \left(H \left(s, \frac{x}{N} \right) - H \left(s, \frac{x+1}{N} \right) \right) - 1 \right] \right) \\ + \sum_{x=-N}^N c(x, \eta_s) \left[\exp \left((1 - 2\eta_s(x)) G \left(s, \frac{x}{N} \right) \right) - 1 \right].$$

expanding the exponential and summing by parts, we get

$$R_N(s) = \frac{1}{2} \left\{ \sum_{x=-N}^{N-1} \left[\eta_s(x) \Delta H \left(s, \frac{x}{N} \right) \right. \right. \\ \left. \left. + \frac{1}{4} \left(\nabla H \left(s, \frac{x}{N} \right) \right)^2 [\eta_s(x)(1 - \eta_s(x+1)) + \eta_s(x+1)(1 - \eta_s(x))] \right] \right\} \\ + \frac{1}{2} \left[\eta_s(N) \nabla H(s, 1) - \eta_s(-N) \nabla H(s, -1) \right] \\ + \sum_{x=-N}^N c(x, \eta_s) \left[\exp \left((1 - 2\eta_s(x)) G \left(s, \frac{x}{N} \right) \right) - 1 \right] + O(1).$$

Thanks to the local equilibrium, the microscopic expressions in $R_N(s)$ can be replaced by their averages. We set

$$B_{\delta,\varepsilon} = B_{\delta,\varepsilon,(\nabla H)^2}(\eta_0(1-\eta_0)) \cap B_{\delta,\varepsilon,(e^{G-1})}(c(0,\eta)\eta_0) \cap B_{\delta,\varepsilon,(e^{-G-1})}(c(0,\eta)(1-\eta_0)) \cap B'_{\delta,\pm}, \quad (4.3)$$

where the sets $B_{\delta,\varepsilon,\varphi}(\cdot)$ were introduced in (3.4) and

$$B'_{\delta,\pm} = \left\{ \left| \int_0^T ds \nabla H(s, \pm 1) (\eta_s(\pm N) - \bar{\rho}_\pm) \right| \leq \delta \right\}.$$

The super-exponential replacement Theorem 3.1 implies

$$\forall \delta > 0, \quad \limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N(B_{\delta,\varepsilon}^c) = -\infty.$$

For trajectories in $B_{\delta,\varepsilon}$, the Radon-Nikodym derivative can be approximated as follows

$$\frac{d\mathbb{P}_{\omega,G,H}^N}{d\mathbb{P}_\gamma^N} = \exp \left[N \left(J_{G,H,\varepsilon}(\rho^N, Q^N, K^N) + h_{\gamma,\omega}(\rho_0^N) + O_{G,H}\left(\frac{1}{N}\right) + O(\delta) \right) \right],$$

where we used the functional

(4.4)

$$\begin{aligned} J_{G,H,\varepsilon}(\rho, Q, K) &= \langle Q_T \nabla H_T \rangle - \int_0^T ds \langle Q_s \frac{d}{ds} \nabla H_s \rangle + \langle K_T G_T \rangle - \int_0^T ds \langle K_s \frac{d}{ds} G_s \rangle \\ &\quad - \frac{1}{2} \int_0^T ds \langle \rho_s \Delta H_s + \sigma(\rho_s * i_\varepsilon)(\nabla H_s)^2 \rangle \\ &\quad - \int_0^T ds \langle C(\rho_s * i_\varepsilon)(e^{G_s} - 1) + A(\rho_s * i_\varepsilon)(e^{-G_s} - 1) \rangle \\ &\quad + \frac{1}{2} \bar{\rho}_+ \int_0^T ds \nabla H(s, 1) - \frac{1}{2} \bar{\rho}_- \int_0^T ds \nabla H(s, -1), \end{aligned}$$

with A, C and σ as in (2.4), (2.5). The function i_ε is an approximation of unity

$$\rho_s^N * i_\varepsilon = \bar{\eta}_s^{\varepsilon N}.$$

For any \mathcal{O} open set in \mathcal{E} , we can then deduce the large deviation upper bound

$$\begin{aligned} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N(\mathcal{O}) &\leq \max \left[\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N((\rho^N, Q^N, K^N) \in \mathcal{O} \cap B_{\delta,\varepsilon}); \right. \\ &\quad \left. \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N(B_{\delta,\varepsilon}^c) \right] \\ &\leq \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_{\omega,G,H}^N \left[\frac{d\mathbb{P}_\gamma^N}{d\mathbb{P}_{\omega,G,H}^N} \mathbf{1}_{\{\mathcal{O} \cap B_{\delta,\varepsilon}\}}(\rho^N, Q^N, K^N) \right] \\ &\leq \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_{\omega,G,H}^N \left[e^{-N(J_{G,H,\varepsilon}(\rho^N, Q^N, K^N) + h_{\gamma,\omega}(\rho_0^N) + O_{G,H}(\frac{1}{N}) + O(\delta))} \mathbf{1}_{\mathcal{O}} \right] \\ &\leq \sup_{(\rho, Q, K) \in \mathcal{O}} \left\{ -J_{G,H,\varepsilon}(\rho, Q, K) - h_{\gamma,\omega}(\rho_0) \right\} + O(\delta). \end{aligned}$$

This is true for any (ϵ, G, H, ω) and any $\delta > 0$ so finally

$$(4.5) \quad \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N(\mathcal{O}) \leq \inf_{\epsilon, G, H, \omega} \sup_{(\rho, Q, K) \in \mathcal{O}} \left\{ -J_{G, H, \epsilon}(\rho, Q, K) - h_{\gamma, \omega}(\rho_0) \right\}.$$

The previous bound can then be extended to any compact set \mathcal{K} by using a finite covering with open sets (see [KL])

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N(\mathcal{K}) \leq - \sup_{(\rho, Q, K) \in \mathcal{K}} \sup_{G, H} \{J_{G, H}(\rho, Q, K)\} = -J(\rho, Q, K).$$

4.2. The upper bound for closed sets. We are going to prove the exponential tightness, i.e. to exhibit a sequence $\{\mathcal{K}_n\}$ of compact sets in \mathcal{E} such that for any n

$$(4.6) \quad \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N(\mathcal{K}_n^c) \leq -n.$$

The large deviation upper bound will then follow for general closed sets F of \mathcal{E} by noticing

$$\begin{aligned} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N(F) &\leq \limsup_{N \rightarrow \infty} \frac{1}{N} \log [\mathbb{P}_\gamma^N(F \cap \mathcal{K}_n) + \mathbb{P}_\gamma^N(\mathcal{K}_n^c)] \\ &\leq \max \left[\sup_{(\rho, Q, K) \in F \cap \mathcal{K}_n} J(\rho, Q, K) ; -n \right]. \end{aligned}$$

Letting n go to ∞ completes the upper bound for closed sets.

In order to build a sequence of compact sets, we need to check first that the measures concentrate on equi-continuous trajectories

Lemma 4.1. *Let ϕ be a C^1 function on $[-1, 1]$. Then we have for all $\epsilon > 0$,*

$$(4.7) \quad \lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N \left[\sup_{|t-s| \leq \delta} |\langle \rho_t^N \phi \rangle - \langle \rho_s^N \phi \rangle| > \epsilon \right] = -\infty,$$

$$(4.8) \quad \lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N \left[\sup_{|t-s| \leq \delta} |\langle Q_t^N \phi \rangle - \langle Q_s^N \phi \rangle| > \epsilon \right] = -\infty,$$

$$(4.9) \quad \lim_{\delta \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N \left[\sup_{|t-s| \leq \delta} |\langle K_t^N \phi \rangle - \langle K_s^N \phi \rangle| > \epsilon \right] = -\infty,$$

where the supremum is taken over s, t in $[0, T]$.

Second, we need estimates on the total variation norm of the empirical currents

Lemma 4.2. *For any time $T > 0$, one has*

$$(4.10) \quad \lim_{a \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N \left(\sup_{0 \leq t \leq T} \frac{1}{N} \sum_{x=-N}^N |K_t^N(x)| \geq a \right) = -\infty,$$

$$(4.11) \quad \lim_{a \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N \left(\sup_{0 \leq t \leq T} \frac{1}{N^2} \sum_{x=-N}^N |Q_t^N(x)| \geq a \right) = -\infty.$$

Should the currents be bounded (as the density), then Lemma 4.1 would be enough to ensure the exponential tightness [KL].

We postpone the derivation of the Lemmas and conclude the proof of the exponential tightness (4.6). We focus on the conservative current as the same strategy applies to ρ, K . Fix a sequence ϕ_l of C^2 functions on $[-1, 1]$ dense in $C([-1, 1])$. We set

$$\mathcal{C}_{l,\delta,m} = \left\{ Q \in D([0, T], \mathcal{M}) ; \quad \sup_{|t-s| \leq \delta} \left| \langle Q_t \phi_l \rangle - \langle Q_s \phi_l \rangle \right| \leq \frac{1}{m} \right\}.$$

Using Lemma 4.1, we have,

$$\forall l \geq 0, \forall m \geq 1, \forall n \geq 1, \exists \delta(n, m, l), \quad \mathbb{P}_\gamma^N(Q^N \notin \mathcal{C}_{l,\delta(n,m,l),m}) \leq \exp(-Nnml).$$

We introduce also $\mathcal{C}'_n = \{Q \in D([0, T], \mathcal{M}) ; \sup_{t \leq T} |Q_t| \leq k(n)\}$, where $|Q_t|$ stands for the total variation norm of the measure Q_t and $k(n)$ is chosen, according to Lemma 4.2, such that

$$\mathbb{P}_\gamma^N(Q^N \notin \mathcal{C}'_n) \leq \exp(-Nn).$$

Now consider

$$(4.12) \quad \mathcal{K}_n = \bigcap_{l \geq 0, m \geq 1} \mathcal{C}_{l,\delta(n,m,l),m} \bigcap \mathcal{C}'_n.$$

Ascoli Theorem (see [EK] Theorem 6.3 page 123) implies that \mathcal{K}_n is a compact set. Combining the previous estimates we see that \mathcal{K}_n satisfies (4.6).

Proof of Lemma 4.1. We start by proving (4.8) and follow the strategy of [BDGJL2]. It is enough to show that the expression below goes to $-\infty$ as δ vanishes

$$\max_{0 \leq k \leq \frac{T}{\delta}} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N \left[\sup_{k\delta \leq t \leq (k+1)\delta} \langle Q_t^N \phi \rangle - \langle Q_{k\delta}^N \phi \rangle > \epsilon \right].$$

Thanks to the computation of the Radon-Nikodym derivative of the modified dynamics (4.2) (with $\nabla H = \frac{1}{N}\phi, G = 0$), we know from (4.1) that for all $a > 0$,

$$\begin{aligned} \mathcal{M}_T = & \exp \left\{ N \langle Q_T^N a \phi \rangle - \frac{1}{2} \int_0^T ds \sum_{x=-N}^{N-1} \left(\eta_s(x) a \nabla \phi \left(\frac{x}{N} \right) \right. \right. \\ & \left. \left. + \frac{1}{4} \left[a \phi \left(\frac{x}{N} \right) \right]^2 [\eta_s(x)(1 - \eta_s(x+1)) + \eta_s(x+1)(1 - \eta_s(x))] \right) \right. \\ & \left. + N [\eta_s(N)\phi(1) - \eta_s(-N)\phi(-1)] + O(1) \right\} \end{aligned}$$

is a mean one positive martingale. One easily checks that the integral term is bounded above by $C_\phi a^2 NT$ (we will take the limit as a goes to infinity), where C_ϕ is a constant depending only on ϕ . Therefore, multiplying by aN , adding and subtracting the integral term of the logarithm of the martingale and exponentiating,

$$(4.13) \quad \mathbb{P}_\gamma^N \left[\sup_{k\delta \leq t \leq (k+1)\delta} \langle Q_t^N \phi \rangle - \langle Q_{k\delta}^N \phi \rangle > \epsilon \right] \leq \mathbb{P}_\gamma^N \left[\sup_{k\delta \leq t \leq (k+1)\delta} \frac{\mathcal{M}_t}{\mathcal{M}_{\delta k}} > \exp \{ \epsilon a N - C_\phi a^2 N \delta \} \right].$$

Taking δ small enough such that $C_\phi a^2 N \delta < \frac{\epsilon a N}{2}$, then Doob's inequality implies that the last expression is bounded above by $\exp[-aN\epsilon/4]$. Letting a go to ∞ completes the proof.

For the non-conservative current, (4.9) follows in the same way by using the martingale

$$\mathcal{N}_t = \exp \left[\langle a\phi K_T^N \rangle - \int_0^T ds \sum_{x=-N}^N c(x, \eta_s) [e^{a(1-2\eta_s(x))\phi(x/N)} - 1] \right].$$

The proof of (4.7) can be found in [KL]. \square

Proof of Lemma 4.2. The probability of the event in (4.10) can be estimated from above by the large deviations of $(2N+1)$ independent Poisson processes thus

$$(4.14) \quad \forall a > 0, \quad \mathbb{P}_\gamma^N \left(\sup_{0 \leq t \leq T} \frac{1}{N} \sum_{x=-N}^N |K_t^N(x)| \geq a \right) \leq \exp(-NC_a),$$

where C_a goes to infinity as a diverges.

We turn now to the bound (4.11) and show that

$$(4.15) \quad \forall a > 0, \quad \mathbb{P}_\gamma^N \left(\sup_{0 \leq t \leq T} \frac{1}{N^2} \sum_{x=-N}^N |Q_t^N(x)| \geq a \right) \leq e^{-NC'_a},$$

where C'_a goes to infinity as a diverges. To prove (4.15), we first use a microscopic identity (which holds at any time)

$$(4.16) \quad \forall x \in \{-N, N\}, \quad Q_t^N(x-1) - Q_t^N(x) = \eta_t^N(x) - \eta_0^N(x) - K_t^N(x).$$

Therefore (4.14) implies that with probability at least $1 - e^{-NC_a}$, the conservative current through the edge $(x, x+1)$ satisfies

$$\forall t \in [0, T], \quad |Q_t^N(x)| \leq |Q_t^N(-N)| + 2N(1+a).$$

From this, we get

$$(4.17) \quad \begin{aligned} & \mathbb{P}_\gamma^N \left(\sup_{0 \leq t \leq T} \frac{1}{N^2} \sum_{x=-N}^{N-1} |Q_t^N(x)| \geq 5a \right) \leq e^{-NC_a} + \mathbb{P}^N \left(\sup_{0 \leq t \leq T} \frac{|Q_t^N(-N)|}{N} \geq 3a - 2 \right), \\ & \leq e^{-NC_a t} + \mathbb{P}_\gamma^N \left(\sup_{0 \leq t \leq T} \frac{Q_t^N(-N)}{N} \geq 3a - 2 \right) + \mathbb{P}_\gamma^N \left(\inf_{0 \leq t \leq T} \frac{Q_t^N(-N)}{N} \leq -3a + 2 \right). \end{aligned}$$

We bound now the first term on the RHS (the second one can be bounded similarly by symmetry). Using again the identity (4.16), we get

$$\mathbb{P}_\gamma^N \left(\sup_{0 \leq t \leq T} \frac{Q_t^N(-N)}{N} \geq 3a - 2 \right) \leq \mathbb{P}_\gamma^N \left(\sup_{0 \leq t \leq T} \frac{1}{N^2} \sum_{x=-N}^{N-1} Q_t^N(x) \geq a - 4 \right) + e^{-NC_a t}.$$

The terms in the RHS can be estimated as in (4.13) by reducing to a martingale estimate. This completes (4.15). \square

4.3. The set \mathcal{A} . To complete the derivation of the upper bound, we prove that the trajectories concentrate exponentially fast on the set \mathcal{A} introduced in (2.10), (2.11).

The conservation law.

Let $C_0^2([-1, 1])$ be the set of twice differentiable functions vanishing at the boundary.

Lemma 4.3. *For any $\phi \in C_0^2([-1, 1])$, we introduce*

$$V_T(\rho^N, Q^N, K^N, \phi) = \frac{1}{N} \sum_{x=-N+1}^{N-1} \phi\left(\frac{x}{N}\right) (\eta_T(x) - \eta_0(x)) - \frac{1}{N} \nabla \phi\left(\frac{x}{N}\right) Q_T^N(x) + \phi\left(\frac{x}{N}\right) K_T(x).$$

Then for any $\delta > 0$,

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N [|V_T(\rho^N, Q^N, K^N, \phi)| > \delta] = -\infty.$$

Proof. For any site x and times $s < t$, the following microscopic relation holds

$$\eta_t^N(x) - \eta_s^N(x) = [Q_t^N(x-1) - Q_s^N(x-1)] - [Q_t^N(x) - Q_s^N(x)] + K_t^N(x) - K_s^N(x).$$

Summing in x and integrating by parts the Q -term gives

$$V_T(\rho^N, Q^N, K^N, \phi) = \frac{1}{N} \sum_{x=-N}^{N-1} W_x Q_T^N(x), \quad \text{with } W_x = \int_{x/N}^{(x+1)/N} du \left(\frac{x+1}{N} - u \right) \phi''(u),$$

where we used that ϕ vanishes at the boundaries. From this identity we get for any $a > 0$

$$(4.18) \quad \mathbb{P}_\gamma^N (V_T(\rho^N, Q^N, K^N, \phi) > \delta) \leq e^{-aN\delta} \mathbb{P}_\gamma^N \left(\exp \left(a \sum_{x=-N}^{N-1} W_x Q_T^N(x) \right) \right).$$

We first note that W_x is of the order $1/N^2$ and that uniformly in x , $W_{x+1} - W_x$ is of the order ϵ_N/N^2 where ϵ_N vanishes to 0 as N goes to infinity. Thus the identity (4.1) implies

$$\frac{d\mathbb{P}_{\gamma,0,H}^N}{d\mathbb{P}_\gamma^N} = \exp \left(a \sum_{x=-N}^{N-1} W_x Q_T^N(x) + a o_\phi(N) \right),$$

where H is such that $H((x+1)/N) - H(x/N) = W_x$. This function is a mean one martingale and (4.18) leads to

$$\frac{1}{N} \log \mathbb{P}_\gamma^N (V_T(\rho^N, Q^N, K^N, \phi) > \delta) \leq -a\delta + a o_\phi(1).$$

Letting N and then a go to ∞ concludes the proof of the lemma. \square

Let (ϕ_k) be a dense sequence of functions in $C_0^2([-1, 1])$ and define

$$\mathcal{A}_n^\delta = \{(\rho, Q, K) \in \mathcal{E} \text{ such that } \forall k \leq n, \quad |V_T(\rho, Q, K, \phi_k)| \leq \delta\}.$$

The previous lemma gives that for all n and $\delta > 0$,

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N [(\rho^N, Q^N, K^N) \in F] \leq - \inf_{(\rho, Q, K) \in F \cap \mathcal{A}_n^\delta} J(\rho, Q, K).$$

This is true for any n and (ϕ_n) is dense. So letting δ go to 0, we can define $I_0 = +\infty$ for the trajectories which do not satisfy the relation (2.10).

Energy condition.

Lemma 4.4. *For any smooth $\varphi : (0, T) \times [-1, 1] \rightarrow \mathbb{R}$. There is a constant c_0 such that*

$$(4.19) \quad \forall k > 0, \quad \limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N \left(\mathcal{Q}_\varphi(\rho^N) \geq k \right) \leq -k + c_0 T,$$

where \mathcal{Q}_φ is defined for some suitable constant c

$$(4.20) \quad \begin{aligned} \mathcal{Q}_\varphi(\rho) &= \int_0^T dt \int_{-1}^1 dx \rho(t, x) \nabla \varphi(t, x) - \int_0^T dt (\bar{\rho}_+ \varphi(t, 1) - \bar{\rho}_- \varphi(t, -1)) \\ &\quad - \frac{1}{2c} \int_0^T dt \int_{-1}^1 dx \varphi(t, x)^2. \end{aligned}$$

The proof of this Lemma follows from [BLM, FLM] and therefore is omitted. By considering a dense sequence of functions $\{\varphi_k\}$, one can deduce from (4.19) [FLM] that the large deviations are infinite for the trajectories ρ such that $\sup_\varphi \mathcal{Q}_\varphi(\rho) = +\infty$. Note that if ρ is such that $\sup_\varphi \mathcal{Q}_\varphi(\rho) < +\infty$, then ρ is in $\mathbb{L}^2([0, T], \mathbb{H}([-1, 1]))$ and Riesz representation theorem implies that there exists $\nabla \rho$ such that for any $\varphi : (0, T) \times [-1, 1] \rightarrow \mathbb{R}$.

$$(4.21) \quad \int_0^T dt \int_{-1}^1 dx \rho(t, x) \nabla \varphi(t, x) - \{\bar{\rho}_+ \varphi(t, 1) - \bar{\rho}_- \varphi(t, -1)\} = \int_0^T dt \int_{-1}^1 dx \nabla \rho(t, x) \varphi(t, x).$$

5. LARGE DEVIATIONS LOWER BOUND

The derivation of the lower bound is split into two parts following the general scheme for hydrodynamic large deviations [KOV, KL]. First we derive the lower bound for regular trajectories. Then under assumptions **(L1, L2)**, we prove that general trajectories can be approximated by regular trajectories. A key feature in this approximation procedure is that the contribution of both currents decouple

$$(5.1) \quad I_0(\rho, Q, K) = I_1(\rho, Q) + I_2(\rho, K) = \sup_{H \in \mathcal{C}^{1,2}} J_H^1(\rho, Q) + \sup_{G \in \mathcal{C}^{1,0}} J_G^2(\rho, K),$$

where J_H^1 and J_G^2 were defined in (2.8) and (2.9). This simplifies some steps in the approximation as both functionals can be analyzed independently contrary to the study in

[JLV]. Assumption **(L1)** provides some convexity properties of the large deviation functional which simplify the proof. We follow closely some arguments of [KL, BDGJL2]. Thus we will sketch the main steps of the proofs and only detail the new aspects related to the non-conservative currents.

5.1. Lower bound for regular trajectories. In this section we derive Theorem 2.4. Suppose ρ, Q, K are regular in time and space and that for any time t and x , $\rho(t, x)$ is bounded away from 0 and 1. To the trajectory (ρ, Q, K) , one can associate the functions $G(t, x)$ and $H(t, x)$ satisfying

$$(5.2) \quad \begin{cases} \partial_t Q(t, x) &= -\frac{1}{2} \nabla \rho(t, x) + \sigma(\rho(t, x)) \nabla H(t, x), \\ \partial_t K(t, x) &= C(\rho(t, x)) e^{G(t, x)} - A(\rho(t, x)) e^{-G(t, x)}. \end{cases}$$

The pointwise existence of G comes from the fact that the polynomial $CX^2 - \dot{K}X - A$ has one positive root for positive C and A . Furthermore H is well defined as long as $\sigma(\rho) = \rho(1 - \rho) \neq 0$. We choose $H(t, -1) = 0$ and the value of H at $x = 1$ is imposed by equation (5.2), contrary to the case of density large deviations where H is equal to 0 at both boundaries (see section 6).

Lemma 5.1. *For regular trajectories (ρ, Q, K) the functional I_0 (2.12) is given by*

$$(5.3) \quad I_0(\rho, Q, K) = J_{G, H}(\rho, Q, K),$$

with G and H as in (5.2). This expression coincides with the explicit form of the functional (2.13).

Proof. As (ρ, Q, K) is regular in time, $J_{\tilde{G}, \tilde{H}}$ can be rewritten after integration by parts as

$$(5.4) \quad \begin{aligned} J_{\tilde{G}, \tilde{H}}(\rho, Q, K) &= \int_0^T \left\langle \left[\partial_s Q_s + \frac{1}{2} \nabla \rho_s - \frac{1}{2} \sigma(\rho_s) \nabla \tilde{H}_s \right] \nabla \tilde{H}_s \right\rangle ds \\ &+ \int_0^T \left\langle \partial_s K_s \tilde{G}_s - C(\rho_s)(e^{\tilde{G}_s} - 1) - A(\rho_s)(e^{-\tilde{G}_s} - 1) \right\rangle ds, \end{aligned}$$

for any (\tilde{G}, \tilde{H}) smooth functions.

For G and H given by (5.2), we are going to check that

$$(5.5) \quad J_{G, H}(\rho, Q, K) \geq \sup_{\tilde{G}, \tilde{H}} J_{\tilde{G}, \tilde{H}}(\rho, Q, K).$$

Indeed, take $\tilde{H} = H + F$ and any \tilde{G} , then it is easy to see that

$$J_{\tilde{G}, \tilde{H}}(\rho, Q, K) = J_{\tilde{G}, H}(\rho, Q, K) - \frac{1}{2} \int_0^T ds \langle \sigma(\rho_s) |\nabla F_s|^2 \rangle \leq J_{\tilde{G}, H}(\rho, Q, K).$$

Moreover, if $\tilde{G} = G + F$ and \tilde{H} any regular function, then from the expression of J_G^2 (2.9) and the identity (5.2) we get

$$\begin{aligned} J_{\tilde{G}, \tilde{H}}(\rho, Q, K) &= J_{G, \tilde{H}}(\rho, Q, K) + \int_0^T ds \langle C(\rho_s) e^{G_s} (1 + F_s - e^{F_s}) \rangle \\ &\quad + \int_0^T ds \langle A(\rho_s) e^{-G_s} (1 - F_s - e^{-F_s}) \rangle \leq J_{G, \tilde{H}}(\rho, Q, K), \end{aligned}$$

where we used that $\exp(x) - x - 1 \geq 0$ for any $x \in \mathbb{R}$. This completes the proof of (5.5).

Finally for G and H given by (5.2), one can rewrite (5.4)

$$\begin{aligned} J_{G, H}(\rho, Q, K) &= \frac{1}{2} \int_0^T \left\langle \frac{1}{\sigma(\rho_s)} (\partial_s Q_s + \frac{1}{2} \nabla \rho_s)^2 \right\rangle ds + \int_0^T \langle C(\rho_s) (1 - e^{G_s} + G_s e^{G_s}) \rangle ds \\ &\quad + \int_0^T \langle A(\rho_s) (1 - e^{-G_s} - G_s e^{-G_s}) \rangle ds. \end{aligned}$$

This leads to the explicit form of the functional (2.13). \square

Remark 5.2. For G and H as in (5.2), then the conservation equation $\partial_t \rho = \partial_t K - \nabla \partial_t Q$ implies that ρ obeys the hydrodynamic limit (3.2) of the modified dynamics $\mathbb{P}_{G, H}^N$ (3.1).

We introduce now the set of regular trajectories

Definition 5.3. Denote by \mathcal{S} the set of trajectories (ρ, Q, K) satisfying $I(\rho, Q, K) < \infty$ and such that

- ρ is bounded away from 0 and 1: there is $\varepsilon > 0$ such that $\varepsilon < \rho(t, x) < 1 - \varepsilon$ for any (t, x) in $[0, T] \times [-1, 1]$.
- There exists two regular functions G and H such that (ρ, Q, K) is a weak solution (in the sense of (7.1)) of

$$(5.6) \quad \begin{cases} \partial_t \rho(t, x) = \frac{1}{2} \Delta \rho(t, x) - \nabla (\sigma(\rho(t, x)) \nabla H(t, x)) + C(\rho(t, x)) e^{G(t, x)} - A(\rho(t, x)) e^{-G(t, x)}, \\ \rho(t, \pm 1) = \bar{\rho}_{\pm}, \quad \rho(t = 0, x) = \gamma(x), \end{cases}$$

- For any smooth test function φ in $C^{1,1}([0, T] \times [-1, 1])$

$$(5.7) \quad \begin{aligned} \langle Q_T \varphi_T \rangle - \int_0^T \left\langle Q_s \frac{d}{ds} \varphi_s \right\rangle ds &= -\frac{1}{2} \bar{\rho}_+ \int_0^T \varphi(s, 1) ds + \frac{1}{2} \bar{\rho}_- \int_0^T \varphi(s, -1) ds \\ &\quad + \frac{1}{2} \int_0^T \langle \rho_s \nabla \varphi_s \rangle ds + \int_0^T \langle \sigma(\rho_s) \nabla H_s \varphi_s \rangle ds, \end{aligned}$$

$$(5.8) \quad \langle K_T \varphi_T \rangle - \int_0^T \left\langle K_s \frac{d}{ds} \varphi_s \right\rangle ds = \int_0^T ds \langle [C(\rho_s) e^{G_s} - A(\rho_s) e^{-G_s}] \varphi_s \rangle.$$

The same argument as in Lemma 5.1 implies that for any trajectory (ρ, Q, K) in \mathcal{S}

$$\begin{aligned}
(5.9) \quad I_0(\rho, Q, K) &= J_{G,H}(\rho, Q, K) \\
&= \frac{1}{2} \int_0^T \langle \sigma(\rho_s) |\nabla H_s|^2 \rangle ds + \int_0^T \langle C(\rho_s) (1 - e^{G_s} + G_s e^{G_s}) \rangle ds \\
&\quad + \int_0^T \langle A(\rho_s) (1 - e^{-G_s} - G_s e^{-G_s}) \rangle ds.
\end{aligned}$$

We prove now the lower bound for trajectories in \mathcal{S} . Theorem 2.4 is a direct consequence of relation (5.9) and of the following Proposition.

Proposition 5.4. *For all open set $\mathcal{O} \in \mathcal{E}$,*

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N [(\rho^N, Q^N, K^N) \in \mathcal{O}] \geq - \inf_{(\rho, Q, K) \in \mathcal{O} \cap \mathcal{S}} I(\rho, Q, K).$$

Proof. Let (ρ, Q, K) be in $\mathcal{O} \cap \mathcal{S}$ and satisfying $I(\rho, Q, K) < \infty$. There is regular (G, H) for which ρ is a weak solution of (5.6). Thanks to Lemma 3.1, there is $\epsilon > 0$ such that the trajectories concentrate in the set $B_{\delta, \epsilon}$ (4.3)

$$(5.10) \quad \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_{\gamma, G, H}^N (B_{\delta, \epsilon}^c) \leq -1.$$

The function $J_{G, H, \epsilon}$ defined in (4.4) is continuous on \mathcal{E} . Moreover, since ρ_0 is bounded away from 0 and 1, the function $\varphi \rightarrow h_{\gamma, \rho_0}(\varphi(0, \cdot))$ is continuous on $D([0, T], \mathcal{M}_0)$. Let $\mathcal{V} \subset \mathcal{O}$ be an open neighborhood of (ρ, Q, K) such that

$$\begin{aligned}
\forall (\varphi, U, L) \in \mathcal{V}, \quad & |J_{G, H, \epsilon}(\varphi, U, L) - J_{G, H, \epsilon}(\rho, Q, K)| < \delta, \\
& |h_{\gamma, \rho_0}(\varphi_0) - h_{\gamma, \rho_0}(\rho_0)| = |h_{\gamma, \rho_0}(\varphi_0) - h_\gamma(\rho_0)| < \delta.
\end{aligned}$$

Using the change of measure (4.1)

$$\begin{aligned}
\mathbb{P}_\gamma^N(\mathcal{O}) &\geq \mathbb{P}_\gamma^N(\mathcal{V}) \geq \mathbb{P}_\gamma^N(\mathcal{V} \cap B_{\delta, \epsilon}) \geq \mathbb{P}_{\rho_0, G, H}^N \left(\frac{d\mathbb{P}_\gamma^N}{d\mathbb{P}_{\rho_0, G, H}^N} \mathbf{1}_{\mathcal{V} \cap B_{\delta, \epsilon}} \right) \\
&\geq \exp \left\{ -N(J_{G, H, \epsilon}(\rho, Q, K) + h_\gamma(\rho_0) + O_{G, H}(N^{-1}, \epsilon, \delta)) \right\} \mathbb{P}_{\rho_0, G, H}^N(\mathcal{V} \cap B_{\delta, \epsilon}).
\end{aligned}$$

Thanks to (5.10) and the hydrodynamical limit for the perturbed process (3.2)

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\rho_0, G, H}^N(\mathcal{V} \cap B_{\delta, \epsilon}) = 1,$$

where we used the uniqueness of the weak solution of (5.6) (see the Appendix) to conclude that the probability of \mathcal{V} converges to 1. This leads to

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_\gamma^N(\mathcal{O}) \geq -J_{G, H, \epsilon}(\rho, Q, K) - h_{\gamma, \rho_0}(\rho_0) + O_{G, H}(\delta, \epsilon).$$

Letting $\epsilon \downarrow 0$ and $\delta \downarrow 0$, and since (G, H) satisfy (5.6), the Proposition is completed. \square

5.2. Approximation for general trajectories. To complete Theorem 2.6 for general open sets, it remains to prove that

Lemma 5.5. *We assume (L1, L2). For any (ρ, Q, K) such that $I(\rho, Q, K) < \infty$ there is a sequence $(\rho^{(n)}, Q^{(n)}, K^{(n)})$ in \mathcal{S} converging weakly to (ρ, Q, K) such that*

$$(5.11) \quad I(\rho, Q, K) = \lim_{n \rightarrow \infty} I(\rho^{(n)}, Q^{(n)}, K^{(n)}).$$

Lemma 5.5 implies that

$$(5.12) \quad \inf_{(\rho, Q, K) \in \mathcal{O} \cap \mathcal{S}} I(\rho, Q, K) = \inf_{(\rho, Q, K) \in \mathcal{O}} I(\rho, Q, K).$$

Combining this identity and Proposition 5.4 proves Theorem 2.6.

5.2.1. Bounding the density away from 0 and 1. We first approximate the density by trajectories bounded away from 0 and 1.

Lemma 5.6. *Let $P = (\rho, Q, K)$ be a path such that $I(P) < +\infty$. There is $P_\delta = (\rho_\delta, Q_\delta, K_\delta)$ with density ρ_δ uniformly bounded away from 0 and 1 which converges to P and such that*

$$I(P) = \lim_{\delta \rightarrow 0} I(P_\delta).$$

Proof. Using Assumption (L1), we first establish a property of the functional I_0 . We use the decomposition (5.1) of I_0 . I_1 is a convex functional of (ρ, Q) as it is the supremum of J_H^1 which are convex functionals of (ρ, Q) (we used that $\sigma(\rho) = \rho(1 - \rho)$ is concave). I_2 is not convex, but we use a trick introduced in [JLV] and decompose I_2 as

$$I_2(\rho, K) = \sup_{G \in C^{1,0}} \tilde{J}_G^2(\rho, K) + \int_0^T dt \langle C(\rho_t) + A(\rho_t) \rangle.$$

Since A and C are concave, $\sup_G \tilde{J}_G^2(\rho, K)$ is a convex functional of (ρ, K) . Thus the large deviation functional can be decomposed into two terms

$$(5.13) \quad I_0(\rho, Q, K) = \tilde{I}_0(\rho, Q, K) + \int_0^T dt \langle C(\rho_t) + A(\rho_t) \rangle,$$

and \tilde{I}_0 is a convex functional of (ρ, Q, K) . We deduce that \tilde{I}_0 is lower semi-continuous for the weak topology.

We turn now to the approximation procedure. Let $\bar{P} = (\varphi, U, M)$ be the solution of the hydrodynamic equation (2.6)

$$\begin{cases} \partial_t \varphi(t, x) &= \frac{1}{2} \Delta \varphi(t, x) + C(\varphi(t, x)) - A(\varphi(t, x)) \\ \partial_t U(t, x) &= -\frac{1}{2} \nabla \varphi(t, x) \\ \partial_t M(t, x) &= C(\varphi(t, x)) - A(\varphi(t, x)) \end{cases}$$

with boundary conditions

$$\varphi_0(0, x) = \gamma(x), \quad U(0, x) = 0, \quad M(0, x) = 0, \quad \varphi(t, -1) = \bar{\rho}_-, \quad \varphi(t, 1) = \bar{\rho}_+.$$

By construction $I(\bar{P}) = 0$. We set $P_\delta = (1 - \delta)P + \delta\bar{P}$ which has a density bounded away from 0 and 1. As $\tilde{I}_0(\bar{P}) < \infty$, the convexity implies that $\tilde{I}_0(P_\delta) \leq (1 - \delta)\tilde{I}_0(P) + \delta\tilde{I}_0(\bar{P})$ so that

$$\limsup_{\delta \rightarrow 0} \tilde{I}_0(P_\delta) \leq \tilde{I}_0(P).$$

As P_δ weakly converges to P , the lower semi-continuity of \tilde{I}_0 implies

$$\tilde{I}_0(P) \leq \liminf_{\delta \rightarrow 0} \tilde{I}_0(P_\delta).$$

Thus $\tilde{I}_0(P_\delta)$ converges to $\tilde{I}_0(P)$. Finally h_γ and $\rho \rightarrow \int_0^T dt \langle C(\rho_t) + A(\rho_t) \rangle$ are continuous for $\|\cdot\|_\infty$. This completes the Lemma. \square

5.2.2. *Time regularisation.* We will prove

Lemma 5.7. *For any path $P = (\rho, Q, K)$ such that $I(P) < \infty$, there is a sequence regular in time $P_\epsilon = (\rho_\epsilon, Q_\epsilon, K_\epsilon)$ converging weakly to (ρ, Q, K) such that $I(P_\epsilon)$ converges to $I(P)$.*

In the following, only the regularity of K is needed in order to construct a drift G adapted to the non-conservative current (5.15). However, the regularizing sequence has to satisfy the conservation law (2.10) so that ρ, Q, K will be approximated simultaneously.

Proof. The proof is based on a time convolution and follows similar steps as in Lemma 5.6. We just recall the salient features of the proof (see [KL] for further details).

Let ψ_ϵ be a C^∞ approximation of unity such that $\psi_\epsilon = 0$ outside $[0, \epsilon]$ and $\int_0^\epsilon \psi_\epsilon = 1$. To take the convolution product of (ρ, Q, K) with ψ_ϵ , we have to extend the path (ρ, Q, K) beyond the time T . Let r be the solution of the hydrodynamic equation (2.6)

$$\partial_s r_s = \frac{1}{2} \Delta r_s + C(r_s) - A(r_s), \quad \text{with} \quad r_0 = \rho_T.$$

We set for $s > T$

$$\rho_s = r_{s-T}, \quad Q_s = Q_T - \frac{1}{2} \int_T^s du \nabla \rho_u, \quad K_s = K_T + \int_T^s du [C(\rho_u) - A(\rho_u)].$$

Now we define

$$(5.14) \quad \begin{cases} \rho_\epsilon(t, x) = \rho * \psi_\epsilon(t, x), \\ Q_\epsilon(t, x) = Q * \psi_\epsilon(t, x) - Q * \psi_\epsilon(0, x), \\ K_\epsilon(t, x) = K * \psi_\epsilon(t, x) - K * \psi_\epsilon(0, x). \end{cases}$$

The path $(\rho_\epsilon, Q_\epsilon, K_\epsilon)$ satisfies the relation (2.10) and has initial currents equal to 0. We use the decomposition (5.1) and approximate independently the functionals I_1 and I_2 by using convexity properties deduced from Assumption **(L1)**. \square

5.2.3. *Non regular drifts.* Thanks to Lemmas 5.6 and 5.7, it is enough to consider a trajectory (ρ, Q, K) regular in time with a density uniformly bounded away from 0 and 1. We are going to associate to (ρ, Q, K) the drifts H, G as for the trajectories in \mathcal{S} introduced in Definition 5.3. Note that the drifts H, G can be non regular in space.

As $\partial_s K$ exists, the drift G is defined (as for the regular trajectories) as the solution of

$$(5.15) \quad \forall x \in [-1, 1], \quad \partial_s K(s, x) = C(\rho(s, x))e^{G(s, x)} - A(\rho(s, x))e^{-G(s, x)}.$$

Note that K, G solve the equation (5.8) and as in the regular case (5.9)

$$(5.16) \quad I_2(\rho, K) = \int_0^T \langle C(\rho_s) (1 - e^{G_s} + G_s e^{G_s}) \rangle + \langle A(\rho_s) (1 - e^{-G_s} - G_s e^{-G_s}) \rangle ds.$$

The functional I_1 is the same as the functional of the SSEP (without Glauber rates). Thus the drift of the conservative dynamics can be approximated thanks to the Riesz representation theorem as in [KL, BDGJL2]. For (ρ, Q) such that $I_1(\rho, Q) < \infty$, there is H in $\mathbb{L}^2([0, T], \mathbb{H}_1([-1, 1]))$ such that

$$(5.17) \quad I_1(\rho, Q) = \int_0^T \int_{-1}^1 \frac{\sigma(\rho(s, x))}{2} (\nabla H(s, x))^2 dx ds.$$

Moreover (5.7) holds. Note that the time regularity is not needed to derive (5.17).

Finally, we check that ρ is a weak solution of (7.1) with the drifts G, H . As the large deviation functional is finite, ρ belongs to \mathcal{A} so that $\mathcal{Q}(\rho)$ is finite and (7.2) is satisfied. Combining the conservation law (2.10) with (5.7) and (5.8) shows that (7.3) holds.

5.2.4. *Approximation paths in \mathcal{S} .* We approximate now H, G by regularized drifts H_n, G_n and prove that the associated sequence of paths $(\rho^{(n)}, Q^{(n)}, K^{(n)})$ in \mathcal{S} converges to (ρ, Q, K) .

Let H_n be a sequence of C^∞ functions converging to H in $\mathbb{L}^2([0, T], \mathbb{H}_1([-1, 1]))$. Since C and A satisfy assumption **(L1)** and ρ is uniformly bounded away from 0 and 1 (Lemma 5.6), C and A are either uniformly bounded away from 0, or uniformly equal to 0. We will consider only the case $A > 0, C > 0$ as the other case follows in the same way. From (5.16), we get that $|G|e^{|G|} \in \mathbb{L}^1([0, T] \times [-1, 1])$, so that there is G_n such that

$$(5.18) \quad \int_0^T \int_{-1}^1 |G_n e^{\pm G_n} - G e^{\pm G}| \xrightarrow{n \rightarrow \infty} 0 \quad \text{and} \quad \int_0^T \int_{-1}^1 |e^{\pm G_n} - e^{\pm G}| \xrightarrow{n \rightarrow \infty} 0.$$

Let $(\rho^{(n)}, Q^{(n)}, K^{(n)})$ be the sequence of paths in \mathcal{S} associated to the regular drifts H_n, G_n . In particular $\rho^{(n)}$ is the weak solution of

$$(5.19) \quad \begin{cases} \partial_t \rho^{(n)} = \frac{1}{2} \Delta \rho^{(n)} - \nabla(\sigma(\rho^{(n)}) \nabla H_n) + C(\rho^{(n)})e^{G_n} - A(\rho^{(n)})e^{-G_n} \\ \rho^{(n)}(t, \pm 1) = \bar{\rho}_\pm, \quad \rho^{(n)}(0, \cdot) = \rho(0, \cdot), \end{cases}$$

and $Q^{(n)}$ and $K^{(n)}$ are defined as

$$(5.20) \quad \begin{aligned} Q^{(n)}(t, x) &= \int_0^t \left(-\frac{1}{2} \nabla \rho^{(n)}(s, x) + \sigma(\rho^{(n)}(s, x)) \nabla H_n(s, x) \right) ds, \\ K^{(n)}(t, x) &= \int_0^t \left(C(\rho^{(n)}(s, x)) e^{G_n(s, x)} - A(\rho^{(n)}(s, x)) e^{-G_n(s, x)} \right) ds. \end{aligned}$$

From energy estimates (using similar bounds as in the Appendix (7.8)) we get

$$(5.21) \quad \int_0^T dt \int_{-1}^1 dx (\nabla \rho^{(n)})^2 \leq C,$$

where C is a constant independent of n . As $\rho^{(n)}$ and $\nabla \rho^{(n)}$ are bounded sequences in \mathbb{L}^2 , they are tight in the weak topology. We want to check the uniqueness of the limiting points. Let R be a limit point of $\rho^{(n)}$. We will prove that R is a weak solution of (7.1). From Section 5.2.3, we know that ρ is also a weak solution of (7.1) so that the uniqueness of the weak solutions (see the Appendix) will imply that $\rho = R$.

We want to take the limit in the weak formulation of (5.19). By construction, for any smooth function φ on $[-1, 1]$, we have

$$\int_0^t ds \int_{-1}^1 dx \rho^{(n)}(s, x) \nabla \varphi(x) = [\bar{\rho}_+ \varphi(1) - \bar{\rho}_- \varphi(-1)] - \int_0^t ds \int_{-1}^1 dx \nabla \rho^{(n)}(s, x) \varphi(x).$$

By weak convergence of the subsequence, one has

$$\int_0^t ds \int_{-1}^1 dx R(s, x) \nabla \varphi(x) = [\bar{\rho}_+ \varphi(1) - \bar{\rho}_- \varphi(-1)] - \int_0^t ds \int_{-1}^1 dx \nabla R(s, x) \varphi(x).$$

Furthermore (5.21) implies that the limit ∇R is also in \mathbb{L}^2 . It remains to take the limit in n in the equation

$$(5.22) \quad \begin{aligned} \int_{-1}^1 \rho^{(n)}(t) \varphi &= \int_{-1}^1 \gamma \varphi - \frac{1}{2} \int_0^t \int_{-1}^1 \nabla \rho^{(n)} \nabla \varphi + \int_0^t \int_{-1}^1 \nabla \varphi \sigma(\rho^{(n)}) \nabla H_n \\ &\quad + \int_0^t \int_{-1}^1 C(\rho^{(n)}) e^{G_n} \varphi - \int_0^t \int_{-1}^1 A(\rho^{(n)}) e^{-G_n} \varphi. \end{aligned}$$

The difficulty is to treat the non linear terms. We proceed term by term and start with the non-linearity C in (5.22) (the term in A can be controlled in the same way). By (5.18), $\exp(G_n)$ converges to $\exp(G)$ in \mathbb{L}^1 thus it is enough to check that

$$(5.23) \quad \lim_{n \rightarrow \infty} \int_0^t \int_{-1}^1 C(\rho^{(n)}) e^{G_n} \varphi = \int_0^t \int_{-1}^1 C(R) e^G \varphi.$$

We write

$$(5.24) \quad \int_0^t \int_{-1}^1 (C(\rho^{(n)}) - C(R))\varphi e^G = \int_0^t \int_{-1}^1 (C(\rho^{(n)}) - C(\rho^{(n)} * \iota_\delta))\varphi e^G \\ + \int_0^t \int_{-1}^1 (C(\rho^{(n)} * \iota_\delta) - C(R * \iota_\delta))\varphi e^G + \int_0^t \int_{-1}^1 (C(R * \iota_\delta) - C(R))\varphi e^G.$$

As δ goes to 0, $R * \iota_\delta$ converges to R . The convergence in the weak topology implies that $\rho^{(n)} * \iota_\delta$ converges a.s. to $R * \iota_\delta$ when n goes to infinity. Suppose that

$$(5.25) \quad \limsup_{\delta \rightarrow 0} \sup_n \left| \int_0^t \int_{-1}^1 (C(\rho^{(n)}) - C(\rho^{(n)} * \iota_\delta))e^G \varphi \right| = 0,$$

then choosing δ small and then n large, the convergence in (5.23) follows.

To derive (5.25), we first note that for $x \in]-1 + \delta, 1 - \delta[$

$$(5.26) \quad \rho^{(n)} * \iota_\delta(x) - \rho^{(n)}(x) \\ = \int_{-1}^1 dy \iota_\delta(x - y)[\rho^{(n)}(y) - \rho^{(n)}(x)] = \int_{-1}^1 dy \iota_\delta(x - y) \int_x^y du \partial_u \rho^{(n)}(u) \\ \leq \sqrt{\int_{-1}^1 du (\partial_u \rho^{(n)}(u))^2} \int_{-1}^1 dy \iota_\delta(x - y) \sqrt{|x - y|} \leq 2\sqrt{\delta} \sqrt{\int_{-1}^1 du (\partial_u \rho^{(n)}(u))^2}.$$

Using the uniform bound (5.21) there is a constant C independent of n such that

$$(5.27) \quad \int_0^t \int_{-1}^1 (\rho^{(n)} * \iota_\delta - \rho^{(n)})^2 \leq C\delta.$$

It remains to check (5.25). Let \mathbf{m} be the uniform measure on $[0, t] \times [-1, 1]$ given by $\frac{1}{2t} ds dx$. Given $\psi(s, x)$ a non-negative function on $[0, t] \times [-1, 1]$ such that $\mathbf{m}(\psi) = 1$, then the entropy inequality implies that for any function φ

$$(5.28) \quad \mathbf{m}(\varphi(s, x)\psi(s, x)) \leq \mathbf{m}(\psi(s, x) \log(\psi(s, x))) + \log \mathbf{m}(\exp(\varphi(s, x))).$$

Since A and C are positive then $\mathbf{m}(e^{|G|}|G|)$ is bounded and (5.28) implies

$$\mathbf{m} \left(|C(\rho^{(n)} * \iota_\delta) - C(\rho^{(n)})| |\varphi| \frac{e^{|G|}}{\mathbf{m}(e^{|G|})} \right) \\ \leq \varepsilon_\delta \mathbf{m} \left(\frac{e^{|G|}}{\mathbf{m}(e^{|G|})} \log \frac{e^{|G|}}{\mathbf{m}(e^{|G|})} \right) + \varepsilon_\delta \log \mathbf{m} \left(\exp \left(\frac{1}{\varepsilon_\delta} |C(\rho^{(n)} * \iota_\delta) - C(\rho^{(n)})| |\varphi| \right) \right),$$

where ε_δ vanishes (in a suitable way to be determined later) as δ goes to 0. Recall that

$$\forall x \in [0, M], \quad \exp(x) \leq 1 + x + \exp(M)x^2 \quad \text{and} \quad \log(x) \leq x - 1.$$

As $|C(\rho^{(n)} * \iota_\delta) - C(\rho^{(n)})||\varphi| \leq 2\|C\|_\infty\|\varphi\|_\infty$, the previous inequalities imply

$$\begin{aligned} & \log \mathbf{m} \left(\exp \left(\frac{1}{\varepsilon_\delta} |C(\rho^{(n)} * \iota_\delta) - C(\rho^{(n)})||\varphi| \right) \right) \\ & \leq \frac{1}{\varepsilon_\delta^2} \exp \left(\frac{\|C\|_\infty\|\varphi\|_\infty}{\varepsilon_\delta} \right) \mathbf{m} \left(|C(\rho^{(n)} * \iota_\delta) - C(\rho^{(n)})||\varphi| + \left(|C(\rho^{(n)} * \iota_\delta) - C(\rho^{(n)})||\varphi| \right)^2 \right), \end{aligned}$$

We choose ε_δ such that

$$\lim_{\delta \rightarrow 0} \varepsilon_\delta = 0, \quad \lim_{\delta \rightarrow 0} \frac{\sqrt{\delta}}{\varepsilon_\delta} \exp \left(\frac{\|C\|_\infty\|\varphi\|_\infty}{\varepsilon_\delta} \right) = 0.$$

Combining the previous inequality and (5.27), we conclude that (5.25) holds.

We turn now to the non-linearity σ . Using the decomposition

$$\begin{aligned} & \int_0^t \int_{-1}^1 [\sigma(\rho^{(n)}) - \sigma(R)] \nabla \varphi_s \nabla H_s = \int_0^t \int_{-1}^1 [\sigma(\rho^{(n)}) - \sigma(\rho^{(n)} * \iota_\delta)] \nabla \varphi_s \nabla H_s \\ & + \int_0^t \int_{-1}^1 [\sigma(\rho^{(n)} * \iota_\delta) - \sigma(R * \iota_\delta)] \nabla \varphi_s \nabla H_s + \int_0^t \int_{-1}^1 [\sigma(R * \iota_\delta) - \sigma(R)] \nabla \varphi_s \nabla H_s \end{aligned}$$

the last two terms vanish in the limit as in (5.24). The first term can be controlled thanks to the uniform bound (5.27) and a Cauchy-Schwartz estimate as $\nabla \varphi \nabla H$ belongs to \mathbb{L}^2 . This concludes the convergence of (5.22).

Following the same proof, (5.20) implies that $(Q^{(n)}, K^{(n)})$ converges weakly to (Q, K) .

5.2.5. I -convergence. We finally complete the proof of Lemma 5.5. Let $(\rho^{(n)}, Q^{(n)}, K^{(n)})$ be the regularizing sequence defined in (5.19). We start by proving

$$(5.29) \quad \limsup_{n \rightarrow \infty} I_0(\rho^{(n)}, Q^{(n)}, K^{(n)}) \leq I_0(\rho, Q, K).$$

From the expression (5.17) of I_1 and the weak convergence of $\rho^{(n)}$ to ρ , we get

$$\begin{aligned} I_1(\rho, Q) &= \int_0^T \int_{-1}^1 \frac{\sigma(\rho(s, x))}{2} \nabla H(s, x)^2 dx ds = \lim_{\delta \rightarrow 0} \int_0^T \int_{-1}^1 \frac{\sigma([\rho_s * \iota_\delta](x))}{2} \nabla H(s, x)^2 dx ds \\ &= \lim_{\delta \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{2} \int_0^T \int_{-1}^1 \nabla H(s, x)^2 \sigma([\rho_s^{(n)} * \iota_\delta](x)) dx ds \\ &\geq \lim_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{2} \int_0^T \int_{-1}^1 (\nabla H(s, x))^2 [\sigma(\rho_s^{(n)} * \iota_\delta(x))] dx ds, \end{aligned}$$

where we used the concavity of σ in the last inequality. Notice that

$$\begin{aligned} & \int_0^T \int_{-1}^1 (\nabla H(s, x))^2 [\sigma(\rho_s^{(n)} * \iota_\delta(x))] dx ds = \int_0^T \int_{-1}^1 (\nabla H(s, x))^2 \sigma(\rho_s^{(n)}(x)) dx ds \\ & + \int_0^T \int_{-1}^1 [(\nabla H_s)^2 * \iota_\delta(x) - (\nabla H(s, x))^2] \sigma(\rho_s^{(n)}(x)) dx ds. \end{aligned}$$

As σ is bounded and $(\nabla H)^2$ is integrable, the last term vanishes uniformly in n when δ goes to 0, so that

$$I_1(\rho, Q) \geq \limsup_{n \rightarrow \infty} \frac{1}{2} \int_0^T \int_{-1}^1 \nabla H(s, x)^2 \sigma(\rho_s^{(n)}(x)) dx ds.$$

As $\nabla H^{(n)}$ converges to ∇H in \mathbb{L}^2 and σ is bounded, we conclude that

$$(5.30) \quad I_1(\rho, Q) \geq \limsup_{n \rightarrow \infty} I_1(\rho^{(n)}, Q^{(n)}).$$

Similarly for I_2 , the concavity and boundedness of C and A imply that

$$I_2(\rho, K) \geq \limsup_{n \rightarrow \infty} \int_0^T \int_{-1}^1 C(\rho^{(n)}) (1 - e^G + Ge^G) + A(\rho^{(n)}) (1 - e^{-G} - Ge^{-G}).$$

Then we use the convergence properties on $G^{(n)}$ (5.18) and the fact that C and A are bounded to conclude that

$$(5.31) \quad I_2(\rho, K) \geq \limsup_{n \rightarrow \infty} I_2(\rho^{(n)}, K^{(n)})$$

Estimates (5.30) and (5.31) imply (5.29).

The converse inequality

$$(5.32) \quad \liminf_{n \rightarrow \infty} I_0(\rho^{(n)}, Q^{(n)}, K^{(n)}) \geq I_0(\rho, Q, K),$$

can be deduced from the weak convergence of $(\rho^{(n)}, Q^{(n)}, K^{(n)})$ to (ρ, Q, K) and from the decomposition (5.13): the limit follows for the term \tilde{I}_0 from its lower semi-continuity and the convergence of the second term $\int_0^T dt \langle C(\rho_t^{(n)}) + A(\rho_t^{(n)}) \rangle$ can be obtained as in (5.23).

6. THE DENSITY LARGE DEVIATIONS

In this section, we recover the density large deviation principle (first derived in [JLV]) by optimizing the functional I over the currents. We assume that the rates A, C satisfy assumptions **(L1, L2)**. The contraction principle [DZ] implies that the density large deviation functional is given by

$$(6.1) \quad \mathcal{F}(\rho) = \inf_{(Q, K)} I(\rho, Q, K),$$

where the infimum is taken over the currents (Q, K) . Using the approximation procedure of section 5.2, we will check that it is enough to consider regular density profiles and the modified functional

$$(6.2) \quad \hat{\mathcal{F}}(\rho) = \inf_{(Q, K)_{\text{reg}}} I(\rho, Q, K),$$

where the infimum is taken now over the regular currents (Q, K) .

Step 1. In this first step, the explicit solution of the variational problem (6.2) is computed for regular trajectories. The functional (6.2) can be rewritten as (see (5.9))

$$(6.3) \quad \hat{\mathcal{F}}(\rho) = \inf_{(G,H)} \left\{ \frac{1}{2} \int_0^T \langle \sigma(\rho_s) |\nabla H_s|^2 \rangle ds + \int_0^T \langle C(\rho_s) (1 - e^{G_s} + G_s e^{G_s}) \rangle ds \right. \\ \left. + \int_0^T \langle A(\rho_s) (1 - e^{-G_s} - G_s e^{-G_s}) \rangle ds \right\},$$

where G, H are smooth functions such that the conservation relation (2.10) holds

$$(6.4) \quad \partial_s \rho_s = \frac{1}{2} \Delta \rho_s - \partial_x (\sigma(\rho_s) \nabla H_s) + (C(\rho_s) e^{G_s} - A(\rho_s) e^{-G_s}).$$

We first check that the infimum is reached for functions H with boundary conditions $H(s, -1) = H(s, 1) = 0$ for any time s . Let f be a smooth function in $[0, T]$. Perturbing H into $H(s, x) + f(s) \int_0^x \frac{1}{\sigma(\rho(s, u))} du$, we see that the conservation law (6.4) is preserved (it simply amounts to adding a constant conservative current) and

$$\int_0^T \left\langle \sigma(\rho_s) \left| \nabla H_s + \frac{\alpha f(s)}{\sigma(\rho(s, x))} \right|^2 \right\rangle ds = \int_0^T \langle \sigma(\rho_s) |\nabla H_s|^2 \rangle ds \\ + 2 \int_0^T (H(s, 1) - H(s, -1)) f(s) + \int_0^T \left\langle \frac{(f(s))^2}{\sigma(\rho(s, x))} \right\rangle ds.$$

As H minimizes the integral, this implies that H vanishes at the boundaries.

Suppose that an extremum is reached at (H, G) and consider a perturbation with the new drifts $H + h$ and $G + g$. Then the constraint (6.4) implies that h, g satisfy the relation

$$(6.5) \quad \partial_x (\sigma(\rho_s) \nabla h_s) + g_s (C(\rho_s) e^{G_s} + A(\rho_s) e^{-G_s}) = 0.$$

A perturbation of (6.3) around the extremum H, G leads to

$$(6.6) \quad 0 = \int_0^T \langle \sigma(\rho_s) \nabla H_s \nabla h_s \rangle ds + \int_0^T \langle g_s G_s (C(\rho_s) e^{G_s} + A(\rho_s) e^{-G_s}) \rangle ds.$$

Since H vanishes at the boundaries, the relation (6.5) combined to (6.6) leads to

$$0 = \int_0^T \langle g_s [G_s - H_s] (C(\rho_s) e^{G_s} + A(\rho_s) e^{-G_s}) \rangle ds.$$

This holds for any g so that the extremum is such that $G = H$ with H determined by

$$(6.7) \quad \partial_t \rho = \frac{1}{2} \Delta \rho - \nabla (\sigma(\rho) \nabla H) + C(\rho) e^H - A(\rho) e^{-H}.$$

Thus if H satisfies (6.7) and $G = H$, then an extremum is reached for the corresponding currents (\hat{Q}, \hat{K}) . Since the functional $I(\rho, Q, K)$ is convex wrt (Q, K) (thanks to the

representation (2.8), (2.9)), the extremum (\hat{Q}, \hat{K}) has to be a global minimum, thus

$$\begin{aligned} \hat{\mathcal{F}}(\rho) &= \frac{1}{2} \int_0^T \langle \sigma(\rho_s) |\nabla H_s|^2 \rangle ds + \int_0^T \langle C(\rho_s) (1 - e^{H_s} + H_s e^{H_s}) \rangle ds \\ &\quad + \int_0^T \langle A(\rho_s) (1 - e^{-H_s} - H_s e^{-H_s}) \rangle ds. \end{aligned}$$

Step 2. To approximate \mathcal{F} (6.1) in terms of $\hat{\mathcal{F}}$ (6.2), we first check that the minimum is reached in (6.1). Consider a sequence (Q^n, K^n) which realizes the infimum. By the tightness argument (section 4.2) the sequence belongs to a compact set and therefore has a weak limit (Q^*, K^*) . From the lower semi-continuity of the functional $(Q, K) \rightarrow I(\rho, Q, K)$, this weak limit is a minimizer

$$(6.8) \quad \mathcal{F}(\rho) = I(\rho, Q^*, K^*).$$

Let $(\rho_\varepsilon, Q_\varepsilon^*, K_\varepsilon^*)$ be a regular sequence (as in (5.19)) converging to (ρ, Q^*, K^*) such that

$$I(\rho_\varepsilon, Q_\varepsilon^*, K_\varepsilon^*) \rightarrow I(\rho, Q^*, K^*).$$

As $\hat{\mathcal{F}}(\rho_\varepsilon) \leq I(\rho_\varepsilon, Q_\varepsilon^*, K_\varepsilon^*)$, one has

$$\limsup_{\varepsilon \rightarrow 0} \hat{\mathcal{F}}(\rho_\varepsilon) \leq \lim_{\varepsilon \rightarrow 0} I(\rho_\varepsilon, Q_\varepsilon^*, K_\varepsilon^*) = \mathcal{F}(\rho).$$

Using (6.7), there are regular $(\hat{Q}_\varepsilon, \hat{K}_\varepsilon)$ such that

$$\hat{\mathcal{F}}(\rho_\varepsilon) = I(\rho_\varepsilon, \hat{Q}_\varepsilon, \hat{K}_\varepsilon).$$

The sequence $(\rho_\varepsilon, \hat{Q}_\varepsilon, \hat{K}_\varepsilon)$ has a bounded large deviation cost and thus it belongs to a compact set. There is a subsequence such that $(\hat{Q}_\varepsilon, \hat{K}_\varepsilon)$ converges weakly to (\hat{Q}, \hat{K}) . One gets

$$\liminf_{\varepsilon \rightarrow 0} \hat{\mathcal{F}}(\rho_\varepsilon) \geq I(\rho, \hat{Q}, \hat{K}) \geq \mathcal{F}(\rho).$$

This limit follows from the decomposition (5.13). The term \tilde{I}_0 converges by lower semi-continuity and the convergence of the second term $\int_0^T dt \langle C(\rho_\varepsilon(t)) + A(\rho_\varepsilon(t)) \rangle$ can be obtained as in (5.23) (ρ_ε is just a subsequence extracted from $\rho^{(n)}$).

Combining both estimates, we deduce that \mathcal{F} can be approximated by $\hat{\mathcal{F}}$.

7. APPENDIX : UNIQUENESS OF THE WEAK SOLUTIONS

Let H be in $\mathbb{L}^2([0, T], \mathbb{H}_1([-1, 1]))$ and $|G| \exp(|G|)$ in $\mathbb{L}^1([0, T] \times]-1, 1[)$. Given an initial data γ , a weak solution of

$$(7.1) \quad \begin{cases} \partial_t \rho &= \frac{1}{2} \Delta \rho - \partial_x (\sigma(\rho) \partial_x H) + C(\rho) e^G - A(\rho) e^{-G} \\ \bar{\rho}(t, \pm 1) &= \bar{\rho}_\pm, \quad \bar{\rho}(0, x) = \gamma(x) \end{cases}$$

is defined as :

- The density ρ is in $\mathbb{L}^2([0, T], \mathbb{H}^1([-1, 1]))$, i.e. there is a function in $\mathbb{L}^2([0, T] \times [-1, 1])$ which will be denoted by $\nabla\rho$ such for every $t \in [0, T]$ and every function $\varphi \in C^1([-1, 1])$

$$(7.2) \quad \int_0^t ds \int_{-1}^1 dx \rho(s, x) \nabla\varphi(x) - \{\bar{\rho}_+\varphi(1) - \bar{\rho}_-\varphi(-1)\}t = \int_0^t ds \int_{-1}^1 dx \nabla\rho(s, x)\varphi(x),$$

where $\bar{\rho}_\pm$ are fixed boundary conditions.

- For every $t \in [0, T]$ and every function $\varphi \in C^1([-1, 1])$ vanishing at the boundaries,

$$(7.3) \quad \int_{-1}^1 dx \rho(t, x)\varphi(x) - \int_{-1}^1 dx \gamma(x)\varphi(x) = - \int_0^t ds \int_{-1}^1 dx \nabla\rho(s, x)\nabla\varphi(x) \\ + \int_0^t ds \int_{-1}^1 dx \sigma(\rho(s, x))\nabla H(s, x)\nabla\varphi(x) \\ + \int_0^t ds \int_{-1}^1 dx (C(\rho(s, x))e^{G(s, x)} - A(\rho(s, x))e^{-G(s, x)})\varphi(x).$$

The hydrodynamic limit (2.6) corresponds to $H = G = 0$.

In this Appendix, we derive the uniqueness of the weak solutions. The main technical difficulty comes from the fact that $G, \partial_x H$ are unbounded (see [E] for bounded drifts). Note that at this stage Assumption **(L1)** is irrelevant. We will rely on Assumption **(L2)** on A, C which can be interpreted as follows. Equation (7.1) in a strong form reads

$$\partial_t \rho(t, x) = \Delta\rho(t, x) - \partial_x(\sigma(\rho(t, x))\nabla H(t, x)) - V'_{G(t, x)}(\rho(t, x)),$$

where the reaction term is determined by the space-time dependent potential $V_{G(t, x)}$ with $V'_g(\rho) = -e^g C(\rho) + e^{-g} A(\rho)$. Assumption **(L2)** ensures that the potential $V_{G(t, x)}$ is convex uniformly in $G(t, x)$. Thus the reaction and the diffusion terms are both contractions and the solution will be unique. We adapt to our framework the argument of [LMS, FLM].

We consider two initial datas ρ_0^1, ρ_0^2 and the corresponding weak solutions ρ^1, ρ^2 . We are going to prove that the \mathbb{L}^1 -norm $\|\rho_t^1 - \rho_t^2\|_1$ decreases in time. For a given $\delta > 0$, we introduce the regularized absolute value

$$(7.4) \quad U_\delta(v) = \frac{v^2}{2\delta} 1_{\{|v| \leq \delta\}} + \left(|v| - \frac{\delta}{2}\right) 1_{\{|v| > \delta\}}.$$

Define

$$(7.5) \quad V_\delta = \left\{ (s, x) \in [0, T] \times [-1, 1], \quad \text{such that } |\rho^1(s, x) - \rho^2(s, x)| \leq \delta \right\}.$$

Step 1. We are going to check that for times $t < t'$

$$\begin{aligned}
(7.6) \quad & \int_{-1}^1 dx U_\delta \left(\rho^1(t', x) - \rho^2(t', x) \right) - \int_{-1}^1 dx U_\delta \left(\rho^1(t, x) - \rho^2(t, x) \right) \\
&= -\frac{1}{\delta} \int_t^{t'} \int_{-1}^1 ds dx 1_{\{(x,s) \in V_\delta\}} \nabla(\rho^1 - \rho^2)(s, x) \left\{ \nabla(\rho^1 - \rho^2)(s, x) + \bar{\sigma}(s, x) \nabla H(s, x) \right\} \\
&\quad + \frac{1}{\delta} \int_t^{t'} \int_{-1}^1 ds dx U'_\delta(\rho^1(s, x) - \rho^2(s, x)) \left\{ \bar{C}(s, x) e^{G(s, x)} - \bar{A}(s, x) e^{-G(s, x)} \right\}
\end{aligned}$$

where we set

$$\begin{aligned}
\bar{A}(t, x) &= A(\rho^1(t, x)) - A(\rho^2(t, x)), \quad \bar{C}(t, x) = C(\rho^1(t, x)) - C(\rho^2(t, x)), \\
\bar{\sigma}(t, x) &= \sigma(\rho^1(t, x)) - \sigma(\rho^2(t, x)).
\end{aligned}$$

To prove (7.6), we follow the regularization scheme introduced in [BLM] (see the proof of their Theorem 4.6). For $\varepsilon > 0$, denote by $R_\varepsilon^D : [-1, 1]^2 \rightarrow \mathbb{R}^+$ (resp R_ε^N) the resolvent of the Dirichlet Laplacian Δ_D (resp Neumann Δ_N)

$$R_\varepsilon^D = (\text{Id} - \varepsilon \Delta_D)^{-1}, \quad R_\varepsilon^N = (\text{Id} - \varepsilon \Delta_N)^{-1}.$$

The mollified trajectory is defined by

$$\rho^\varepsilon(t, x) = \bar{\rho}(x) + R_\varepsilon^D(\rho_t - \bar{\rho})(x),$$

where $\bar{\rho}$ stands for the linear profile between $\bar{\rho}_+$ and $\bar{\rho}_-$. Note that the resolvent of the Dirichlet Laplacian preserves the boundary conditions of the mollified trajectory.

As ρ is a weak solution of (7.3), one has

$$\begin{aligned}
\partial_t \rho^\varepsilon(t, x) &= \partial_t \int_{-1}^1 dy R_\varepsilon^D(x, y) (\rho(t, y) - \bar{\rho}(y)) \\
&= \int_{-1}^1 dy \partial_y R_\varepsilon^D(x, y) \left\{ -\partial_y(\rho(t, y) - \bar{\rho}(y)) + \sigma(\rho(t, y)) \nabla H(t, y) \right\} \\
&\quad + \int_{-1}^1 dy R_\varepsilon^D(x, y) (C(\rho(t, y)) e^{G(t, y)} - A(\rho(t, y)) e^{-G(t, y)}),
\end{aligned}$$

where we used the fact that $R_\varepsilon^D(x, 1) = R_\varepsilon^D(x, -1) = 0$ for any x .

From the relation $\partial_y R_\varepsilon^D(x, y) = -\partial_x R_\varepsilon^N(x, y)$, one has

$$\begin{aligned}
(7.7) \quad \partial_t \rho^\varepsilon(t, x) &= \partial_x \int_{-1}^1 dy R_\varepsilon^N(x, y) \left\{ \partial_y(\rho(t, y) - \bar{\rho}(y)) - \sigma(\rho(t, y)) \nabla H(t, y) \right\} \\
&\quad + \int_{-1}^1 dy R_\varepsilon^D(x, y) (C(\rho(t, y)) e^{G(t, y)} - A(\rho(t, y)) e^{-G(t, y)}).
\end{aligned}$$

Taking the time derivative we obtain

$$\partial_t \int_{-1}^1 dx U_\delta \left(\rho^{1, \varepsilon}(t, x) - \rho^{2, \varepsilon}(t, x) \right) = \int_{-1}^1 dx U'_\delta \left(\rho^{1, \varepsilon}(t, x) - \rho^{2, \varepsilon}(t, x) \right) \partial_t \left(\rho^{1, \varepsilon}(t, x) - \rho^{2, \varepsilon}(t, x) \right)$$

As $\partial_x R_\varepsilon^D(x, y) = -\partial_y R_\varepsilon^N(x, y)$, we get

$$\begin{aligned} \partial_x (\rho^{1,\varepsilon}(t, x) - \rho^{2,\varepsilon}(t, x)) &= - \int_{-1}^1 dy \partial_y R_\varepsilon^N(x, y) (\rho^1(t, y) - \rho^2(t, y)) \\ &= \int_{-1}^1 dy R_\varepsilon^N(x, y) (\partial_y \rho^1(t, y) - \partial_y \rho^2(t, y)). \end{aligned}$$

where we used that ρ^1, ρ^2 are in $\mathbb{L}^1([0, T], \mathbb{H}_1([-1, 1]))$. Thus we can write

$$\partial_x \left[U'_\delta(\rho^{1,\varepsilon}(t, x) - \rho^{2,\varepsilon}(t, x)) \right] = 1_{\{x \in V_{\delta,t}^\varepsilon\}} R_\varepsilon^N(\partial_x(\rho^1 - \rho^2))(t, x),$$

with the notation

$$V_{\delta,t}^\varepsilon = \left\{ x \in [-1, 1], \quad \text{such that} \quad |\rho^{1,\varepsilon}(t, x) - \rho^{2,\varepsilon}(t, x)| \leq \delta \right\}.$$

Using relation (7.7), one obtains

$$\begin{aligned} &\partial_t \int_{-1}^1 dx U_\delta(\rho^{1,\varepsilon}(t, x) - \rho^{2,\varepsilon}(t, x)) \\ &= - \int_{-1}^1 dx \int_{-1}^1 dy 1_{\{x \in V_{\delta,t}^\varepsilon\}} R_\varepsilon^N(\partial_x(\rho^1 - \rho^2))(t, x) \\ &\quad R_\varepsilon^N(x, y) \left\{ \partial_y(\rho^1(t, y) - \rho^2(t, y)) + \bar{\sigma}(t, y) \nabla H(t, y) \right\} \\ &\quad + \iint_{-1}^1 dx dy U'_\delta((\rho^{1,\varepsilon} - \rho^{2,\varepsilon})(t, x)) R_\varepsilon^D(x, y) (\bar{C}(t, y) e^{G(t,y)} - \bar{A}(t, y) e^{-G(t,y)}). \end{aligned}$$

Taking the limit as ε tends to 0, we recover (7.6).

Step 2.

First note that from assumption **(L2)**, one has

$$U'_\delta((\rho^1 - \rho^2)(t, x)) (\bar{C}(t, y) e^{G(t,y)} - \bar{A}(t, y) e^{-G(t,y)}) \leq 0.$$

Thus the reaction term acts as a contraction. As $\sigma(\rho) = \rho(1 - \rho)$, then

$$\forall (s, x) \in V_\delta, \quad |\bar{\sigma}(s, x)| \leq \delta.$$

Thus (7.6) implies

$$(7.8) \quad \begin{aligned} \langle U_\delta(\rho_{t'}^1 - \rho_{t'}^2) \rangle - \langle U_\delta(\rho_t^1 - \rho_t^2) \rangle &\leq -\frac{1}{\delta} \int_t^{t'} ds \langle 1_{\{(x,s) \in V_\delta\}} (\nabla \rho_s^1 - \nabla \rho_s^2)^2 \rangle \\ &\quad + \int_t^{t'} ds \langle 1_{\{(x,s) \in V_\delta\}} |\nabla(\rho_s^1 - \rho_s^2)| |\nabla H_s| \rangle, \end{aligned}$$

using the fact that

$$2|\nabla(\rho^1 - \rho^2)(s, x)| |\nabla H(s, x)| \leq \frac{1}{\delta} |\nabla(\rho^1 - \rho^2)(s, x)|^2 + \delta |\nabla H(s, x)|^2,$$

we obtain

$$\begin{aligned} \langle U_\delta(\rho_{t'}^1 - \rho_{t'}^2) \rangle - \langle U_\delta(\rho_t^1 - \rho_t^2) \rangle &\leq -\frac{1}{2\delta} \int_t^{t'} ds \langle 1_{\{(x,s) \in V_\delta\}} (\nabla \rho_s^1 - \rho_s^2)^2 \rangle + \delta \int_t^{t'} ds \langle |\nabla H_s|^2 \rangle \\ &\leq \delta \int_t^{t'} ds \langle |\nabla H_s|^2 \rangle. \end{aligned}$$

Recall that H belongs to $\mathbb{L}^2([0, T], \mathbb{H}_1(\cdot) - 1, 1)$ thus as δ tends to 0, the LHS converges to 0. Furthermore U_δ converges to the absolute value function. This implies that

$$\forall t \leq t', \quad \int_{-1}^1 dx \left| \rho^1(t', x) - \rho^2(t', x) \right| \leq \int_{-1}^1 dx \left| \rho^1(t, x) - \rho^2(t, x) \right|,$$

from which the uniqueness of the weak solutions follows. \square

REFERENCES

- [BDGJL1] L. Bertini, A. De Sole, D. Gabrielli, G. Jona Lasinio, C. Landim, *Large deviations for the boundary driven symmetric simple exclusion process*, Math. Phys. Anal. Geom. **6**, 231–267 (2003).
- [BDGJL2] L. Bertini, A. De Sole, D. Gabrielli, G. Jona Lasinio, C. Landim, *Large deviations of the empirical current in interacting particle systems*, Theory Probab. Appl. **51**, 2–27 (2007).
- [BDGJL3] L. Bertini, A. De Sole, D. Gabrielli, G. Jona Lasinio, C. Landim, *Stochastic interacting particle systems out of equilibrium*, J. Stat. Mech. P07014 (2007).
- [BLM] L. Bertini, C. Landim, M. Mourragui, *Dynamical large deviations for the boundary driven weakly asymmetric exclusion process*, Ann. Probab. **37**, no. 6, 2357–2403 (2009).
- [B] E. Bertin, *An exactly solvable dissipative transport model*, J. Phys. A: Math. Gen. **39**, 1539 (2006).
- [BD] T. Bodineau, B. Derrida, *Cumulants and large deviations of the current through non-equilibrium steady states*, C.R. Physique **8**, 540–555 (2007).
- [BL] T. Bodineau, M. Lagouge, *Current large deviations in a driven dissipative model*, J. Stat. Phys. **139**, no 2, 201–219 (2010).
- [DFL] A. De Masi, P. Ferrari, J. Lebowitz, *Rigorous Derivation of Reaction-Diffusion Equations with Fluctuations*, Phys. Rev. Lett., 55:1947–1949, (1985); *Reaction-Diffusion Equations for Interacting Particle Systems*, J. Stat. Phys., **44**, 589–644, (1986).
- [D] B. Derrida, *Non-equilibrium steady states: fluctuations and large deviations of the density and of the current*, J. Stat. Mech. P07023 (2007).
- [DV] M.D. Donsker, S.R.S. Varadhan, *Large deviations from a hydrodynamic scaling limit*, Comm. Pure Appl. Math. **42**, no. 3, 243–270 (1989).
- [DZ] A. Dembo, O. Zeitouni, *Large deviations techniques and applications*, Stochastic Modelling and Applied Probability, **38** Springer (2010).
- [EK] S. Ethier, T. Kurtz, *Markov processes. Characterization and convergence*, Wiley Series in Probability and Mathematical Statistics, John Wiley & Sons (1986).
- [E] L. Evans, *Partial differential equations*, Graduate Studies in Mathematics, **19** American Mathematical Society, Providence (1998).
- [FLM] J.S. Farfan Vargas, C. Landim, M. Mourragui, *Hydrodynamic behavior of boundary driven exclusion processes in dimension $d > 1$* , arXiv:0903.5526 (2009).
- [G] G. Gallavotti, *Fluctuation relation, fluctuation theorem, thermostats and entropy creation in non equilibrium statistical Physics*, Comptes Rendus de l’Académie de Sciences - Physique, (CR-Physique), **8**, 486–494, (2007).
- [JLV] G. Jona-Lasinio, C. Landim, M. E. Vares, *Large deviations for a reaction diffusion model*, Probab. Theory Related Fields **97**, no. 3, 339–361 (1993).

- [KL] C. Kipnis, C. Landim, *Scaling limits of interacting particle systems*, Springer (1999).
- [KOV] C. Kipnis, S. Olla, S.R.S. Varadhan, *Hydrodynamics and large deviation for simple exclusion processes*, Comm. Pure Appl. Math. **42**, no. 2, 115–137 (1989).
- [LMS] C. Landim, M. Mourragui, S. Sellami, *Hydrodynamic limit for a nongradient interacting particle system with stochastic reservoirs*, Theory Probab. Appl. **45**, no. 4, 604–623 (2002).
- [LL] D. Levanony, D. Levine, *Correlation and response in a driven dissipative model*, Phys. Rev. E **73**, 055102(R) (2006).
- [QRV] J. Quastel, F. Rezakhanlou, S.R.S. Varadhan, *Large deviations for the symmetric simple exclusion process in dimensions $d \geq 3$* , Probab. Theory Related Fields **113**, no. 1, 1–84 (1999).
- [S] H. Spohn, *Large scale dynamics of interacting particles*, Springer (1991).
- [SL] Y. Shokef, D. Levine, *Energy distribution and effective temperatures in a driven dissipative model*, Phys. Rev. E **74**, 051111 (2006).

ECOLE NORMALE SUPÉRIEURE, DMA, 45 RUE D'ULM 75230 PARIS CEDEX 05, FRANCE