

DJKM ALGEBRAS I: THEIR UNIVERSAL CENTRAL EXTENSION.

BEN COX AND VYACHESLAV FUTORNY

ABSTRACT. The purpose of this paper is to explicitly describe in terms of generators and relations the universal central extension of the infinite dimensional Lie algebra, $\mathfrak{g} \otimes \mathbb{C}[t, t^{-1}, u | u^2 = (t^2 - b^2)(t^2 - c^2)]$, appearing in the work of Date, Jimbo, Kashiwara and Miwa in their study of integrable systems arising from Landau-Lifshitz differential equation.

1. INTRODUCTION

In this paper the authors explicitly describe in terms of generators and relations and three families of polynomials, the universal central extension of an algebra appearing in work of Date, Jimbo, Kashiwara and Miwa (see [DJKM83, DJKM85]), where they study integrable systems arising from Landau-Lifshitz differential equation. Two of these families of polynomials are described below in terms of elliptic integrals and the other family is a variant of certain ultraspherical polynomials. The authors Date, Jimbo, Kashiwara and Miwa solved the Landau-Lifshitz equation using methods developed in some of their previous work on affine Lie algebras. The hierarchy of this equation is written in terms of free fermions on an elliptic curve. The infinite-dimensional Lie algebra mentioned above is shown to act on solutions of the Landau-Lifshitz equation as infinitesimal Bäcklund transformations where they derive an N -soliton formula. These authors arrive at an algebra that is a one dimensional central extension of $\mathfrak{g} \otimes \mathbb{C}[t, t^{-1}, u | u^2 = (t^2 - b^2)(t^2 - c^2)]$ where $b \neq \pm c$ are complex constants and \mathfrak{g} is a simple finite dimensional Lie algebra defined over the complex numbers. Below we explicitly describe its four dimensional universal central extension. Modulo the center this algebra is a particular example of a Krichever-Novikov current algebra (see ([KN87b], [KN87a], [KN89])). A fair amount of interesting and fundamental work has been done by Krichever, Novikov, Schlichenmaier, and Sheinman on the representation theory of certain one dimensional central extensions of these latter current algebras and of analogues of the Virasoro algebra. In particular Wess-Zumino-Witten-Novikov theory and analogues of the Knizhnik-Zamolodchikov equations are developed for these algebras (see the

2000 *Mathematics Subject Classification.* Primary 17B37, 17B67; Secondary 81R10, 81B50.

Key words and phrases. Krichever-Novikov Algebras, Landau-Lifshitz differential equation, Date-Jimbo-Miwa-Kashiwara algebras, universal central extension, ultraspherical polynomials, elliptic integrals.

The second author were partially supported by Fapesp (processo 2005/60337-2) and CNPq (processo 301743/2007-0). The first author is grateful to the Fapesp (processo 2009/17533-6) and the University of Sao Paulo for the support and hospitality during his visit to Sao Paulo. The first author was also partially supported by a research and development grant from the College of Charleston.

survey article [She05], and for example [SS99], [SS99],[She03],[Sch03a],[Sch03b], and [SS98]).

M. Bremner on the other hand has explicitly described in terms of generators, relations and certain families of polynomials (ultraspherical and Pollacyk) the structure constants for the universal central extension of algebras of the form $\mathfrak{g} \otimes \mathbb{C}[t, t^{-1}, u | u^2 = p(t)]$ where $p(t) = t^2 - 2bt + 1$ and $p(t) = t^3 - 2bt^2 + t$ (see [Bre95, Bre94]). He determined more generally the dimension of the universal central extension for affine Lie algebras of the form $\mathfrak{g} \otimes R$ where R is the ring of regular functions defined on an algebraic curve with any number of points removed. He obtained this using C. Kassel's result ([Kas84]) where one knows that the center is isomorphic as a vector space to Ω_R^1/dR (the *space of Kähler differentials of R modulo exact forms*). We will review this material below as needed.

In previous work of the authors (see [Cox08, BCF09]) we used Bremner's aforementioned description to obtain certain free field realizations of the four point and elliptic affine algebras depending on a parameter $r = 0, 1$ which correspond to two different normal orderings. These later realizations are analogues of Wakimoto type realizations which have been used by Schechtman and Varchenko and various other authors in the affine setting to pin down integral solutions to the Knizhnik-Zamolodchikov differential equations (see for example [ATY91], [Kur91], [EFK98], [SV90]). Such realizations have also been used in the study of Drinfeld-Sokolov reduction in the setting of W -algebras and in E. Frenkel's and B. Feigin's description of the center of the completed enveloping algebra of an affine Lie algebra (see [FFR94], [Fre05], and [FF92]). In future work we plan to use results of this paper to describe free field realizations of the universal central extension of the algebras of Date, Jimbo, Kashiwara and Miwa (which, since this is a mouth full, we will call DJKM algebras).

2. UNIVERSAL CENTRAL EXTENSIONS OF CURRENT ALGEBRAS

Let R be a commutative algebra defined over \mathbb{C} . Consider the left R -module with action $f(g \otimes h) = fg \otimes h$ for $f, g, h \in R$ and let K be the submodule generated by the elements $1 \otimes fg - f \otimes g - g \otimes f$. Then $\Omega_R^1 = F/K$ is the module of Kähler differentials. The element $f \otimes g + K$ is traditionally denoted by fdg . The canonical map $d : R \rightarrow \Omega_R^1$ by $df = 1 \otimes f + K$. The *exact differentials* are the elements of the subspace dR . The coset of fdg modulo dR is denoted by \overline{fdg} . As C. Kassel showed the universal central extension of the current algebra $\mathfrak{g} \otimes R$ where \mathfrak{g} is a simple finite dimensional Lie algebra defined over \mathbb{C} , is the vector space $\hat{\mathfrak{g}} = (\mathfrak{g} \otimes R) \oplus \Omega_R^1/dR$ with Lie bracket given by

$$[x \otimes f, Y \otimes g] = [xy] \otimes fg + (x, y) \overline{fdg}, [x \otimes f, \omega] = 0, [\omega, \omega'] = 0,$$

where $x, y \in \mathfrak{g}$, and $\omega, \omega' \in \Omega_R^1/dR$ and (x, y) denotes the Killing form on \mathfrak{g} .

Consider the polynomial

$$p(t) = t^n + a_{n-1}t^{n-1} + \cdots + a_0$$

where $a_i \in \mathbb{C}$ and $a_n = 1$. Fundamental to the description of the universal central extension for $R = \mathbb{C}[t, t^{-1}, u | u^2 = p(t)]$ is the following:

Theorem 2.0.1 ([Bre94], Theorem 3.4). *Let R be as above. The set*

$$\{\overline{t^{-1} dt}, \overline{t^{-1}u dt}, \dots, \overline{t^{-n}u dt}\}$$

forms a basis of Ω_R^1/dR (omitting $\overline{t^{-n}u dt}$ if $a_0 = 0$).

Set $u^m = p(t)$. Then $u d(u^m) = mu^m du$ and

$$\sum_{j=1}^n ja_j t^{j-1} u dt - m \left(\sum_{j=0}^n a_j t^j du \right) = 0$$

or

$$p'(t)u dt - mp(t)du = 0.$$

Multiplying by t^i we get

$$(2.1) \quad \sum_{j=1}^n ja_j t^{i+j-1} u dt - m \left(\sum_{j=0}^n a_j t^{i+j} du \right) = 0$$

Lemma 2.0.2. *If $u^m = p(t)$ and $R = \mathbb{C}[t, t^{-1}, u | u^m = p(t)]$, then in Ω_R^1/dR , one has*

$$(2.2) \quad ((m+1)n + im)t^{n+i-1} u dt \equiv - \sum_{j=0}^{n-1} ((m+1)j + mi)a_j t^{i+j-1} u dt \pmod{dR}$$

Proof. We have expanding $d(t^{i+j}u)$

$$(i+j)t^{i+j-1}u dt \equiv -t^{i+j} du \pmod{dR}.$$

so that (2.1) implies

$$(2.3) \quad \sum_{j=0}^n ja_j t^{i+j-1} u dt + m \left(\sum_{j=0}^n (i+j)a_j t^{i+j-1} u dt \right) = 0 \pmod{dR}$$

or

$$(2.4) \quad \sum_{j=0}^n ((m+1)j + mi)a_j t^{i+j-1} u dt \equiv 0 \pmod{dR}$$

This gives (2.2). □

3. DESCRIPTION OF THE UNIVERSAL CENTRAL EXTENSION OF DATE-JIMBO-MIWA-KASHIWARA ALGEBRAS

In the Date-Jimbo-Miwa-Kashiwara setting one takes $m = 2$ and $p(t) = (t^2 - a^2)(t^2 - b^2) = t^4 - (a^2 + b^2)t^2 + (ab)^2$ with $a \neq \pm b$ and neither a nor b is zero. We fix from here onward $R = \mathbb{C}[t, t^{-1}, u | u^2 = (t^2 - a^2)(t^2 - b^2)]$. As in this case $a_0 = (ab)^2$, $a_1 = 0$, $a_2 = -(a^2 + b^2)$, $a_3 = 0$ and $a_4 = 1$, then letting $k = i - 2$ the recursion relation in (2.2) looks like

$$(6 + 2k)\overline{t^k u dt} = -2(k-3)\overline{(ab)^2 t^{k-4} u dt} + 2k(a^2 + b^2)\overline{t^{k-2} u dt}.$$

After a change of variables we may assume that $a^2 b^2 = 1$. Then the recursion relation looks like

$$(3.1) \quad (6 + 2k)\overline{t^k u dt} = -2(k-3)\overline{t^{k-4} u dt} + 4kc\overline{t^{k-2} u dt},$$

after setting $c = (a^2 + b^2)/2$, so that $p(t) = t^4 - 2ct^2 + 1$. Let $P_k := P_k(c)$ be the polynomial in c satisfy the recursion relation

$$(6 + 2k)P_k(c) = 4kcP_{k-2}(c) - 2(k-3)P_{k-4}(c)$$

for $k \geq 0$. Then set

$$P(c, z) := \sum_{k \geq -4} P_k(c) z^{k+4} = \sum_{k \geq 0} P_{k-4}(c) z^k.$$

so that after some straightforward rearrangement of terms we have

$$\begin{aligned} 0 &= \sum_{k \geq 0} (6 + 2k) P_k(c) z^k - 4c \sum_{k \geq 0} k P_{k-2}(c) z^k + 2 \sum_{k \geq 0} (k - 3) P_{k-4}(c) z^k \\ &= (-2z^{-4} + 8cz^{-2} - 6)P(c, z) + (2z^{-3} - 4cz^{-1} + 2z) \frac{d}{dz} P(c, z) \\ &\quad + (2z^{-4} - 8cz^{-2})P_{-4}(c) - 4cP_{-3}(c)z^{-1} - 2P_{-2}(c)z^{-2} - 4P_{-1}(c)z^{-1}. \end{aligned}$$

We then multiply the above through by z^4 to get

$$\begin{aligned} 0 &= (-2 + 8cz^2 - 6z^4)P(c, z) + (2z - 4cz^3 + 2z^5) \frac{d}{dz} P(c, z) \\ &\quad + (2 - 8cz^2)P_{-4}(c) - 4cP_{-3}(c)z^3 - 2P_{-2}(c)z^2 - 4P_{-1}(c)z^3. \end{aligned}$$

Hence $P(c, z)$ must satisfy the differential equation

$$(3.2) \quad \frac{d}{dz} P(c, z) - \frac{3z^4 - 4cz^2 + 1}{z^5 - 2cz^3 + z} P(c, z) = \frac{2(P_{-1} + cP_{-3})z^3 + P_{-2}z^2 + (4cz^2 - 1)P_{-4}}{z^5 - 2cz^3 + z}$$

This has integrating factor

$$\begin{aligned} \mu(z) &= \exp \int \left(\frac{-2(z^3 - cz)}{1 - 2cz^2 + z^4} - \frac{1}{z} \right) dz \\ &= \exp\left(-\frac{1}{2} \ln(1 - 2cz^2 + z^4) - \ln(z)\right) = \frac{1}{z\sqrt{1 - 2cz^2 + z^4}}. \end{aligned}$$

3.1. Elliptic Case 1. If we take initial conditions $P_{-3}(c) = P_{-2}(c) = P_{-1}(c) = 0$ and $P_{-4}(c) = 1$ then we arrive at a generating function

$$P_{-4}(c, z) := \sum_{k \geq -4} P_{-4,k}(c) z^{k+4} = \sum_{k \geq 0} P_{-4,k-4}(c) z^k,$$

defined in terms of an elliptic integral

$$P_{-4}(c, z) = z\sqrt{1 - 2cz^2 + z^4} \int \frac{4cz^2 - 1}{z^2(z^4 - 2cz^2 + 1)^{3/2}} dz.$$

One way to interpret the right hand integral is to expand $(z^4 - 2cz^2 + 1)^{-3/2}$ as a Taylor series about $z = 0$ and then formally integrate term by term and multiply the result by the Taylor series of $z\sqrt{1 - 2cz^2 + z^4}$. More precisely one integrates formally with zero constant term

$$\int (4c - z^{-2}) \sum_{n=0}^{\infty} Q_n^{(3/2)}(c) z^{2n} dz = \sum_{n=0}^{\infty} \frac{4cQ_n^{(3/2)}(c)}{2n+1} z^{2n+1} - \sum_{n=0}^{\infty} \frac{Q_n^{(3/2)}(c)}{2n-1} z^{2n-1}$$

where $Q_n^{(\lambda)}(c)$ is the n -th Gegenbauer polynomial. After multiplying this by

$$z\sqrt{1 - 2cz^2 + z^4} = \sum_{n=0}^{\infty} Q_n^{(-1/2)}(c) z^{2n+1}$$

one arrives at the series $P_{-4}(c, z)$.

3.2. Elliptic Case 2. If we take initial conditions $P_{-4}(c) = P_{-3}(c) = P_{-1}(c) = 0$ and $P_{-2}(c) = 1$ then we arrive at a generating function defined in terms of another elliptic integral:

$$P_{-2}(c, z) = z\sqrt{1 - 2cz^2 + z^4} \int \frac{1}{(z^4 - 2cz^2 + 1)^{3/2}} dz.$$

3.3. Gegenbauer Case 3. If we take $P_{-1}(c) = 1$, and $P_{-2}(c) = P_{-3}(c) = P_{-4}(c) = 0$ and set

$$P_{-1}(c, z) = \sum_{n \geq 0} P_{-1, n-4} z^n,$$

then we get a solution which after solving for the integration constant can be turned into a power series solution

$$\begin{aligned} P_{-1}(c, z) &= (z\sqrt{1 - 2cz^2 + z^4}) \left(\int \frac{2cz^3}{t\sqrt{1 - 2cz^2 + z^4}(z^5 - 2cz^3 + z)} dt + C \right) \\ &= \frac{z(c - z^3)}{c^2 - 1} - \frac{c}{c^2 - 1} z\sqrt{z^4 - 2cz^2 + 1} \\ &= \frac{1}{c^2 - 1} \left(cz - z^3 - cz\sqrt{z^4 - 2cz^2 + 1} \right) \\ &= \frac{1}{c^2 - 1} \left(cz - z^3 - \sum_{k=0}^{\infty} cQ_n^{(-1/2)}(c)z^{2n+1} \right) \\ &= \frac{1}{c^2 - 1} \left(cz - z^3 - cz + c^2z^3 - \sum_{k=2}^{\infty} cQ_n^{(-1/2)}(c)z^{2n+1} \right) \end{aligned}$$

where $Q_n^{(-1/2)}(c)$ is the n -th Gegenbauer polynomial. Hence

$$\begin{aligned} P_{-1, -4}(c) &= P_{-1, -3}(c) = P_{-1, -2}(c) = P_{-1, 2m}(c) = 0, \\ P_{-1, -1}(c) &= 1, \\ P_{-1, 2n-3}(c) &= \frac{-cQ_n(c)}{c^2 - 1}, \end{aligned}$$

for $m \geq 0$ and $n \geq 2$. The $Q_n^{(-1/2)}(c)$ are known to satisfy the second order differential equation:

$$(1 - c^2) \frac{d^2}{dc^2} Q_n^{(-1/2)}(c) + n(n-1) Q_n^{(-1/2)}(c) = 0$$

so that the $P_{-1, k} := P_{-1, k}(c)$ satisfy the second order differential equation

$$(c^4 - c^2) \frac{d^2}{dc^2} P_{-1, 2n-3} + 2c(c^2 + 1) \frac{d}{dc} P_{-1, 2n-3} + (-c^2n(n-1) - 2) P_{-1, 2n-3} = 0$$

for $n \geq 2$.

3.4. Gegenbauer Case 4. Next we consider the initial conditions $P_{-1}(c) = 0 = P_{-2}(c) = P_{-4}(c) = 0$ with $P_{-3}(c) = 1$ and set

$$P_{-3}(c, z) = \sum_{n \geq 0} P_{-3, n-4}(c) z^n,$$

then we get a power series solution

$$\begin{aligned}
P_{-3}(c, z) &= (z\sqrt{1-2cz^2+z^4}) \left(\int \frac{2cz^3}{z\sqrt{1-2cz^2+z^4}(z^5-2cz^3+z)} dz + C \right) \\
&= \frac{cz(c-z^3)}{c^2-1} - \frac{1}{c^2-1} z\sqrt{z^4-2cz^2+1} \\
&= \frac{1}{c^2-1} \left(c^2z - cz^3 - z\sqrt{z^4-2cz^2+1} \right) \\
&= \frac{1}{c^2-1} \left(c^2z - cz^3 - \sum_{k=0}^{\infty} Q_n^{(-1/2)}(c) z^{2n+1} \right) \\
&= \frac{1}{c^2-1} \left(c^2z - cz^3 - z + cz^3 - \sum_{k=2}^{\infty} Q_n^{(-1/2)}(c) z^{2n+1} \right)
\end{aligned}$$

where $Q_n^{(-1/2)}(c)$ is the n -th Gegenbauer polynomial. Hence

$$\begin{aligned}
P_{-3,-4}(c) &= P_{-3,-2}(c) = P_{-3,-1}(c) = P_{-1,2m}(c) = 0, \\
P_{-3,-3}(c) &= 1, \\
P_{-3,2n-3}(c) &= \frac{-Q_n(c)}{c^2-1},
\end{aligned}$$

for $m \geq 0$ and $n \geq 2$ and hence

$$(c^2-1) \frac{d^2}{dc^2} P_{-3,2n-3} + 4c \frac{d}{dc} P_{-3,2n-3} - (n+1)(n-2) P_{-3,2n-3} = 0$$

for $n \geq 2$ and $P_{-1,2n-3} = cP_{-3,2n-3}$ for $n \geq 2$.

4. MAIN RESULT

First we give an explicit description of the cocycles contributing to the *even* part of the DJKM algebra.

Proposition 4.0.1 (cf. [Bre94], Prop. 4.2). *Set $\omega_0 = \overline{t^{-1} dt}$. For $i, j \in \mathbb{Z}$ one has*

$$(4.1) \quad t^i d(t^j) = j\delta_{i+j,0}\omega_0$$

and

$$(4.2) \quad t^{i-1} u d(t^{j-1} u) = (\delta_{i+j,-2}(j+1) - 2cj\delta_{i+j,0} + (j-1)\delta_{i+j,2})\omega_0.$$

Proof. First observe that $2u du = d(u^2) = (4t^3 - 4ct) dt$. The second congruence then follows from

$$\begin{aligned}
t^{i-1} u d(t^{j-1} u) &= (j-1)t^{i+j-3} u^2 dt + t^{i+j-2} u du \\
&= (j-1)t^{i+j-3} (t^4 - 2ct^2 + 1) dt + 2t^{i+j-2} (t^3 - ct) dt \\
&= (j-1)(t^{i+j+1} - 2ct^{i+j-1} + t^{i+j-3}) dt + 2(t^{i+j+1} - ct^{i+j-1}) dt \\
&= (j+1)t^{i+j+1} dt - 2cj t^{i+j-1} dt + (j-1)t^{i+j-3} dt.
\end{aligned}$$

□

The map $\sigma : R \rightarrow R$ given by $\sigma(t) = t^{-1}$, $\sigma(u) = t^{-2}u$ is an algebra automorphism as $\sigma(u^2) = t^{-4}u^2 = 1 - 2ct^{-2} + t^{-4} = \sigma(1 - 2ct^2 + t^4)$. This descends to a linear map $\sigma : \Omega_R^1/dR$ where

$$\begin{aligned}\sigma(\overline{t^{-1} dt}) &= \overline{-t^{-1} dt}, \\ \sigma(\overline{t^{-1}u dt}) &= \overline{t(t^{-2}u)d(t^{-1})} = \overline{-t^{-3}u dt}, \\ \sigma(\overline{t^{-2}u dt}) &= \overline{t^2(t^{-2}u)d(t^{-1})} = \overline{-t^{-2}u dt}, \\ \sigma(\overline{t^{-3}u dt}) &= \overline{-t^{-1}u dt}, \\ \sigma(\overline{t^{-4}u dt}) &= \overline{t^4(t^{-2}u)d(t^{-1})} = \overline{-u dt} = \overline{-t^{-4}u dt},\end{aligned}$$

whereby the last identity follows from the recursion relation (3.1) with $k = 0$. Setting $\omega_{-k} = \overline{t^{-k}u dt}$, $k = 1, 2, 3, 4$, then $\sigma(\omega_{-1}) = -\omega_{-3}$, and $\sigma(\omega_{-l}) = -\omega_{-l}$ for $l = 2, 4$.

Theorem 4.0.2. *Let \mathfrak{g} be a simple finite dimensional Lie algebra over the complex numbers with the Killing form $(\cdot | \cdot)$ and define $\psi_{ij}(c) \in \Omega_R^1/dR$ by*

$$(4.3) \quad \psi_{ij}(c) = \begin{cases} \omega_{i+j} & \text{for } i+j = 1, 0, -1, -2 \\ P_{-3, i+j-2}(c)(\omega_{-3} + c\omega_{-1}) & \text{for } i+j = 2n-1 \geq 3, n \in \mathbb{Z}, \\ P_{-3, i+j-2}(c)(c\omega_{-3} + \omega_{-1}) & \text{for } i+j = -2n+1 \leq -3, n \in \mathbb{Z}, \\ P_{-4, |i+j|-2}(c)\omega_{-4} + P_{-2, |i+j|-2}(c)\omega_{-2} & \text{for } |i+j| = 2n \geq 2, n \in \mathbb{Z}. \end{cases}$$

The universal central extension of the Date-Jimbo-Kashiwara-Miwa algebra is the \mathbb{Z}_2 -graded Lie algebra

$$\widehat{\mathfrak{g}} = \widehat{\mathfrak{g}}^0 \oplus \widehat{\mathfrak{g}}^1,$$

where

$$\widehat{\mathfrak{g}}^0 = (\mathfrak{g} \otimes \mathbb{C}[t, t^{-1}]) \oplus \mathbb{C}\omega_0, \quad \widehat{\mathfrak{g}}^1 = (\mathfrak{g} \otimes \mathbb{C}[t, t^{-1}]u) \oplus \mathbb{C}\omega_{-4} \oplus \mathbb{C}\omega_{-3} \oplus \mathbb{C}\omega_{-2} \oplus \mathbb{C}\omega_{-1}$$

with bracket

$$[x \otimes t^i, y \otimes t^j] = [x, y] \otimes t^{i+j} + \delta_{i+j, 0} j(x, y)\omega_0,$$

$$\begin{aligned}[x \otimes t^{i-1}u, y \otimes t^{j-1}u] &= [x, y] \otimes (t^{i+j+2} - 2ct^{i+j} + t^{i+j-2}) \\ &\quad + (\delta_{i+j, -2}(j+1) - 2cj\delta_{i+j, 0} + (j-1)\delta_{i+j, 2})(x, y)\omega_0,\end{aligned}$$

$$[x \otimes t^{i-1}u, y \otimes t^j] = [x, y]u \otimes t^{i+j-1} + j(x, y)\psi_{ij}(c).$$

Proof. The first two equalities follow from Proposition 4.0.1. For the last one we first observe that for $k = i + j - 2 \neq -3$,

$$\begin{aligned}j\omega_{ij}(c) &= \overline{t^{i-1}u d(t^j)} = \overline{jt^{i+j-2}u dt} \\ &= j \left(\frac{-2(k-3)\overline{t^{k-4}u dt} + 4kct^{k-2}u dt}{6+2k} \right),\end{aligned}$$

where the last equality is derived from (3.1). Then by setting $k = 0, 1, 2, 3, 4, 5$ in (3.1)

$$(6+2k)\overline{t^k u dt} = -2(k-3)\overline{t^{k-4}u dt} + 4kct^{k-2}u dt.$$

gives us

$$\begin{aligned}
\overline{6u dt} &= \overline{6t^{-4}u dt}, \\
\overline{8tu dt} &= \overline{4t^{-3}u dt} + \overline{4ct^{-1}u dt}, \\
\overline{10t^2u dt} &= \overline{2t^{-2}u dt} + \overline{8cu dt}, \\
\overline{12t^3u dt} &= \overline{12ctu dt}, \\
\overline{14t^4u dt} &= \overline{-2u dt} + \overline{16ct^2u dt}, \\
\overline{16t^5u dt} &= \overline{-4tu dt} + \overline{20ct^3u dt}, \\
(6+2k)\overline{t^k u dt} &= \overline{-2(k-3)t^{k-4}u dt} + \overline{4kct^{k-2}u dt}.
\end{aligned}$$

Hence when $i+j-2 = k = 0, 1, 2, 3, 4, 5$

$$\begin{aligned}
\overline{u dt} &= \omega_{-4}, \\
\overline{tu dt} &= \frac{1}{2}(\omega_{-3} + c\omega_{-1}), \\
\overline{t^2u dt} &= \frac{1}{5}\omega_{-2} + \frac{4c}{5}\omega_{-4}, \\
\overline{t^3u dt} &= \frac{c}{2}(\omega_{-3} + c\omega_{-1}), \\
\overline{t^4u dt} &= -\frac{1}{7}\overline{u dt} + \frac{8}{7}\overline{ct^2u dt} = -\frac{1}{7}\omega_{-4} + \frac{8}{7}c\left(\frac{1}{5}\omega_{-2} + \frac{4c}{5}\omega_{-4}\right) \\
&= \left(\frac{32c^2 - 5}{35}\right)\omega_{-4} + \frac{8}{35}c\omega_{-2}, \\
\overline{t^5u dt} &= -\frac{1}{8}(\omega_{-3} + c\omega_{-1}) + \frac{5c^2}{8}(\omega_{-3} + c\omega_{-1}) \\
&= \frac{5c^2 - 1}{8}(\omega_{-3} + c\omega_{-1}), \\
\overline{t^k u dt} &= \frac{-2(k-3)\overline{t^{k-4}u dt} + \overline{4kct^{k-2}u dt}}{6+2k}.
\end{aligned}$$

Thus by induction using the last equation above for $i+j-2 = k = 2n-3 \geq 1$, $n \in \mathbb{Z}$, we have

$$(4.4) \quad \omega_{ij}(c) = P_{-3, i+j-2}(c)(\omega_{-3} + c\omega_{-1}),$$

and for $i+j-2 = k = 2n-2 \geq 0$, $n \in \mathbb{Z}$, we have

$$(4.5) \quad \omega_{ij}(c) = P_{-4, i+j-2}(c)\omega_{-4} + P_{-2, i+j-2}(c)\omega_{-2}.$$

Applying σ to (4.4) for $i+j-2 = k = 2n-3 \geq 1$ to obtain

$$\begin{aligned}
j\sigma(\omega_{ij}(c)) &= \overline{t^{-i+1}u d(t^{-j})} = -j\overline{t^{-i-j-2}u dt} \\
&= j\sigma(P_{-3, i+j-2}(c)(\omega_{-3} + c\omega_{-1})) \\
&= -jP_{-3, i+j-2}(c)(\omega_{-1} + c\omega_{-3}).
\end{aligned}$$

Hence for $i+j-2 = 2n-3 \geq 1$

$$\omega_{-i, -j}(c) = \overline{t^{-i-j-2}u dt} = P_{-3, i+j-2}(c)(\omega_{-1} + c\omega_{-3}).$$

Setting $i' = -i$ and $j' = -j$ we get for $i'+j'-2 = -k-4 = -2n+3 \leq -5$

$$\omega_{i'j'}(c) = \overline{t^{i'+j'-2}u dt} = P_{-3, |i'+j'|-2}(c)(\omega_{-1} + c\omega_{-3}).$$

Similarly if we apply σ to (4.5) for $i + j = 2n \geq 2$, $n \in \mathbb{Z}$, we obtain

$$\begin{aligned} j\sigma(\omega_{ij}(c)) &= \overline{t^{-i+1}u d(t^{-j})} = -j\overline{t^{-i-j-2}u dt} \\ &= j\sigma(P_{-4,i+j-2}(c)\omega_{-4} + P_{-2,i+j-2}(c)\omega_{-2}) \\ &= -j(P_{-4,i+j-2}(c)\omega_{-4} + P_{-2,i+j-2}(c)\omega_{-2}) \end{aligned}$$

Hence for $i + j = 2n \geq 2$

$$\omega_{-i,-j}(c) = \overline{t^{-i-j-2}u dt} = P_{-4,i+j-2}(c)\omega_{-4} + P_{-2,i+j-2}(c)\omega_{-2}.$$

Setting $i' = -i$ and $j' = -j$ we get for $i' + j' = -2n \leq -2$

$$\omega_{i',j'}(c) = \overline{t^{i'+j'-2}u dt} = P_{-4,|i'+j'|-2}(c)\omega_{-4} + P_{-2,|i'+j'|-2}(c)\omega_{-2}.$$

□

One might want to compare the above theorem with the results that M. Bremner obtained for the elliptic and four point affine Lie algebra cases ([Bre94, Theorem 4.6] and [Bre95, Theorem 3.6] respectively).

REFERENCES

- [ATY91] Hidetoshi Awata, Akihiro Tsuchiya, and Yasuhiko Yamada. Integral formulas for the WZNW correlation functions. *Nuclear Phys. B*, 365(3):680–696, 1991.
- [BCF09] André Bueno, Ben Cox, and Vyacheslav Futorny. Free field realizations of the elliptic affine Lie algebra $\mathfrak{sl}(2, \mathbf{R}) \oplus (\Omega_R/dR)$. *J. Geom. Phys.*, 59(9):1258–1270, 2009.
- [Bre94] Murray Bremner. Universal central extensions of elliptic affine Lie algebras. *J. Math. Phys.*, 35(12):6685–6692, 1994.
- [Bre95] Murray Bremner. Four-point affine Lie algebras. *Proc. Amer. Math. Soc.*, 123(7):1981–1989, 1995.
- [Cox08] Ben Cox. Realizations of the four point affine Lie algebra $\mathfrak{sl}(2, R) \oplus (\Omega_R/dR)$. *Pacific J. Math.*, 234(2):261–289, 2008.
- [DJKM83] Etsurō Date, Michio Jimbo, Masaki Kashiwara, and Tetsuji Miwa. Landau-Lifshitz equation: solitons, quasiperiodic solutions and infinite-dimensional Lie algebras. *J. Phys. A*, 16(2):221–236, 1983.
- [DJKM85] Etsurō Date, Michio Jimbo, Masaki Kashiwara, and Tetsuji Miwa. On Landau-Lifshitz equation and infinite-dimensional groups. In *Infinite-dimensional groups with applications (Berkeley, Calif., 1984)*, volume 4 of *Math. Sci. Res. Inst. Publ.*, pages 71–81. Springer, New York, 1985.
- [EFK98] Pavel I. Etingof, Igor B. Frenkel, and Alexander A. Kirillov, Jr. *Lectures on representation theory and Knizhnik-Zamolodchikov equations*, volume 58 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1998.
- [FF92] Boris Feigin and Edward Frenkel. Affine Kac-Moody algebras at the critical level and Gel’fand-Dikiĭ algebras. In *Infinite analysis, Part A, B (Kyoto, 1991)*, volume 16 of *Adv. Ser. Math. Phys.*, pages 197–215. World Sci. Publ., River Edge, NJ, 1992.
- [FFR94] Boris Feigin, Edward Frenkel, and Nikolai Reshetikhin. Gaudin model, Bethe ansatz and critical level. *Comm. Math. Phys.*, 166(1):27–62, 1994.
- [Fre05] Edward Frenkel. Wakimoto modules, opers and the center at the critical level. *Adv. Math.*, 195(2):297–404, 2005.
- [Kas84] Christian Kassel. Kähler differentials and coverings of complex simple Lie algebras extended over a commutative algebra. In *Proceedings of the Luminy conference on algebraic K-theory (Luminy, 1983)*, volume 34, pages 265–275, 1984.
- [KN87a] Igor Moiseevich Krichever and S. P. Novikov. Algebras of Virasoro type, Riemann surfaces and strings in Minkowski space. *Funktsional. Anal. i Prilozhen.*, 21(4):47–61, 96, 1987.
- [KN87b] Igor Moiseevich Krichever and S. P. Novikov. Algebras of Virasoro type, Riemann surfaces and the structures of soliton theory. *Funktsional. Anal. i Prilozhen.*, 21(2):46–63, 1987.

- [KN89] Igor Moiseevich Krichever and S. P. Novikov. Algebras of Virasoro type, the energy-momentum tensor, and operator expansions on Riemann surfaces. *Funktsional. Anal. i Prilozhen.*, 23(1):24–40, 1989.
- [Kur91] Gen Kuroki. Fock space representations of affine Lie algebras and integral representations in the Wess-Zumino-Witten models. *Comm. Math. Phys.*, 142(3):511–542, 1991.
- [Sch03a] Martin Schlichenmaier. Higher genus affine algebras of Krichever-Novikov type. *Mosc. Math. J.*, 3(4):1395–1427, 2003.
- [Sch03b] Martin Schlichenmaier. Local cocycles and central extensions for multipoint algebras of Krichever-Novikov type. *J. Reine Angew. Math.*, 559:53–94, 2003.
- [She03] O. K. Sheĭnman. Second-order Casimirs for the affine Krichever-Novikov algebras $\widehat{\mathfrak{gl}}_{g,2}$ and $\widehat{\mathfrak{sl}}_{g,2}$. In *Fundamental mathematics today (Russian)*, pages 372–404. Nezasvis. Mosk. Univ., Moscow, 2003.
- [She05] O. K. Sheĭnman. Highest-weight representations of Krichever-Novikov algebras and integrable systems. *Uspekhi Mat. Nauk*, 60(2(362)):177–178, 2005.
- [SS98] M. Schlichenmaier and O. K. Scheinman. The Sugawara construction and Casimir operators for Krichever-Novikov algebras. *J. Math. Sci. (New York)*, 92(2):3807–3834, 1998. Complex analysis and representation theory, 1.
- [SS99] M. Shlikhenmaier and O. K. Sheĭnman. The Wess-Zumino-Witten-Novikov theory, Knizhnik-Zamolodchikov equations, and Krichever-Novikov algebras. *Uspekhi Mat. Nauk*, 54(1(325)):213–250, 1999.
- [SV90] V. V. Schechtman and A. N. Varchenko. Hypergeometric solutions of Knizhnik-Zamolodchikov equations. *Lett. Math. Phys.*, 20(4):279–283, 1990.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHARLESTON, 66 GEORGE ST., CHARLESTON, SC 29424, USA

E-mail address: `coxbl@cofc.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SÃO PAULO, SÃO PAULO, BRAZIL

E-mail address: `futorny@ime.usp.br`