

L_1 -uniqueness of degenerate elliptic operators

Derek W. Robinson¹ and Adam Sikora²

September 2010

Abstract

Let Ω be an open subset of \mathbf{R}^d with $0 \in \Omega$. Further let $H_\Omega = -\sum_{i,j=1}^d \partial_i c_{ij} \partial_j$ be a second-order partial differential operator with domain $C_c^\infty(\Omega)$ where the coefficients $c_{ij} \in W_{\text{loc}}^{1,\infty}(\overline{\Omega})$ are real, $c_{ij} = c_{ji}$ and the coefficient matrix $C = (c_{ij})$ satisfies bounds $0 < C(x) \leq c(|x|)I$ for all $x \in \Omega$. If

$$\int_0^\infty ds s^{d/2} e^{-\lambda \mu(s)^2} < \infty$$

for some $\lambda > 0$ where $\mu(s) = \int_0^s dt c(t)^{-1/2}$ then we establish that H_Ω is L_1 -unique, i.e. it has a unique L_1 -extension which generates a continuous semigroup, if and only if it is Markov unique, i.e. it has a unique L_2 -extension which generates a submarkovian semigroup. Moreover these uniqueness conditions are equivalent with the capacity of the boundary of Ω , measured with respect to H_Ω , being zero. We also demonstrate that the capacity depends on two gross features, the Hausdorff dimension of subsets A of the boundary the set and the order of degeneracy of H_Ω at A .

AMS Subject Classification: 47B25, 47D07, 35J70.

- | | |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------|
| 1. Centre for Mathematics
and its Applications
Mathematical Sciences Institute
Australian National University
Canberra, ACT 0200
Australia
derek.robinson@anu.edu.au | 2. Department of Mathematics
Macquarie University
Sydney, NSW 2109
Australia
sikora@ics.mq.edu.au |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------|

1 Introduction

In a recent paper [RS10b] we established that Markov uniqueness and L_1 -uniqueness are equivalent properties for a second-order, symmetric, elliptic operator with bounded Lipschitz continuous coefficients c_{ij} on an open subset Ω of \mathbf{R}^d . Moreover, these properties hold if and only if the corresponding capacity of the boundary $\partial\Omega$ of Ω is zero. In this note we extend these results to operators with locally bounded coefficients with a possible growth at infinity. As an illustration of our results we establish that Markov uniqueness, L_1 -uniqueness and the capacity condition are equivalent if the matrix $C = (c_{ij})$ satisfies $\|C(x)\| \sim |x|^2(\log|x|)^\alpha$ as $|x| \rightarrow \infty$ with $\alpha \in [0, 1]$. In addition we give an example with $\|C(x)\| \sim |x|^2(\log|x|)^{1+\varepsilon}$ as $|x| \rightarrow \infty$, where $\varepsilon > 0$ is arbitrarily small, which is Markov unique but not L_1 -unique. Our results extend uniqueness criteria previously established for the special case $\Omega = \mathbf{R}^d$ (see [Dav85], [Ebe99] Chapter 2, [Sta99] Section 2, and references therein).

Let Ω be an open subset of \mathbf{R}^d and choose coordinates such that $0 \in \Omega$. Define H_Ω as the positive symmetric operator on $L_2(\Omega)$ with domain $D(H_\Omega) = C_c^\infty(\Omega)$ and action

$$H_\Omega\varphi = - \sum_{i,j=1}^d \partial_i c_{ij} \partial_j \varphi \quad (1)$$

where $\partial_i = \partial/\partial x_i$ and the coefficients c_{ij} satisfy

$$\left. \begin{array}{l} 1. \quad c_{ij} = c_{ji} \in W_{\text{loc}}^{1,\infty}(\overline{\Omega}) \text{ are real,} \\ 2. \quad C(x) = (c_{ij}(x)) \text{ is a strictly positive-definite matrix for all } x \in \Omega \end{array} \right\} \quad (2)$$

where $W_{\text{loc}}^{s,p}(\Omega)$ denotes the local version of the usual Sobolev spaces and $W_{\text{loc}}^{s,p}(\overline{\Omega})$ denotes the restriction to Ω of functions in $W_{\text{loc}}^{s,p}(\mathbf{R}^d)$. The class of operators defined by (1) and (2) will be denoted by \mathcal{E}_Ω .

It follows that each $H_\Omega \in \mathcal{E}_\Omega$ is locally strongly elliptic, i.e. for each relatively compact $V \subset \Omega$ there are $\mu_V, \lambda_V > 0$ such that $\mu_V I \leq C(x) \leq \lambda_V I$ for all $x \in V$. There are, however, two potential sources of degeneracy. It is possible that $c_{ij}(x) \rightarrow 0$ as $x \rightarrow \partial\Omega$ or that $c_{ij}(x) \rightarrow \infty$ as $|x| \rightarrow \infty$.

In order to control the possible growth of the coefficients at infinity we introduce the strictly positive non-decreasing function c by

$$r \in \langle 0, \infty \rangle \mapsto c(r) = \sup\{\|C(x)\| : x \in \Omega, |x| < r\} \quad (3)$$

where $\|C(x)\|$ denotes the norm of the matrix $C(x) = (c_{ij}(x))$. Then $\|C(x)\| \leq c(|x|)$ and $c(0_+) > 0$. The growth conditions will be expressed either explicitly or implicitly in terms of the asymptotic properties of the positive increasing function μ given by

$$s \in \langle 0, \infty \rangle \mapsto \mu(s) = \int_0^s dt c(t)^{-1/2}. \quad (4)$$

This function is a lower bound on the Riemannian distance to infinity measured with respect to the metric C^{-1} . If, for example, $c(s) \sim s^2 (\log s)^\alpha$ as $s \rightarrow \infty$ with $\alpha \in [0, 2)$ then $\mu(s) \sim (\log s)^{1-\alpha/2} \rightarrow \infty$ as $s \rightarrow \infty$.

We are interested in criteria for various uniqueness properties of H_Ω and adopt the terminology of [Ebe99]. In particular H_Ω , viewed as an operator on $L_p(\Omega)$ for $p \in [1, \infty]$, is defined to be L_p -unique if it has a unique extension which generates an L_p -continuous semigroup. Moreover, it is defined to be Markov unique if it has a unique self-adjoint extension on $L_2(\Omega)$ which generates a submarkovian semigroup, i.e. an L_2 -continuous contraction semigroup S with the property that $0 \leq S_t \varphi \leq \mathbf{1}$ whenever $0 \leq \varphi \leq \mathbf{1}$. It follows that H_Ω is L_2 -unique if and only if it is essentially self-adjoint (see [Ebe99], Corollary 1.1.2). Then the self-adjoint closure is automatically submarkovian and H_Ω is Markov unique. Moreover, if H_Ω is L_1 -unique then it is Markov unique ([Ebe99], Lemma 1.1.6).

First, introduce the positive quadratic form h_Ω associated with H_Ω by

$$h_\Omega(\varphi) = \sum_{i,j=1}^d \int_{\Omega} dx c_{ij}(x) (\partial_i \varphi)(x) (\partial_j \varphi)(x) = (\varphi, H_\Omega \varphi)$$

with domain $D(h_\Omega) = D(H_\Omega) = C_c^\infty(\Omega)$. Since h_Ω is the form of the symmetric operator H_Ω it is closable with respect to the graph norm $\varphi \mapsto \|\varphi\|_{D(h_\Omega)} = (h_\Omega(\varphi) + \|\varphi\|_2^2)^{1/2}$. In the sequel we use the well known relationship between positive closed quadratic forms and positive self-adjoint operators (see [Kat80], Chapter 6) together with the corresponding theory of Dirichlet forms and submarkovian operators (see [BH91] [MR92] [FOT94]). The closure \bar{h}_Ω of h_Ω is automatically a Dirichlet form and the corresponding positive self-adjoint operator, the Friedrichs extension H_Ω^F of H_Ω , is submarkovian. Formally H_Ω^F corresponds to the self-adjoint extension of H_Ω with Dirichlet conditions on the boundary $\partial\Omega$ of Ω . In order to emphasize this interpretation we adopt the alternative notation $H_{\Omega,D} = H_\Omega^F$ and $h_{\Omega,D} = \bar{h}_\Omega$.

Secondly, we introduce a positive self-adjoint extension of H_Ω related to Neumann boundary conditions. Let $\chi \in C_c^\infty(\Omega)$ with $0 \leq \chi \leq \mathbf{1}_\Omega$ and define $h_{\Omega,\chi}$ as the form of the symmetric operator on $L_2(\Omega)$ with coefficients χc_{ij} . Then $h_{\Omega,\chi}$ is closable, its closure $\bar{h}_{\Omega,\chi}$ is a Dirichlet form and $\bar{h}_{\Omega,\chi} \leq h_{\Omega,D}$. Next set $\mathcal{C}_\Omega = \{\chi \in C_c^\infty(\Omega), 0 \leq \chi \leq \mathbf{1}_\Omega\}$. It follows that \mathcal{C}_Ω is a convex set which is directed with respect to the natural order and if $\chi, \eta \in \mathcal{C}_\Omega$ with $\chi \leq \eta$ then $\bar{h}_{\Omega,\chi} \leq \bar{h}_{\Omega,\eta}$. Now we define $h_{\Omega,N}$ by

$$h_{\Omega,N}(\varphi) = \lim\{\bar{h}_{\Omega,\chi}(\varphi) : \chi \in \mathcal{C}_\Omega\} = \sup\{\bar{h}_{\Omega,\chi}(\varphi) : \chi \in \mathcal{C}_\Omega\}. \quad (5)$$

Since $h_{\Omega,N}$ is the limit of quadratic forms it is a quadratic form and since it is the supremum of a family of closed forms it is a closed form. It is automatically a Dirichlet form satisfying $h_{\Omega,N} \leq h_{\Omega,D}$. If $H_{\Omega,N}$ is the positive self-adjoint operator associated with $h_{\Omega,N}$ it readily follows that $H_{\Omega,N}$ is a submarkovian extension of H_Ω and $H_{\Omega,N} \leq H_{\Omega,D}$. If $\partial\Omega$ is smooth, or even Lipschitz continuous, then $H_{\Omega,N}$ corresponds to the extension of H_Ω with Neumann boundary conditions but we adopt this definition for general open Ω .

In order to formulate our main result on uniqueness properties we need two extra definitions.

The operator $H_\Omega \in \mathcal{E}_\Omega$ is defined to be conservative if the submarkovian semigroup $S^{\Omega,D}$ generated by $H_{\Omega,D}$ is conservative, i.e. if $S_t^{\Omega,D} \mathbf{1}_\Omega = \mathbf{1}_\Omega$ for all $t > 0$. Moreover, the capacity of the measurable subset $A \subset \bar{\Omega}$ relative to the operator H_Ω is defined by

$$\text{cap}_\Omega(A) = \inf \left\{ \|\psi\|_{D(h_{\Omega,N})}^2 : \psi \in D(h_{\Omega,N}) \text{ and there exists an open set } U \subset \mathbf{R}^d \text{ such that } U \supseteq A \text{ and } \psi \geq 1 \text{ a. e. on } U \cap \Omega \right\}.$$

Thus cap_Ω corresponds to the capacity relative to the Dirichlet form $h_{\Omega,N}$ as defined in [BH91] or [FOT94].

Theorem 1.1 *Assume $H_\Omega \in \mathcal{E}_\Omega$. Consider the following conditions:*

- I. H_Ω is conservative,
- II. H_Ω is L_1 -unique,
- III. H_Ω is Markov unique,
- IV. $\text{cap}_\Omega(\partial\Omega) = 0$.

Then $\text{I} \Leftrightarrow \text{II} \Rightarrow \text{III} \Rightarrow \text{IV}$.

Conversely, if $\mu(s) \rightarrow \infty$ as $s \rightarrow \infty$ then $\text{IV} \Rightarrow \text{III}$ or if

$$\int_0^\infty ds s^{d/2} e^{-\lambda \mu(s)^2} < \infty \tag{6}$$

for one $\lambda > 0$ then $\text{IV} \Rightarrow \text{III} \Rightarrow \text{I}$ and all four conditions are equivalent.

Since μ is a positive increasing function with $\mu(0_+) = 0$ the finiteness restriction (6) is a condition on the growth μ at infinity, i.e. an implicit condition on the possible growth of the coefficients of H_Ω . If the coefficients are uniformly bounded then $\mu(s) = O(s)$ as $s \rightarrow \infty$ and (6) is satisfied. Then the four conditions of the theorem are equivalent. This retrieves the results of Theorems 1.2 and 1.3 of [RS10b].

The theorem is in part a restatement of standard results. The equivalence of Conditions I and II was established by Davies [Dav85], Theorem 2.2, whose arguments were based on earlier results of Azencott [Aze74]. Although Davies assumptions were somewhat different his arguments apply with little modification to the current setting. The implication $\text{II} \Rightarrow \text{III}$ is a straightforward result which is established, for example, in [Ebe99] Lemma 1.16. The implication $\text{III} \Rightarrow \text{IV}$ follows as in the proof of Theorem 1.2 in [RS10b] for operators with $c_{ij} \in W^{1,\infty}(\Omega)$.

In the special case $c(s) \sim s^2 (\log s)^\alpha$ for large s it follows that $\mu(s) \sim (\log s)^{1-\alpha/2}$ and $\mu(s) \rightarrow \infty$ as $s \rightarrow \infty$ for $\alpha \in [0, 2)$. On the other hand if $\alpha \in [0, 1]$ then (6) is satisfied for all sufficiently large $\lambda > 0$. Thus if $\alpha \in [0, 2)$ then Markov uniqueness of H_Ω is equivalent to the capacity of the boundary being zero and if $\alpha \in [0, 1]$ then it is also equivalent to L_1 -uniqueness of H_Ω .

Note that if $\Omega = \mathbf{R}^d$ the capacity condition is clearly satisfied and one concludes that $H_{\mathbf{R}^d}$ is L_1 -unique whenever (6) is satisfied for one large $\lambda > 0$. More generally we establish in Section 4.2 that the capacity condition depends on the Hausdorff dimension of bounded subsets $A \subset \partial\Omega$ and the order of degeneracy of H_Ω at A .

2 Submarkovian extensions

The Friedrichs extension $H_{\Omega,D}$ of H_Ω is well known to be the largest submarkovian extension, i.e. the extension with the minimal form domain. In this section we examine some basic properties of the smallest submarkovian extension, i.e. the extension with the maximal form domain (see, [FOT94] Section 3.3.3, [Ebe99] Section 3c or [RS10b], Section 3). In particular we identify $H_{\Omega,N}$ as the smallest submarkovian extension.

We begin by discussing the imposition of Neumann boundary conditions on a general submarkovian extension K_Ω of $H_\Omega \in \mathcal{E}_\Omega$. Let k_Ω be the Dirichlet form corresponding to K_Ω . Then $D(k_\Omega) \cap L_\infty(\Omega)$ is an algebra. Clearly $C_c^\infty(\Omega) \subseteq D(k_\Omega) \cap L_\infty(\Omega)$. Thus one can define the truncated form $k_{\Omega,\chi}$ for each $\chi \in C_c^\infty(\Omega)$ by $D(k_{\Omega,\chi}) = D(k_\Omega) \cap L_\infty(\Omega)$ and

$$k_{\Omega,\chi}(\varphi) = k_\Omega(\varphi, \chi\varphi) - 2^{-1}k_\Omega(\chi, \varphi^2) \quad (7)$$

for $\varphi \in D(k_{\Omega,\chi})$. The $k_{\Omega,\chi}$ have many properties similar to those of the forms $h_{\Omega,\chi}$. In particular the $k_{\Omega,\chi}$ are Markovian forms satisfying $0 \leq k_{\Omega,\chi} \leq k_\Omega$. Moreover, if $\chi_1, \chi_2 \in \mathcal{C}_\Omega$ and $\chi_1 \leq \chi_2$ then $k_{\Omega,\chi_1} \leq k_{\Omega,\chi_2}$ (see [BH91], Proposition I.4.1.1). But it is not evident that the $k_{\Omega,\chi}$ are closable. This, however, is part of our first result.

Theorem 2.1 *Let $H_\Omega \in \mathcal{E}_\Omega$. Further let K_Ω be a submarkovian extension of H_Ω and k_Ω the corresponding Dirichlet form.*

If $\chi \in C_c^\infty(\Omega)$ then the truncated form $k_{\Omega,\chi}$ defined by (7) is closable and the closure $\overline{k}_{\Omega,\chi}$ satisfies $\overline{k}_{\Omega,\chi} = \overline{h}_{\Omega,\chi}$. Therefore

$$h_{\Omega,N} \leq k_\Omega \leq h_{\Omega,D} .$$

In particular H_Ω is Markov unique if and only if $h_{\Omega,N} = h_{\Omega,D}$.

Proof The first step in the proof is a regularity property which extends a similar result for operators with bounded coefficients given by Theorem 1.1.IV in [RS10b].

Lemma 2.2 *Let K_Ω be a positive, self-adjoint extension of H_Ω . Then*

$$C_c^\infty(\Omega)D(K_\Omega) \subseteq D(\overline{H}_\Omega) .$$

Proof First, if K_Ω is a self-adjoint extension of H_Ω then $H_\Omega \subseteq K_\Omega \subseteq H_\Omega^*$. Therefore it suffices to establish that $C_c^\infty(\Omega)D(H_\Omega^*) \subseteq D(\overline{H}_\Omega)$. This property was proved for operators with bounded Lipschitz coefficients in Theorem 2.1 of [RS10b] but the proof is also valid for operators with coefficients which are only locally bounded. For example, if $\eta \in C_c^\infty(\Omega)$ with $\text{supp } \eta = K$ and V is a relatively compact subset of Ω with $K \subset V$ then to deduce that $\eta D(H_\Omega^*) \subseteq D(\overline{H}_\Omega)$ it suffices to prove that $\eta D(H_V^*) \subseteq D(\overline{H}_V)$ where H_V is the restriction of H_Ω to $C_c^\infty(V)$. Since the coefficients of H_V are uniformly bounded the result follows from Theorem 2.1 of [RS10b]. \square

Next we prove the first statement of Theorem 2.1.

Lemma 2.3 *If $\chi \in C_c^\infty(\Omega)$ then $k_{\Omega,\chi}$ is closable and the closure $\overline{k}_{\Omega,\chi}$ satisfies $\overline{k}_{\Omega,\chi} = \overline{h}_{\Omega,\chi}$.*

Proof First $C_c^\infty(\Omega)D(K_\Omega) \subseteq D(\overline{H}_\Omega)$ by Lemma 2.2. Now fix $\varphi \in D(K_\Omega) \cap L_\infty(\Omega)$. Then for each $\chi \in \mathcal{C}_\Omega$ one has $\chi\varphi \in D(\overline{H}_\Omega)$. Moreover,

$$k_\Omega(\varphi, \chi\varphi) = (K_\Omega\varphi, \chi\varphi) = (\varphi, \overline{H}_\Omega\chi\varphi)$$

and

$$k_\Omega(\chi, \varphi^2) = (K_\Omega\chi, \varphi^2) = (H_\Omega\chi, \varphi^2) .$$

Therefore

$$k_{\Omega,\chi}(\varphi) = (\varphi, \overline{H}_\Omega\chi\varphi) - 2^{-1}(H_\Omega\chi, \varphi^2) .$$

Next choose a $\chi_1 \in \mathcal{C}_\Omega$ with $\chi_1 = 1$ on $\text{supp } \chi$ and set $\varphi_1 = \chi_1 \varphi$. It follows from Lemma 2.2 that $\varphi_1 \in D(\overline{H}_\Omega) \cap L_\infty(\Omega)$. Moreover,

$$k_{\Omega, \chi}(\varphi) = (\varphi, \overline{H}_\Omega \chi \varphi_1) - 2^{-1}(H_\Omega \chi, \varphi_1^2) = (\varphi_1, \overline{H}_\Omega \chi \varphi_1) - 2^{-1}(H_\Omega \chi, \varphi_1^2) = \overline{h}_{\Omega, \chi}(\varphi_1).$$

The first equality is obvious since $\text{supp } \overline{H}_\Omega \chi = \text{supp } \chi$. The second equality follows by approximating φ in $L_2(\Omega)$ by a sequence $\varphi_n \in C_c^\infty(\Omega)$ and noting that

$$(\varphi_n, \overline{H}_\Omega \chi \varphi_1) = (H_\Omega \varphi_n, \chi \varphi_1) = (H_\Omega \chi_1 \varphi_n, \chi \varphi_1) = (\chi_1 \varphi_n, \overline{H}_\Omega \chi \varphi_1).$$

The third equality is also obvious. But for χ and φ fixed $\overline{h}_{\Omega, \chi}(\chi_1 \varphi)$ is independent of the choice of χ_1 . Moreover, if χ_2 is a second choice, with $\chi_2 = 1$ on $\text{supp } \chi$ then $\chi_1 - \chi_2 = 0$ on $\text{supp } \chi$ and $\overline{h}_{\Omega, \chi}((\chi_1 - \chi_2)\varphi) = 0$. Therefore if $\chi_1 \leq \chi_2 \leq \dots \leq \mathbf{1}_\Omega$ is an increasing family of C_c^∞ -functions with $\chi_n = 1$ on $\text{supp } \chi$ then $\overline{h}_{\Omega, \chi}((\chi_n - \chi_m)\varphi) = 0$ but $\|\chi_n \varphi - \varphi\|_2 \rightarrow 0$. This establishes that $\varphi \in D(\overline{h}_{\Omega, \chi})$ and $\overline{h}_{\Omega, \chi}(\varphi) = k_{\Omega, \chi}(\varphi)$. Then, however,

$$\overline{h}_{\Omega, \chi}(\varphi) = k_{\Omega, \chi}(\varphi) \leq k_\Omega(\varphi)$$

for all $\varphi \in D(K_\Omega) \cap L_\infty(\Omega)$. Since $D(K_\Omega)$ is a core of k_Ω it follows by continuity that $\overline{h}_{\Omega, \chi}(\varphi) = k_{\Omega, \chi}(\varphi)$ for all $\varphi \in D(k_\Omega) \cap L_\infty(\Omega) = D(k_{\Omega, \chi})$. Therefore $\overline{h}_{\Omega, \chi}$ is a closed extension of $k_{\Omega, \chi}$. Thus $k_{\Omega, \chi}$ is closable and its closure $\overline{k}_{\Omega, \chi} \subseteq \overline{h}_{\Omega, \chi}$.

But $h_{\Omega, \chi}(\psi) = k_{\Omega, \chi}(\psi)$ for all $\psi \in C_c^\infty(\Omega)$ and $C_c^\infty(\Omega)$ is a core of $\overline{h}_{\Omega, \chi}$ by definition. Therefore $\overline{k}_{\Omega, \chi} \supseteq \overline{h}_{\Omega, \chi}$. Combination of these conclusions gives $\overline{k}_{\Omega, \chi} = \overline{h}_{\Omega, \chi}$. \square

One can now immediately deduce Theorem 2.1.

Proof of Theorem 2.1 The first statement of the theorem has been established by Lemma 2.3. Hence

$$h_{\Omega, N} = \sup_{\chi \in \mathcal{C}_\Omega} \overline{k}_{\Omega, \chi}.$$

But $k_{\Omega, \chi} \leq k_\Omega$ for all $\chi \in \mathcal{C}_\Omega$. Therefore $h_{\Omega, N} \leq k_\Omega$.

Finally $k_\Omega \supseteq h_\Omega$. Hence $k_\Omega \leq \overline{h}_\Omega = h_{\Omega, D}$. \square

The form $h_{\Omega, N}$ possesses a *carré du champ* in the sense of [BH91], Section I.4. This is initially defined as the bilinear form from $W_{\text{loc}}^{1,2}(\Omega) \times W_{\text{loc}}^{1,2}(\Omega)$ into $L_{1,\text{loc}}(\Omega)$ given by

$$\Gamma(\varphi; \psi)(x) = \sum_{i,j=1}^d c_{ij}(x) (\partial_i \varphi)(x) (\partial_j \psi)(x)$$

and $\Gamma(\varphi) = \Gamma(\varphi; \varphi)$. Then

$$D(h_{\Omega, N}) = \left\{ \varphi \in W_{\text{loc}}^{1,2}(\Omega) : \sup_V \int_V dx \Gamma(\varphi)(x) < \infty \right\}$$

where the supremum is over the relatively compact subsets V of Ω and

$$h_{\Omega, N}(\varphi) = \sup_V \int_V dx \Gamma(\varphi)(x)$$

for all $\varphi \in D(h_{\Omega, N})$. It follows readily that if $\varphi \in D(h_{\Omega, N})$ then $\Gamma(\varphi)$ is a positive $L_1(\Omega)$ -function with $\|\Gamma(\varphi)\|_1 = h_{\Omega, N}(\varphi)$. The foregoing explicit identification of the form of the

minimal extension has been used in previous discussions of Markov uniqueness, [FOT94] Section 3.3.3, [Ebe99] Section 3c or [RS10b], Section 3.

A number of properties of general submarkovian extension follows from the identification of the minimal extension. If k_Ω is the form of the submarkovian extension K_Ω of H_Ω it follows from Theorem 2.1 that $D(k_\Omega) \subseteq D(h_{\Omega,N})$. Therefore k_Ω possesses a *carré du champ* since $\Gamma(\varphi) \in L_1(\Omega)$ for all $\varphi \in D(k_\Omega)$. Moreover, $k_\Omega(\varphi) = \|\Gamma(\varphi)\|_1$ for all $\varphi \in D(k_\Omega)$. Further the form $h_{\Omega,N}$ is strongly local in the sense of [FOT94] and hence the restriction k_Ω is also strongly local.

Subsequently we need two Dirichlet form implications of the elliptic regularity property.

Corollary 2.4 *Let K_Ω be a submarkovian extension of $H_\Omega \in \mathcal{E}_\Omega$ and k_Ω the corresponding Dirichlet form. Then*

$$C_c^\infty(\Omega)D(k_\Omega) \subseteq D(\bar{h}_\Omega) .$$

Proof If $\eta \in C_c^\infty(\Omega)$ and $\varphi \in D(K_\Omega)$ then it follows from Lemma 2.2 that $\eta\varphi \in D(\bar{H}_\Omega) \subset D(K_\Omega) \subseteq D(k_\Omega) \subseteq D(h_{\Omega,N})$. Moreover,

$$\bar{h}_\Omega(\eta\varphi) = h_{\Omega,N}(\eta\varphi) \leq 2 \int_\Omega \Gamma(\eta) \varphi^2 + 2 \int_\Omega \eta^2 \Gamma(\varphi) \leq 2 (\|\Gamma(\eta)\|_\infty + \|\eta\|_\infty^2) \|\varphi\|_{D(k_\Omega)}^2 .$$

Since $D(K_\Omega)$ is a core of k_Ω with respect to the $D(k_\Omega)$ -graph norm this estimate extends to all $\varphi \in D(k_\Omega)$ by continuity. The statement of the corollary follows immediately. \square

Corollary 2.5 *If $H_\Omega \in \mathcal{E}_\Omega$ then $C_c^\infty(\mathbf{R}^d)D(h_{\Omega,N}) \subseteq D(h_{\Omega,N})$.*

Proof Fix $\rho \in C_c^\infty(\mathbf{R}^d)$ and $\chi \in \mathcal{C}_\Omega$. If $\varphi \in C_c^\infty(\Omega)$. Then $\rho\varphi \in C_c^\infty(\Omega)$ and $\|\rho\varphi\|_2 \leq \|\rho\|_\infty \|\varphi\|_2$. Moreover,

$$h_{\Omega,\chi}(\rho\varphi) \leq 2 \|\rho\|_\infty^2 h_{\Omega,\chi}(\varphi) + 2 a_\rho \|\nabla\rho\|_\infty^2 \|\varphi\|_2^2$$

where $a_\rho = \sup_{x \in \text{supp } \rho} \|C(x)\|$. Therefore, by continuity, $\rho D(\bar{h}_{\Omega,\chi}) \subseteq D(\bar{h}_{\Omega,\chi})$ and

$$\|\rho\varphi\|_{D(\bar{h}_{\Omega,\chi})} \leq a(\rho) \|\varphi\|_{D(\bar{h}_{\Omega,\chi})}$$

for all $\varphi \in D(\bar{h}_{\Omega,\chi})$ with $a(\rho) = 2 (a_\rho \|\nabla\rho\|_\infty^2 + \|\rho\|_\infty^2)$. Since this estimate is uniform for $\chi \in \mathcal{C}_\Omega$ it follows that $\rho D(h_{\Omega,N}) \subseteq D(h_{\Omega,N})$ and $\|\rho\varphi\|_{D(h_{\Omega,N})} \leq a(\rho) \|\varphi\|_{D(h_{\Omega,N})}$. \square

3 L_1 -uniqueness

In this section we prove Theorem 1.1. Much of the proof consists of refinements of previous arguments.

I \Leftrightarrow II This equivalence was established by Davies [Dav85], Theorem 2.2, for a large class of second-order elliptic operators with smooth coefficients. But his arguments extend to the current situation with only minor modifications. We omit further details

II \Rightarrow III This is a general structural result which is proved, for example, in Lemma 1.1.6 of [Ebe99].

III \Rightarrow IV First note that Markov uniqueness of H_Ω is equivalent to the identity $h_{\Omega,N} = h_{\Omega,D}$ by Theorem 2.1. But in general $h_{\Omega,N} \supseteq h_{\Omega,D}$ and $C_c^\infty(\Omega)$ is a core of $h_{\Omega,D}$. Therefore Markov uniqueness of H_Ω implies that $C_c^\infty(\Omega)$ is a core of $h_{\Omega,N}$.

Secondly, let $\psi \in D(h_{\Omega,N}) \cap L_\infty(\Omega)$ with $\psi = 1$ on $U \cap \Omega$ where U is an open subset containing $\partial\Omega$. Then since $C_c^\infty(\Omega)$ is a core of $h_{\Omega,N}$ there is a sequence $\psi_n \in C_c^\infty(\Omega)$ such that $\|\psi - \psi_n\|_{D(h_{\Omega,N})} \rightarrow 0$ as $n \rightarrow \infty$. Set $\varphi_n = \psi - \psi_n$. Then $\varphi_n \in D(h_{\Omega,N})$, $\|\varphi_n\|_{D(h_{\Omega,N})} \rightarrow 0$ and since ψ_n has compact support there is an open subset U_n containing $\partial\Omega$ such that $\varphi_n = 1$ on $(U \cap U_n) \cap \Omega$. Therefore $\text{cap}_\Omega(\partial\Omega) = 0$.

Combination of the foregoing observations establishes the first statement of Theorem 1.1. Now we turn to the proof of the second statement.

IV \Rightarrow III Assume $\mu(s) \rightarrow \infty$ as $s \rightarrow \infty$ where μ is defined by (3) and (4). Then H_Ω is Markov unique if and only if $C_c^\infty(\Omega)$ is a core of $h_{\Omega,N}$. Thus it is necessary to demonstrate that each $\varphi \in D(h_{\Omega,N}) \cap L_\infty(\Omega)$ can be approximated in the $D(h_{\Omega,N})$ -graph norm by a sequence $\varphi_n \in C_c^\infty(\Omega)$.

Fix $\varphi \in D(h_{\Omega,N}) \cap L_\infty(\Omega)$. Next fix $\rho \in C_c^\infty(\mathbf{R})$ with $0 \leq \rho \leq 1$, $\rho(s) = 1$ if $s \leq 1$ and $\rho(s) = 0$ if $s \geq 2$. Then define ρ_n by $\rho_n(x) = \rho(n^{-1}\mu(|x|))$. It follows that $\rho_n(x) = 1$ if $\mu(|x|) \leq n$ and $\rho_n(x) = 0$ if $\mu(|x|) \geq 2n$. Moreover, $\|\Gamma(\rho_n)\|_\infty \leq b^2 n^{-2}$ with $b = \|\rho'\|_\infty$. Then $\rho_n \varphi \in D(h_{\Omega,N}) \cap L_\infty(\Omega)$ by Corollary 2.5 and

$$\begin{aligned} \|\varphi - \rho_n \varphi\|_{D(h_{\Omega,N})}^2 &\leq 2 \int_\Omega \varphi^2 \Gamma(\rho_n) + 2 \int_\Omega (\mathbf{1}_\Omega - \rho_n)^2 \Gamma(\varphi) + \|(\mathbf{1}_\Omega - \rho_n)\varphi\|_2^2 \\ &\leq 2b^2 n^{-2} \|\varphi\|_2^2 + \int_\Omega (\mathbf{1}_\Omega - \rho_n)^2 (2\Gamma(\varphi) + \varphi^2). \end{aligned}$$

The first term on the right hand side clearly tends to zero as $n \rightarrow \infty$. But it follows by construction that $(\mathbf{1}_\Omega - \rho_n)^2 \rightarrow 0$ pointwise on Ω . Therefore the second term also tends to zero by the Lebesgue dominated convergence theorem. Thus φ is approximated by the sequence $\rho_n \varphi$ in the $D(h_{\Omega,N})$ -graph norm.

Next since $\text{cap}_\Omega(\partial\Omega) = 0$ one may choose $\chi_n \in D(h_{\Omega,N})$ and open subsets $U_n \supset \partial\Omega$ such that $0 \leq \chi_n \leq 1$, $\|\chi_n\|_{D(h_{\Omega,N})} \leq n^{-1}$ and $\chi_n \geq 1$ on $U_n \cap \Omega$. But since $h_{\Omega,N}$ is a Dirichlet form one may assume $\chi_n = 1$ on $U_n \cap \Omega$. Then with $\varphi_n = (\mathbf{1}_\Omega - \chi_n)\rho_n \varphi$ one has

$$\lim_{n \rightarrow \infty} \|\varphi - \varphi_n\|_{D(h_{\Omega,N})} \leq \lim_{n \rightarrow \infty} \|\chi_n \rho_n \varphi\|_{D(h_{\Omega,N})}$$

by the Cauchy–Schwarz estimate and the conclusion of the previous paragraph. But

$$\|\chi_n \rho_n \varphi\|_{D(h_{\Omega,N})}^2 = h_{\Omega,N}(\chi_n \rho_n \varphi) + \|\chi_n \rho_n \varphi\|_2^2$$

and the second term on the right hand side tends to zero as $n \rightarrow \infty$ because $\|\chi_n \rho_n \varphi\|_2 \leq \|\chi_n\|_2 \|\varphi\|_\infty$. The first term on the right can, however, be estimated by

$$\begin{aligned} h_{\Omega,N}(\chi_n \rho_n \varphi) &\leq 2 \int_\Omega \varphi^2 \Gamma(\chi_n) + 4 \int_\Omega \varphi^2 \Gamma(\rho_n) + 4 \int_\Omega \chi_n^2 \Gamma(\varphi) \\ &\leq 2 \|\varphi\|_\infty^2 h_{\Omega,N}(\chi_n) + 4b^2 n^{-2} \|\varphi\|_2^2 + 4 \int_\Omega \chi_n^2 \Gamma(\varphi) \end{aligned}$$

since $\|\Gamma(\rho_n)\|_\infty \leq b^2 n^{-2}$. The first term on the right hand side tends to zero because $h_{\Omega,N}(\chi_n) \leq n^{-1}$ and the second obviously tends to zero. Finally the third tends to zero by

an equicontinuity estimate because $\chi_n^2 \leq 1$ and $\Gamma(\varphi) \in L_1(\Omega)$. Thus one now concludes that φ is approximated by the sequence φ_n in the $D(h_{\Omega,N})$ -graph norm.

Finally, $\text{supp } \varphi_n \subseteq \Omega_n = ((\text{supp } \rho_n) \cap \Omega) \cap (\Omega \setminus (U_n \cap \Omega))$ and Ω_n is a relatively compact subset of Ω . Therefore $H_{\Omega,N}$ is strongly elliptic in restriction to Ω_n . Consequently φ_n , and hence φ , can be approximated by a sequence of $C_c^\infty(\Omega_n)$ -functions in the $D(h_{\Omega,N})$ -graph norm.

This completes the proof of the second statement of Theorem 1.1. Now we turn to the proof of the third statement. By the foregoing it suffices to prove the following.

III \Rightarrow I The proof is an elaboration of the argument used to demonstrate the comparable implication in Theorem 1.3 in [RS10b].

Proposition 3.1 *Assume $H_\Omega \in \mathcal{E}_\Omega$ is Markov unique. Further assume that*

$$\int_0^\infty ds s^{d/2} e^{-\lambda \mu(s)^2} < \infty$$

for one $\lambda > 0$ where $\mu(s) = \int_0^s c^{-1/2}$ with c defined by (3).

Then H_Ω is conservative.

Proof The proof is in several steps.

Step 1: Ω bounded. If Ω is bounded then H_Ω is conservative by Step 1 in the proof of Theorem 1.3 in [RS10b]. Therefore we now assume that Ω is unbounded.

Step 2: Bounded approximation. The second step consists of introducing an increasing sequence of bounded sets Ω_n and conservative operators $H_{\Omega;n} \in \mathcal{E}_{\Omega_n}$ which approximate H_Ω in a suitable manner.

First, fix $\rho \in C_c^\infty(\mathbf{R})$ with $0 \leq \rho \leq 1$, $\rho(s) = 1$ if $|s| \leq 1$ and $\rho(s) = 0$ if $|s| \geq 2$. Then introduce the sequence ρ_n by $\rho_n(x) = \rho(n^{-1}|x|)$. Thus $\rho_n(x) = 1$ if $|x| \leq n$ and $\rho_n(x) = 0$ if $|x| \geq 2n$. The family of functions ρ_n is monotonically increasing. Set $B_n = \{x \in \mathbf{R}^d : |x| < n\}$ and $\Omega_n = \Omega \cap B_{2n}$. Note that Ω_n is bounded.

Secondly, define $H_{\Omega;n} \in \mathcal{E}_{\Omega_n}$ as the operator with coefficients $\rho_n c_{ij}$ acting on $L_2(\Omega_n)$. Then it follows that $H_{\Omega;n}$ is Markov unique since the capacity of $\partial\Omega_n$ with respect to the Neumann form associated with $H_{\Omega;n}$ is zero. Therefore $H_{\Omega;n}$ is conservative by Step 1. Then if H_n is the extension to $L_2(\Omega)$ of the unique submarkovian extension $H_{\Omega;n,N}(= H_{\Omega;n,D})$ of $H_{\Omega;n}$ acting on $L_2(\Omega_n)$, i.e. if $H_n = H_{\Omega;n,N} \oplus 0$ with $L_2(\Omega) = L_2(\Omega_n) \oplus L_2(\Omega_n)^\perp$, then H_n is conservative.

Step 3: L_2 -convergence. The third step is to establish strong convergence on $L_2(\Omega)$ of the semigroups $S^{(n)}$ generated by the H_n to the semigroup S generated by the unique submarkovian extension $H_{\Omega,N}(= H_{\Omega,D})$ of H_Ω . This follows by a monotone convergence argument. The closed form h_n corresponding to H_n on $L_2(\Omega)$ is given by $h_n(\varphi) = h_{\Omega;n,N}(\varphi)$ for all $\varphi \in C_c^\infty(\Omega)$ and then by closure for all $\varphi \in D(h_n)$. Since the ρ_n are a monotonically increasing family of functions on \mathbf{R}^d the forms h_n are a monotonically increasing family of Dirichlet forms. If $h = \sup_{n \geq 1} h_n$ then h is a Dirichlet form.

It follows from the monotonic increase of the forms h_n that the operators H_n converge in the strong resolvent sense on $L_2(\Omega)$ to the operator H corresponding to h (see, for example, [Kat80], Section VIII.4, or [MR92], Section I.3). Moreover, the semigroups $S^{(n)}$ converge strongly on $L_2(\Omega)$ to the submarkovian semigroup S generated by H . It also

follows readily that H is a submarkovian extension of H_Ω . Therefore $H = H_{\Omega,N}(= H_{\Omega,D})$, by Markov uniqueness.

Our next aim is to prove that the semigroups $S^{(n)}$ converge strongly to S on $L_1(\Omega)$. Following a tactic used in [RS08] [RS10b], we convert the L_2 -convergence of the semigroups into L_1 -convergence by the use of suitable off-diagonal bounds.

Step 4: L_2 -off-diagonal bounds. Let

$$D_n = \{ \psi \in W^{1,\infty}(\Omega) : \sum_{i,j=1}^d \rho_n c_{ij} (\partial_i \psi)(\partial_j \psi) \leq 1 \} . \quad (8)$$

The corresponding Riemannian (pseudo-)distance is defined by

$$d_n(x; y) = \sup_{\psi \in D_n} (\psi(x) - \psi(y)) \quad (9)$$

for all $x, y \in \Omega$. This function has the metric properties of a distance but it takes the value infinity if either x or y is not in Ω_n . Secondly, introduce the corresponding set-theoretic distance by

$$d_n(A; B) = \inf_{x \in A, y \in B} d_n(x; y)$$

where A and B are general measurable subsets of Ω . Finally define D by setting $\rho_n = \mathbf{1}_\Omega$ in (8). Then $D \subseteq D_n$ and the corresponding Riemannian distance $d(\cdot; \cdot)$, defined in analogy with (9), satisfies $d(x; y) \leq d_n(x; y)$.

Lemma 3.2 *If A, B are open subsets of Ω then*

$$\sup_{n \geq 1} |(\varphi_A, S_t^{(n)} \varphi_B)| \vee |(\varphi_A, S_t \varphi_B)| \leq e^{-d(A;B)^2(4t)^{-1}} \|\varphi_A\|_2 \|\varphi_B\|_{2S}$$

for all $\varphi_A \in L_2(A)$, $\varphi_B \in L_2(B)$ and $t > 0$ with the convention $e^{-\infty} = 0$.

Bounds of this type have now been derived by many authors (see, for example, [Aus07] [CGT82] [Dav92] [Gri99] [Stu95] [Stu98]) under a variety of ellipticity assumptions. A proof applicable in the current context can be found in [RS08], Section 4. The bounds for $S^{(n)}$ are initially in terms of $d_n(A; B)$ but $d_n(A; B) \leq d(A; B)$. Then since the $S_t^{(n)}$ are L_2 -convergent to S_t the bounds also hold for S .

Next $C(x) \leq c(|x|) I$ for all $x \in \Omega$. Therefore

$$D_n \supseteq \widehat{D}_n = \{ \psi \in W^{1,\infty}(\Omega) : \rho(n^{-1}|x|) c(|x|) |(\nabla \psi)(x)|^2 \leq 1 \} . \quad (10)$$

Consequently

$$\widehat{d}_n(x; y) = \sup_{\psi \in \widehat{D}_n} (\psi(x) - \psi(y)) \leq d_n(x; y) \quad (11)$$

for all $x, y \in \Omega$. Moreover, if \widehat{D} is defined by setting $\rho = \mathbf{1}_\Omega$ in (10) and $\widehat{d}(\cdot; \cdot)$ is defined in analogue with (11) then $\widehat{d}(x; y) \leq d(x; y)$ for all $x, y \in \Omega$. Thus the bounds of Lemma 3.2 are also valid with $d(A; B)$ replaced by $\widehat{d}(A; B)$.

If A, B are bounded open sets with $A \subset \Omega \cap B_m$ and $B \subset \Omega \cap (B_M)^c$ where $M > m \geq 1$ define

$$\rho_m = \sup_{x \in \Omega \cap B_m} \widehat{d}(x; 0) \quad \text{and} \quad \nu_M = \inf_{x \in \Omega \cap (B_M)^c} \widehat{d}(x; 0) .$$

Then it follows from the triangle inequality $\hat{d}(x; 0) \leq \hat{d}(x; y) + \hat{d}(y; 0)$ that

$$\nu_M \leq \inf_{x \in B} \hat{d}(x; 0) \leq \inf_{x \in B} \hat{d}(x; y) + \hat{d}(y; 0) \leq \inf_{x \in B} \hat{d}(x; y) + \rho_m$$

for all $y \in A$. Therefore

$$\hat{d}(A; B) \geq \nu_M - \rho_m \geq 0$$

where the last inequality follows because $M > m$. But it follows directly from the definition of $\hat{d}(\cdot; \cdot)$ that

$$\hat{d}(x; 0) = \int_0^{|x|} ds c(s)^{-1/2}$$

for all $x \in \Omega$. Therefore

$$\rho_m = \int_0^m ds c(s)^{-1/2} = \mu(m) \quad \text{and} \quad \hat{\nu}_M = \int_0^M ds c(s)^{-1/2} = \mu(M).$$

Hence

$$\hat{d}(A; B) \geq \mu(M) - \mu(m) \geq 0.$$

Consequently one has the following variation of Lemma 3.2.

Lemma 3.3 *If $M > m \geq 1$ and A, B are bounded open sets with $A \subset \Omega \cap B_m$ and $B \subset \Omega \cap (B_M)^c$ then*

$$\sup_{n \geq 1} |(\varphi_A, S_t^{(n)} \varphi_B)| \vee |(\varphi_A, S_t \varphi_B)| \leq e^{\mu(m)^2(4t)^{-1}} e^{-\mu(M)^2(8t)^{-1}} \|\varphi_A\|_2 \|\varphi_B\|_2$$

for all $\varphi_A \in L_2(A)$, $\varphi_B \in L_2(B)$ and $t > 0$.

Proof The bounds on $|(\varphi_A, S_t \varphi_B)|$ follow directly from the bounds of Lemma 3.2, the foregoing observation that $\hat{d}(A; B) \geq \mu(M) - \mu(m) \geq 0$ and the estimate

$$(\mu(M) - \mu(m))^2 \geq 2^{-1} \mu(M)^2 - \mu(m)^2.$$

The bounds on $\sup_{n \geq 1} |(\varphi_A, S_t^{(n)} \varphi_B)|$ follow by similar reasoning since $\hat{d}_n(A; B) \geq \hat{d}(A; B)$. We omit further details. \square

Now we are prepared for the key estimate.

Lemma 3.4 *There is a $b > 0$ such that if $M > m \geq 1$ then*

$$\sup_{n \geq 1} |(\mathbf{1}_{(B_M)^c}, S_t^{(n)} \varphi)| \vee |(\mathbf{1}_{(B_M)^c}, S_t \varphi)| \leq b e^{\mu(m)^2(4t)^{-1}} \int_M^\infty ds s^{d/2} e^{-\mu(s)^2(8t)^{-1}} \|\varphi\|_2$$

for all $\varphi \in L_2(\Omega \cap B_m)$ and $t > 0$.

Proof The proof is a variation of an argument of [ERSZ07]. Let $C_p = B_{p+1} \setminus B_p$. It follows that $(B_M)^c = \bigcup_{p \geq M} C_p$. If A is a bounded open set with $\text{supp } \varphi \subset A \subseteq \Omega \cap B_m$ then by Lemma 3.3

$$|(\mathbf{1}_{(B_M)^c}, S_t \varphi)| = \left| \sum_{p \geq M} (\mathbf{1}_{C_p}, S_t \varphi) \right| \leq e^{\mu(m)^2(4t)^{-1}} \sum_{p \geq M} e^{-\mu(p)^2(8t)^{-1}} |B_{p+1}|^{1/2} \|\varphi\|_2.$$

But the sum is a Riemann approximation to the integral occurring in the statement of the lemma. Therefore the bounds for $|(\mathbf{1}_{(B_M)^c}, S_t \varphi)|$ follow immediately. The bounds for $|(\mathbf{1}_{(B_M)^c}, S_t^{(n)} \varphi)|$ follow by similar reasoning. \square

Step 5: L_1 -convergence. The fifth step consists of proving that the semigroups $S_t^{(n)}$ are strongly convergent on $L_1(\Omega)$ to S_t (see [RS08], Proposition 6.2, for a similar result).

Since the semigroups $S_t^{(n)}$ and S_t are all submarkovian it suffices to prove convergence on a subset of $L_1(\Omega)$ whose span is dense. In particular it suffices to prove convergence on positive functions in $L_1(A) \cap L_2(A)$ for each bounded open subset A of Ω .

Fix $A \subset \Omega \cap B_m$ and $\varphi \in L_1(A) \cap L_2(A)$. Assume φ is positive. Then

$$\begin{aligned} \| (S_t^{(n)} - S_t) \varphi \|_1 &\leq \| \mathbf{1}_{B_M} (S_t^{(n)} - S_t) \varphi \|_1 + \| \mathbf{1}_{(B_M)^c} S_t^{(n)} \varphi \|_1 + \| \mathbf{1}_{(B_M)^c} S_t \varphi \|_1 \\ &\leq |B_M|^{1/2} \| (S_t^{(n)} - S_t) \varphi \|_2 + (\mathbf{1}_{(B_M)^c}, S_t^{(n)} \varphi) + (\mathbf{1}_{(B_M)^c}, S_t \varphi) \\ &\leq |B_M|^{1/2} \| (S_t^{(n)} - S_t) \varphi \|_2 + 2b e^{\mu(m)^2(4t)^{-1}} \int_M^\infty ds s^{d/2} e^{-\mu(s)^2(8t)^{-1}} \| \varphi \|_2 \end{aligned}$$

where we have used the positivity of the semigroups and the functions to express the L_1 -norms as pairings between L_1 and L_∞ . The last step uses Lemma 3.4. But the integral is convergent for one $t = t_0 > 0$, by assumption. Therefore it is convergent for all $t \in \langle 0, t_0 \rangle$. Then since $S_t^{(n)}$ is L_2 -convergent to S_t for all $t > 0$ and since the last term on the right hand side converges to zero as $M \rightarrow \infty$ for each $t \in \langle 0, t_0 \rangle$ it follows that $S_t^{(n)}$ is L_1 -convergent to S_t for all $t \in \langle 0, t_0 \rangle$. Finally it follows from the semigroup property and contractivity that $S_t^{(n)}$ is L_1 -convergent to S_t for all $t > 0$.

Step 6: Conservation. The conservation property for S now follows because the approximating semigroups $S^{(n)}$ are conservative, by Step 2, and are L_1 -convergent to S , by Step 5. Therefore

$$(\mathbf{1}_\Omega, S_t \varphi) = \lim_{n \rightarrow \infty} (\mathbf{1}_\Omega, S_t^{(n)} \varphi) = \lim_{n \rightarrow \infty} (S_t^{(n)} \mathbf{1}_\Omega, \varphi) = (\mathbf{1}_\Omega, \varphi)$$

for all $\varphi \in L_1(\Omega)$. Hence $S_t \mathbf{1}_\Omega = \mathbf{1}_\Omega$. \square

This completes the proof of Proposition 3.1 and the third statement of Theorem 1.1.

4 Illustrations and examples

In this section we illustrate the foregoing results with some applications and examples.

Theorem 1.1 established that L_1 -uniqueness of $H_\Omega \in \mathcal{E}_\Omega$ is a consequence of two distinct properties, a capacity condition on the boundary and a growth condition on the coefficients. Therefore we separate the initial discussion into two parts each concentrating one of these conditions.

4.1 Growth properties

If $\Omega = \mathbf{R}^d$ then the capacity condition plays no role and so we begin by considering this case. We continue to use the function c and the corresponding function μ defined by (3) and (4), respectively, as measures of the coefficient growth. The following statement combines the L_1 -properties which follow from the foregoing with the comparable L_2 -properties established earlier by Davies et al. (see [Dav85] and references therein).

Proposition 4.1 *Let $H \in \mathcal{E}_{\mathbf{R}^d}$. Then the following are valid:*

- I. *If $\mu(s) \rightarrow \infty$ as $s \rightarrow \infty$ then H is L_2 -unique,*
- II. *If $\int_0^\infty ds s^{d/2} e^{-\lambda \mu(s)^2} < \infty$ for one $\lambda > 0$ then H is L_1 -unique.*

The second statement is a direct consequence of the second statement of Theorem 1.1 since one automatically has $\text{cap}_\Omega(\partial\Omega) = 0$. The first statement follows from [Dav85], Theorem 3.2. This theorem asserts that H is essentially self-adjoint, i.e. L_2 -unique, if there exists a strictly positive differentiable function η over \mathbf{R}^d with $\|\Gamma(\eta)\|_\infty < \infty$ such that $\eta(x) \rightarrow \infty$ as $x \rightarrow \infty$. But it follows by assumption that $\eta(x) = \int_0^{|x|} dt c(t)^{-1/2}$ satisfies these properties.

Next we consider a special case of the growth property examined earlier by Davies [Dav85] and Eberle [Ebe99].

Example 4.2 Assume that $c(s) \leq a s^2 (\log s)^\alpha$ for some $a > 0$, $\alpha \geq 0$ and all large s . In this case $\mu(s) \geq b (\log s)^{1-\alpha/2}$ with $b > 0$ for all large s . Therefore $\mu(s) \rightarrow \infty$ as $s \rightarrow \infty$ if $\alpha < 2$ and H is L_2 -unique by the first statement of Proposition 4.1. But Davies has demonstrated by specific example, [Dav85] Example 3.5, that if $d \geq 2$ and $\alpha > 2$ then L_2 -uniqueness can fail (see also [Ebe99] Chapter 2, Section c). Finally Theorem 2.3 in [Ebe99] treats the borderline case $\alpha = 2$. This theorem establishes that if $c(s) \leq a s^2 (\log s)^2$ for all large s then H is not only L_2 -unique but also L_p -unique for all $p \in \langle 1, 2 \rangle$.

Next if $\alpha \leq 1$ then the second statement of Proposition 4.1 establishes that the bound H is L_1 -unique. Indeed if $\alpha \leq 1$ then $\mu(s) \geq b (\log s)^{1-\alpha/2}$ for s large and the integral (6) is finite for large λ . Therefore Proposition 4.1 establishes that H is L_1 -unique. But L_1 -uniqueness can fail if $c(s) \sim s^2 (\log s)^\alpha$ with $\alpha > 1$ for large s . To verify this let $d = 1$ and define the positive L_∞ -function ψ on \mathbf{R} by

$$\psi(x) = 1 - a (\log(\log 2)) (\log(\log(2 + |x|)))^{-1} \quad (12)$$

with $a \in \langle 0, 1 \rangle$. Thus $\psi(0) = 1 - a > 0$ and $\psi(x) \rightarrow 1$ as $|x| \rightarrow \infty$. Then define c by $c(x) = \psi'(x)^{-1} \int_0^x ds \psi(s)$. It is evident that c is strictly positive and $c \in W_{\text{loc}}^{1,\infty}(\mathbf{R})$. Moreover, $c(x) \sim |x|^2 (\log |x|) (\log(\log |x|))$ as $|x| \rightarrow \infty$. But if H is the corresponding operator on $C_c^\infty(\mathbf{R})$, i.e. if $H\varphi = -(c\varphi)'$, then $(I + H)\psi = \psi - (c\psi)' = 0$. Therefore the range of $I + H$ is not L_1 -dense and H is not L_1 -unique.

Therefore within this class of examples the growth bound $c(s) \leq a s^2 (\log s)$ is optimal for L_1 -uniqueness and the bound $c(s) \leq a s^2 (\log s)^2$ is optimal for L_2 -uniqueness. \square

Note that if $d = 1$, $\Omega = \langle 0, \infty \rangle$ and one repeats the foregoing construction with ψ given by (12) but with $a = 1$ then c is strictly positive and $c(x) = O(x)$ as $x \rightarrow 0$. Moreover, $c(x) \sim x^2 (\log x) (\log(\log x))$ for all large x . Therefore the corresponding operator H is Markov unique, by [RS10a] Theorem 2.7, but again it is not L_1 -unique. In fact it is not L_p -unique for any $p \in [1, \infty)$.

The function μ is a lower bound on the Riemmanian distance to infinity measured with respect to the metric C^{-1} associated with the operator H_Ω . If one has more detailed information on the geometry one can obtain stronger conclusion by the same general reasoning. This is illustrated by the following example of a Grušin-type operator.

Let $d = 2$ and $\Omega = \Omega_+ \cup \Omega_-$ with $\Omega_{\pm} = \{x = (x_1, x_2) : \pm x_1 > 0\}$. Define the Grušin operator H by $D(H) = C_c^\infty(\Omega)$ and

$$(H\varphi)(x) = -\partial_1(c_1(x_1)\partial_1\varphi)(x) - c_2(x_1)(\partial_2^2\varphi)(x)$$

where $c_1, c_2 \in W_{\text{loc}}^{1,\infty}(\mathbf{R} \setminus \{0\})$ are strictly positive and $c_i(x) \sim |x|^{(2\delta_i, 2\delta'_i)}$ with $\delta_i, \delta'_i \geq 0$. Here we use the notation of [RS06] [RS08]. Specifically $s^{(\alpha, \alpha')} = s^\alpha$ if $\alpha \leq 1$ and $s^{(\alpha, \alpha')} = s^{\alpha'}$ if $\alpha \geq 1$ and functions f, g satisfy the relation $f \sim g$ if there are $a, a' > 0$ such that $a f \leq g \leq a' f$. We assume that $\delta_1, \delta'_1 < 1$ but there are no upper bounds on δ_2 and δ'_2 . Thus $H \in \mathcal{E}_\Omega$ with

$$C(x) = \begin{pmatrix} c_1(x_1) & 0 \\ 0 & c_2(x_1) \end{pmatrix}.$$

Therefore $\|C(x)\| = c_1(x_1) \vee c_2(x_1) \leq a |x_1|^{2(\delta'_1 \vee \delta'_2)} \leq a |x_1|^{2(1 \vee \delta'_2)}$ for all $|x_1| \geq 1$. Although the asymptotic growth of C is dictated by c_2 , which behaves asymptotically like $|x_1|^{2\delta'_2}$, the uniqueness properties are independent of the magnitude of δ'_2 .

Proposition 4.3 *Let H denote the Grušin operator defined above.*

- I. *If $\delta_1 \in [0, 1/2)$ then H is not Markov unique and consequently not L_1 -unique.*
- II. *If $\delta_1 \in [1/2, 1)$ then H is L_1 -unique and consequently Markov unique.*

Proof The first statement of the proposition follows from the observations at the end of Section 6 in [RS08] and in particular from Proposition 6.10.

Some care has to be taken in comparing the current statements with those of [RS08]. The operator H is defined on $C_c^\infty(\mathbf{R}^2 \setminus \{x_1 = 0\})$ but the operator H_δ studied in [RS08] corresponds to the extension of H to $C_c^\infty(\mathbf{R}^2)$. The Friedrichs' extension $H_{\delta, D}$ of H_δ is the self-adjoint extension H_N of H which satisfies the Neumann-type boundary condition $(c_1\partial_1)(0_+, x_2) = (c_1\partial_1)(0_-, x_2)$ on the line of degeneracy $x_1 = 0$. The Friedrichs' extension H_D of H is, however, the self-adjoint extension with the Dirichlet-type boundary condition $\varphi(0_+, x_2) = \varphi(0_-, x_2)$. If $\delta_1 \in [0, 1/2)$ then these extensions are distinct and, in addition there are extensions with analogous Robin boundary conditions sandwiched between the minimal extension $H_N (= H_{\delta, D})$ and the maximal extension H_D . But if $\delta_1 \in [1/2, 1)$ then $H_N = H_D$ and all the operators coincide (see [RS08], Proposition 6.10).

The proof of L_1 -uniqueness for $\delta_1 \in [1/2, 1)$ is by reasoning similar to that used to prove Proposition 3.1 and it does not require an upper bound on δ'_2 . The argument follows the lines of the proof of Theorem 6.1 in [RS08], details of which are given in [RS06]. First, Markov uniqueness follows from Proposition 6.10 of [RS08]. Secondly, one deduces that H is conservative by the arguments given in [RS06]. The semigroup S generated by $H_N (= H_D)$ is approximated on $L_2(\Omega)$ by semigroups $S^{(N, \varepsilon)}$ generated by the Grušin operators with coefficients $(C \wedge NI) + \varepsilon I$. Then S and $S^{N, \varepsilon}$ satisfy L_2 -off-diagonal bounds with respect to the corresponding Riemannian distances by [RS08], Proposition 4.1. But if $N \geq 1 \geq \varepsilon > 0$ then these distances are all larger than the Riemannian distance $d_1(\cdot; \cdot)$ corresponding to the Grušin operator with coefficients $(c_1 + 1, c_2 + 1)$. Therefore S and the approximants $S^{(N, \varepsilon)}$ all satisfy L_2 -off-diagonal bounds with respect to $d_1(\cdot; \cdot)$. Since the operator with coefficients $(c_1 + 1, c_2 + 1)$ has $\delta_1 = 0 = \delta_2$ it follows that $d_1(\cdot; \cdot)$ is independent of δ_1 and δ_2 .

Next let $B_{1,r} = \{x \in \mathbf{R}^2 : d_1(0; x) < r\}$. Then if $\varphi \in L_2(B_{1,m})$ it follows by L_2 -off-diagonal bounds, similar to those of Lemma 3.2, as in the proof of Lemma 3.4 that

$$|(\mathbf{1}_{(B_M)^c}, S_t^{(N,\varepsilon)}\varphi)| \vee |(\mathbf{1}_{(B_M)^c}, S_t\varphi)| \leq \sum_{p \geq M} |B_{1,p+1}|^{1/2} e^{-d_1(C_p; B_{1,m})^2(4t)^{-1}} \|\varphi\|_2$$

where $C_p = B_{1,p+1} \setminus B_{1,p}$. But $d_1(C_p; B_{1,m}) \geq p - m$ by the triangle inequality. Moreover, it follows from Proposition 5.1 of [RS08] that there is an $a > 0$ such that $|B_{1,p}| \leq a^2 p^{D'}$ with $D' = 1 + (1 + \delta'_2 - \delta'_1)(1 - \delta'_1)^{-1}$. Therefore

$$|(\mathbf{1}_{(B_M)^c}, S_t^{(N,\varepsilon)}\varphi)| \vee |(\mathbf{1}_{(B_M)^c}, S_t\varphi)| \leq a \sum_{p \geq M} p^{D'/2} e^{-(p-m)^2(4t)^{-1}} \|\varphi\|_2$$

and the estimate is uniform for all $N \geq 1$ and $\varepsilon \in \langle 0, 1 \rangle$. Finally the $S^{(N,\varepsilon)}$ are conservative, since their generators are strongly elliptic, and they are L_2 -convergent to S . But

$$\|(S_t^{(N,\varepsilon)} - S_t)\varphi\|_1 \leq |B_{1,M}|^{1/2} \|(S_t^{(N,\varepsilon)} - S_t)\varphi\|_2 + (\mathbf{1}_{(B_M)^c}, S_t^{(N,\varepsilon)}\varphi) + (\mathbf{1}_{(B_M)^c}, S_t\varphi)$$

for all positive $\varphi \in L_2(B_{1,m})$. Therefore taking the limits $N \rightarrow \infty$, $\varepsilon \rightarrow 0$ and $M \rightarrow \infty$ one deduces that the $S^{(N,\varepsilon)}$ are L_1 -convergent to S . Hence S is conservative and H is L_1 -unique by Theorem 1.1. \square

4.2 Capacity estimates

In this subsection we suppose that Ω is a strict subset of \mathbf{R}^d and examine the capacity condition $\text{cap}_\Omega(\partial\Omega) = 0$. It follows from the general properties of the capacity [BH91] or [FOT94] that $\text{cap}_\Omega(\partial\Omega) = 0$ if and only if $\text{cap}_\Omega(A) = 0$ for all bounded measurable subsets A of $\partial\Omega$. Moreover, $\text{cap}_\Omega(A) = 0$ implies $|A| = 0$. The capacity $\text{cap}_\Omega(A)$ depends on two gross features of A and H_Ω , the dimension of the set and the order of degeneracy of H_Ω at A .

First let $A \subset \Omega$ be a bounded measurable subset with $|A| = 0$ and let $\dim_H(A)$ and $\dim_M(A)$ denote the Hausdorff and Minkowski dimensions of the set, respectively. If A is a general measurable subset the corresponding dimensions are defined by $\dim_H(A) = \sup\{\dim_H(B) : B \subset A\}$ and $\dim_M(A) = \sup\{\dim_M(B) : B \subset A\}$ where the suprema are over all bounded measurable subsets B . In general $\dim_H(A) \leq \dim_M(A)$.

Secondly, the order of degeneracy of the operator $H_\Omega \in \mathcal{E}_\Omega$ on the bounded set A is defined to be the largest $\gamma_\Omega(A) \geq 0$ for which there is an open subset U containing A and an $a > 0$ such that $0 < C(x) \leq a d(x; A)^{\gamma_\Omega(A)} I$ for all $x \in U \cap \Omega$ where $d(x; A)$ denotes the Euclidean distance of x from A . Again the order of degeneracy of an unbounded set is defined as a supremum over bounded subsets.

It follows from the proof of Proposition 4.3 in [RS10b] that one has the following property.

Lemma 4.4 *Let $A \subset \overline{\Omega}$ be a bounded measurable subset with $|A| = 0$. Assume $H_\Omega \in \mathcal{E}_\Omega$ is degenerate of order $\gamma_\Omega(A)$ on A .*

If $d - 2 \geq \dim_M(A) - \gamma_\Omega(A)$ then $\text{cap}_\Omega(A) = 0$. In particular if $\gamma_\Omega(A) \geq 2$ then $\text{cap}_\Omega(A) = 0$.

The proof of the first statement is similar to the proof of Proposition 4.3 in [RS10b] because the estimates in the proof of the latter proposition are all local. Any possible

growth of the coefficients plays no role. The second statement follows immediately since $\dim_M(A) < d$.

The lemma is not optimal and we next establish a similar but stronger statement involving the Hausdorff dimension.

Proposition 4.5 *Let $A \subset \overline{\Omega}$ be a bounded measurable subset with $|A| = 0$. Assume $H_\Omega \in \mathcal{E}_\Omega$ is degenerate of order $\gamma_\Omega(A)$ on A .*

If $d - 2 \geq \dim_H(A) - \gamma_\Omega(A)$ then $\text{cap}_\Omega(A) = 0$. In particular if $d \geq 2$ and $\dim_H(A) \leq d - 2$ then $\text{cap}_\Omega(A) = 0$.

Proof First, if $\gamma_\Omega(A) \geq 2$ then $\text{cap}_\Omega(A)$ by Lemma 4.4. Therefore we assume $\gamma_\Omega(A) < 2$.

Secondly, let $B_r = B(y; r)$ be a Euclidean ball of radius r centred at y and assume $A \cap B_r \neq \emptyset$. Then there is an $a > 0$ such that $d(x; A) \leq ar$ for all $x \in B_{2r}$ uniformly in y . Therefore $\|C(x)\| \leq br^{\gamma(A)}$ for all $x \in B_{2r} \cap \Omega$ with $b = ar^{\gamma(A)}$ independent of y . Next fix $\eta_r \in C_c^\infty(B_{2r})$ with $0 \leq \eta_r \leq 1$ and $\eta_r = 1$ on B_r . One may assume $|\nabla \eta_r| \leq 2r^{-1}$ on $B_{2r} \setminus B_r$. Then

$$\text{cap}_\Omega(B_r \cap \Omega) \leq \int_{B_{2r} \cap \Omega} \left(\|C(x)\| |\nabla \eta_r(x)|^2 + |\eta_r|^2 \right) \leq |B_{2r}| (4br^{\gamma(A)-2} + 1)$$

Since $\gamma(A) < 2$ it follows that there is a $c > 0$ such that $\text{cap}_\Omega(B_r \cap \Omega) \leq cr^{d+\gamma(A)-2}$ for all $r \leq 1$. Again the estimate is uniform in y , the centre point of B_r .

Thirdly, let $B_{r_i} = B(y_i; r_i)$ be a countable family of balls with $r_i \leq \delta \leq 1$ such that $B_{r_i} \cap A \neq \emptyset$ and $A \subset \bigcup_i B_{r_i}$. Then

$$\text{cap}_\Omega(A) \leq \sum_i \text{cap}_\Omega(B_{r_i} \cap \Omega) \leq c \sum_i r_i^{\gamma(A)+d-2}$$

by the foregoing estimate. Therefore

$$\text{cap}_\Omega(A) \leq c H_{\gamma(A)+d-2}(A)$$

where H_s is the Hausdorff measure. It follows immediately that $\text{cap}_\Omega(A) = 0$ if $\gamma(A) + d - 2 \geq \dim_H(A)$ or, equivalently, if $d - 2 \geq \dim_H(A) - \gamma(A)$. Finally since $\gamma(A) \geq 0$ it follows that $\text{cap}_\Omega(A) = 0$ whenever $d \geq 2$ and $\dim_H(A) \leq d - 2$. \square

Combination of Theorem 1.1 and Lemma 4.4 immediately gives the following criterion for L_1 -uniqueness.

Proposition 4.6 *If the growth condition (6) is satisfied and if $d - 2 \geq \dim_H(A) - \gamma_\Omega(A)$ for each bounded measurable subset $A \subset \partial\Omega$ with $|A| = 0$ then H_Ω is L_1 -unique.*

Lemma 4.4 and Proposition 4.5 give a quantitative assessment of the two distinct effects which lead to zero capacity. First if the operator H_Ω has a second-order degeneracy on the set A , i.e. if $\gamma_\Omega(A) \geq 2$, then $\text{cap}_\Omega(A) = 0$ independently of the dimension of A . Secondly, if $\dim_H(A) \leq d - 2$ then $\text{cap}_\Omega(A) = 0$ independently of the order of degeneracy of H_Ω on A . Alternatively, if $\dim_H(A) \leq d - 1$ then a first-order degeneracy, $\gamma_\Omega(A) \geq 1$, is sufficient to ensure that $\text{cap}_\Omega(A) = 0$. If, for example, A is Lipschitz continuous then $\dim_H(A) = d - 1$ and this last case is applicable.

In the first of the foregoing cases one even has a simple criterion for L_2 -uniqueness.

Lemma 4.7 *Assume $|\partial\Omega| = 0$. Let $H_\Omega \in \mathcal{E}_\Omega$. If $\gamma_\Omega(A) \geq 2$ for all bounded measurable $A \subset \partial\Omega$ and $\mu(s) \rightarrow \infty$ as $s \rightarrow \infty$ then H is L_2 -unique.*

Proof Let ρ_n denote the functions introduced in the proof of IV \Rightarrow III in Theorem 1.1. Since $\mu(s) \rightarrow \infty$ as $s \rightarrow \infty$ it follows that ρ_n converges pointwise to $\mathbf{1}_\Omega$ as $n \rightarrow \infty$. Moreover, $\|\Gamma(\rho_n)\|_\infty \leq b^2 n^{-2}$. Next define χ_n on $[0, \infty)$ by $\chi_n(s) = 1$ if $s \in [0, n^{-1})$, $\chi_n(s) = -\log s / \log n$ if $s \in [n^{-1}, 1]$ and $\chi_n(s) = 0$ if $s \geq 1$. Then define ξ_n on Ω by $\xi_n(x) = \chi_n(d(x; \partial\Omega))$. Finally define η_n by $\eta_n = \rho_n (\mathbf{1}_\Omega - \xi_n)$. It follows that η_n converges pointwise to $\mathbf{1}_\Omega$ as $n \rightarrow \infty$. In addition

$$\|\Gamma(\eta_n)\|_\infty \leq 2 \|\Gamma(\rho_n)\|_\infty + 2a \|\Gamma(\xi_n)\|_\infty \leq 2b^2 n^{-2} + 2a (\log n)^{-2} \rightarrow 0$$

as $n \rightarrow \infty$ where we have used $\|\Gamma(\xi_n)\|_\infty \leq a (\log n)^{-2}$. The latter estimate follows from the degeneracy assumption. Hence H_Ω is L_2 -unique by Proposition 6.1 of [RS10b], with $p = 2$, and a standard regularization argument. \square

Note that in Lemma 4.7 there is no restraint on the dimension of the boundary $\partial\Omega$. Therefore the conclusion is valid for sets Ω with arbitrarily rough boundaries, in particular for fractal boundaries. The result is, however, not optimal if the boundary is smooth. Indeed if $d = 1$ and $\Omega = \langle 0, \infty \rangle$ then a degeneracy of order 1 at the origin is necessary and sufficient for L_1 -uniqueness and a degeneracy of order $3/2$ is necessary and sufficient for L_2 -uniqueness (see, [RS10a], Theorem 2.7).

4.3 Negligible sets

The foregoing discussion indicates that sets of Hausdorff dimension lower than $d - 2$ are insignificant for L_1 -uniqueness. In this subsection we verify that this is indeed the case for non-degenerate operators and also establish that sets with Hausdorff dimension less than $d - 4$ are negligible for L_2 -uniqueness.

Proposition 4.8 *Assume $H \in \mathcal{E}_{\mathbf{R}^d}$ with $d \geq 2$ and let $\Gamma \subset \mathbf{R}^d$ be a closed subset with $|\Gamma| = 0$. Further assume the growth condition $\int_0^\infty ds s^{d/2} e^{-\lambda \mu(s)^2} < \infty$ for one $\lambda > 0$.*

The following conditions are equivalent:

- I. $\dim_H(\Gamma) \leq d - 2$,
- II. $C_c^\infty(\mathbf{R}^d \setminus \Gamma)$ is an L_1 -core of H .

Proof The proof is based on the observation that both conditions of the proposition are equivalent to the capacity of the set Γ being zero. But there are three different capacities involved in the argument.

Let $A \subset \Gamma$ be a bounded measurable subset. Then we define $\text{cap}(A)$ as the capacity of the set measured with respect to H . Explicitly

$$\text{cap}(A) = \inf \left\{ \|\psi\|_{D(h)}^2 : \psi \in D(h) \text{ and there exists an open set } U \subset \mathbf{R}^d \text{ such that } U \supseteq A \text{ and } \psi \geq 1 \text{ a. e. on } U \right\} \quad (13)$$

with h the closed quadratic form associated with H . Next let $\Omega = \mathbf{R}^d \setminus \Gamma$. Set $\mathcal{D} = C_c^\infty(\mathbf{R}^d \setminus \Gamma)$ and $H_\Omega = H|_{\mathcal{D}}$. Then $H_\Omega \in \mathcal{E}_\Omega$. Let $\text{cap}_\Omega(A)$ denote the capacity measured with respect to H_Ω . Thus $\text{cap}_\Omega(A)$ is given by (13) with h replaced by $h_{\Omega, N}$. Since $h_{\Omega, N} \supseteq h$

it follows that $\text{cap}_\Omega(A) \leq \text{cap}(A)$. But both capacities can be calculated with functions which are equal to one in an open neighbourhood of A and on such functions the two forms coincide. Therefore $\text{cap}_\Omega(A) = \text{cap}(A)$.

Next define $\text{cap}_{1,2}(A)$ as the capacity of the set A measured with respect to the Laplacian. This latter capacity is defined by (13) but $D(h)$ is replaced by $W^{1,2}(\mathbf{R}^d)$. Now assume $\text{cap}(A) = 0$. Then $C_c^\infty(\mathbf{R}^d)$ is a core of h , by definition. Therefore there exist a sequence $\chi_n \in C_c^\infty(\mathbf{R}^d)$ and a decreasing sequence of bounded open subsets $U_n \supset A$ such that $0 \leq \chi_n \leq 1$, $\chi_n = 1$ on U_n and $h(\chi_n) + \|\chi_n\|_2^2 \leq n^{-1}$. Then fix an $\eta \in C_c^\infty(\mathbf{R}^d)$ such that $0 \leq \eta \leq 1$ and $\eta = 1$ on U_1 and hence on each U_n . Set $\varphi_n = \chi_n \eta$. It follows that $\varphi_n \in D(h)$, $0 \leq \varphi_n \leq 1$, $\varphi_n = 1$ on U_n and $h(\varphi_n) + \|\varphi_n\|_2^2 \leq a n^{-1}$ with $a = 2(\|\nabla \eta\|_\infty^2 + 1)$. Moreover, $\text{supp } \varphi_n \subseteq K$ for all n . But it follows from strict positivity of the matrix of coefficients C that there exists a $\mu_K > 0$ such that

$$\|\varphi\|_{D(h)}^2 \geq \mu_K \|\varphi\|_{W^{1,2}(\mathbf{R}^d)}^2$$

for all $\varphi \in W^{1,2}(K)$. Therefore $\text{cap}_{1,2}(A) = 0$.

After these preliminaries we turn to the proof of equivalence of the conditions of the proposition. We prove that both conditions are equivalent to $\text{cap}(\Gamma) = 0$.

I \Leftrightarrow $\text{cap}(\Gamma) = 0$. It suffices to prove that Condition I is equivalent to $\text{cap}(A) = 0$ for all bounded measurable subsets A of Γ . But Proposition 4.5 establishes that Condition I implies $\text{cap}_\Omega(A) = 0$ or, equivalently, $\text{cap}(A) = 0$. Conversely, $\text{cap}(A) = 0$ implies $\text{cap}_{1,2}(A) = 0$ by the foregoing discussion. But $\text{cap}_{1,2}(A) = 0$ implies $\dim_H(\Gamma) \leq d - 2$ by standard properties of the Laplacian (see, for example, Theorem 2.26 of [HKM93], Corollary 5.1.15 of [AH96] or Section 2.1.7 of [MZ97]).

II \Leftrightarrow $\text{cap}(\Gamma) = 0$. First, observe that H is L_1 -unique by the second statement of Proposition 4.1. But H is L_1 -unique if and only if \overline{H}^1 is the generator of an L_1 -continuous semigroup. Now suppose Condition II is valid. Then $\overline{H}^1 = \overline{H}_\Omega^1$. Therefore \overline{H}_Ω^1 is the generator of an L_1 -continuous semigroup. Consequently, H_Ω is L_1 -unique. But L_1 -uniqueness of H_Ω is equivalent to $\text{cap}_\Omega(\Gamma) = 0$ by Theorem 1.1 which in turn is equivalent to $\text{cap}(\Gamma) = 0$.

Conversely, if H_Ω is L_1 -unique then \overline{H}_Ω^1 is the generator of an L_1 -continuous semigroup. But \overline{H}^1 is also a generator and $\overline{H}^1 \supseteq \overline{H}_\Omega^1$ by definition. Since a semigroup generator cannot have a proper generator extension one must have $\overline{H}^1 = \overline{H}_\Omega^1 = \overline{H}|_{\mathcal{D}}$. Thus \mathcal{D} is an L_1 -core of H . \square

It follows from the assumptions of Proposition 4.8 and the first statement of Proposition 4.1 that H is L_2 -unique, i.e. H is essentially self-adjoint. Then the closed form h associated with H is the form of the L_2 -closure \overline{H} of H . Therefore $h(\varphi) = \|\overline{H}^{1/2} \varphi\|_2^2$ for all $\varphi \in D(h) = D(\overline{H}^{1/2})$. This observation provides a relation between L_1 - and L_2 -cores.

Corollary 4.9 *Adopt the assumptions of Proposition 4.8 .*

The following conditions are equivalent:

- I. $C_c^\infty(\mathbf{R}^d \setminus \Gamma)$ is an L_1 -core of H ,
- II. $C_c^\infty(\mathbf{R}^d \setminus \Gamma)$ is an L_2 -core of $\overline{H}^{1/2}$.

Proof Again set $\Omega = \mathbf{R}^d \setminus \Gamma$, $\mathcal{D} = C_c^\infty(\mathbf{R}^d \setminus \Gamma)$ and $H_\Omega = H|_{\mathcal{D}}$. Then $H_\Omega \in \mathcal{E}_\Omega$. Moreover, it follows from the proof of Proposition 4.8 that Condition I is equivalent to the condition

$\text{cap}_\Omega(\Gamma) = 0$. But Condition II is also equivalent to this capacity condition by the proof of Theorem 1.1. \square

There is also an L_2 -version of Proposition 4.8.

Proposition 4.10 *Assume $H \in \mathcal{E}_{\mathbf{R}^d}$ with $d \geq 4$ and let $\Gamma \subset \mathbf{R}^d$ be a closed subset with $|\Gamma| = 0$. Further assume that $\mu(s) \rightarrow \infty$ as $s \rightarrow \infty$.*

The following conditions are equivalent:

- I. $\dim_H(\Gamma) \leq d - 4$,
- II. $C_c^\infty(\mathbf{R}^d \setminus \Gamma)$ is an L_2 -core of H .

Proof It follows from the first statement of Proposition 4.1 that H is essentially self-adjoint. Then if $A \subset \mathbf{R}^d$ is a measurable subset we define the capacity $\text{Cap}(A)$ associated with the self-adjoint L_2 -closure \overline{H} by

$$\text{Cap}(A) = \inf \left\{ \|\psi\|_{D(\overline{H})}^2 : \psi \in D(\overline{H}) \text{ and there exists an open set } U \text{ such that } U \supseteq A \text{ and } \psi \geq 1 \text{ a. e. on } U \right\}. \quad (14)$$

We will argue that both conditions of the proposition are equivalent to $\text{Cap}(\Gamma) = 0$ or, equivalently, $\text{Cap}(A) = 0$ for all bounded measurable subsets A of Γ .

If A is bounded and $\text{Cap}(A) = 0$ then there exist a sequence $\chi_n \in C_c^\infty(\mathbf{R}^d)$ and a decreasing sequence of bounded open subsets $U_n \supset A$ such that $0 \leq \chi_n \leq 1$, $\chi_n = 1$ on U_n and $\|\chi_n\|_{D(\overline{H})}^2 \leq n^{-1}$. But the sequence χ_n can be modified by a variation of the argument used in the proof of Proposition 4.8 to yield a sequence with similar properties but with each element of the sequence supported by a fixed compact set. Explicitly, fix $\eta \in C_c^\infty(\mathbf{R}^d)$ such that $0 \leq \eta \leq 1$ and $\eta = 1$ on U_1 and hence on each U_n . Then set $\varphi_n = \chi_n \eta$. It follows that $0 \leq \varphi_n \leq 1$, $\varphi_n = 1$ on U_n and $\|\varphi_n\|_2 \leq \|\chi_n\|_2$. But

$$\overline{H}\varphi_n = (\overline{H}\chi_n)\eta + \chi_n(\overline{H}\eta) + 2\Gamma(\chi_n, \eta)$$

where $\Gamma(\cdot; \cdot)$ is the *carré du champ* associated with H . Therefore

$$\|\overline{H}\varphi_n\|_2^2 \leq 3(\|\overline{H}\eta\|_\infty^2 + \|\eta\|_\infty^2) \|\chi_n\|_{D(\overline{H})}^2 + 12 \int \Gamma(\chi_n; \eta)^2.$$

But

$$\int \Gamma(\chi_n; \eta)^2 \leq \int \Gamma(\chi_n) \Gamma(\eta) \leq |\Gamma(\eta)|_\infty h(\chi_n) \leq 2^{-1} |\Gamma(\eta)|_\infty \|\chi_n\|_{D(\overline{H})}^2.$$

Combining these estimates one deduces that there is an $a > 0$ such that $\|\varphi_n\|_{D(\overline{H})}^2 \leq a \|\chi_n\|_{D(\overline{H})}^2 \leq a n^{-1}$ for all n .

Next since the coefficients $c_{ij} \in W_{\text{loc}}^{1,\infty}(\Omega)$ and $C = (c_{ij}) > 0$ it follows that there exists a $\mu_K > 0$ such that

$$\|\varphi\|_{D(\overline{H})}^2 \geq \mu_K \|\varphi\|_{W^{2,2}(\mathbf{R}^d)}^2$$

for all $\varphi \in C_c^\infty(K)$ (see, for example, the appendix of [RS10b]). Therefore replacing φ by φ_n and taking the limit $n \rightarrow \infty$ one deduces that $\text{cap}_{2,2}(A) = 0$ where $\text{cap}_{2,2}$ is the capacity measured with respect to the $W^{2,2}(\mathbf{R}^d)$ -norm, i.e. $\text{cap}_{2,2}(A)$ is given by (14) but with $D(\overline{H})$ replaced by $W^{2,2}(\mathbf{R}^d)$.

$I \Leftrightarrow \text{Cap}(\Gamma) = 0$. It suffices to prove that Condition I is equivalent to $\text{Cap}(A) = 0$ for all bounded measurable subsets A of Γ . But a slight variation of the proof of Proposition 4.5 establishes that Condition I implies $\text{Cap}(A) = 0$. Indeed define B_r and η_r as in the proof of Proposition 4.5. One may also assume that $|\Delta\eta_r| \leq ar^{-2}$ on $B_{2r} \setminus B_r$. Then one estimates straightforwardly that there are $b, b' > 0$ such that

$$\text{Cap}(B_r) \leq b \int_{B_{2r}} \left(|\Delta\eta_r|^2 + |\eta_r|^2 \right) \leq b' (r^{d-4} + r^d) \leq 2b' r^{d-4}$$

for all $r \leq 1$. The rest of the proof remains unchanged.

Conversely, if $\text{Cap}(A) = 0$ then $\text{cap}_{2,2}(A) = 0$ by the foregoing discussion. But then $\dim_H(A) \leq d - 4$ by another application of Theorem 2.26 of [HKM93].

$II \Leftrightarrow \text{Cap}(\Gamma) = 0$. First suppose Condition II is valid. Secondly, fix $\psi \in C_c^\infty(\mathbf{R}^d) \subset D(H)$ with $\psi = 1$ on an open neighbourhood of $A \subset \Gamma$. Then, by II, there is a sequence $\psi_n \in C_c^\infty(\mathbf{R}^d \setminus \Gamma)$ such that $\|\psi - \psi_n\|_{D(\overline{H})} \rightarrow 0$ as $n \rightarrow \infty$. Set $\varphi_n = \psi - \psi_n$. It follows that $\varphi_n \in D(H)$, $\varphi_n = 1$ on an open neighbourhood U_n of A and $\|\varphi_n\|_{D(\overline{H})} \rightarrow 0$ as $n \rightarrow \infty$. Therefore $\text{Cap}(A) = 0$. Since this holds for an arbitrary bounded subset A of Γ it follows that $\text{Cap}(\Gamma) = 0$.

Conversely, suppose $\text{Cap}(\Gamma) = 0$. Therefore $\text{Cap}(A) = 0$ for each bounded measurable subset A of Γ . Then since $C_c^\infty(\mathbf{R}^d)$ is a core of H , by definition, there exist a sequence $\chi_n \in C_c^\infty(\mathbf{R}^d)$ and a sequence of open subsets U_n of A such that $\|\chi_n\|_{D(\overline{H})} \rightarrow 0$ as $n \rightarrow \infty$. Now fix $\psi \in C_c^\infty(\mathbf{R}^d)$ and set $\psi_n = (\mathbf{1} - \chi_n)\psi$. It follows that $\psi_n \in C_c^\infty(\mathbf{R}^d \setminus \Gamma)$. Moreover, $\psi - \psi_n = \chi_n\psi$. But arguing as in the proof of Proposition 4.8 one has

$$H(\psi - \psi_n) = (H\chi_n)\psi + \chi_n(H\psi) + 2\Gamma(\chi_n; \psi)$$

and consequently

$$\|H(\psi - \psi_n)\|_2^2 \leq 3 \left((\|H\psi\|_\infty^2 + \|\psi\|_\infty^2) + 2\|\Gamma(\psi)\|_\infty \right) \|\chi_n\|_{D(\overline{H})}^2.$$

Since $\|\psi - \psi_n\|_2 \leq \|\psi\|_\infty \|\chi_n\|_2$ it follows that $\|(\psi - \psi_n)\|_{D(\overline{H})} \rightarrow 0$ as $n \rightarrow \infty$. Therefore each $\psi \in C_c^\infty(\mathbf{R}^d)$ can be approximated by a sequence $\psi_n \in C_c^\infty(\mathbf{R}^d \setminus \Gamma)$ in the $D(\overline{H})$ -graph norm. But as $C_c^\infty(\mathbf{R}^d)$ is a core of H one concludes that $C_c^\infty(\mathbf{R}^d \setminus \Gamma)$ is also a core. \square

References

- [AH96] ADAMS, D. R., and HEDBERG, L. I., *Function spaces and potential theory*. Grundlehren der mathematische Wissenschaften 314. Springer, Berlin, 1996.
- [Aus07] AUSCHER, P., On necessary and sufficient conditions for L^p -estimates of Riesz transforms associated to elliptic operators on \mathbf{R}^n and related estimates. *Mem. Amer. Math. Soc.* **186**, No. 871 (2007).
- [Aze74] AZENCOTT, R., Behavior of diffusion semi-groups at infinity. *Bull. Soc. Math. France* **102** (1974), 193–240.
- [BH91] BOULEAU, N., and HIRSCH, F., *Dirichlet forms and analysis on Wiener space*, vol. 14 of de Gruyter Studies in Mathematics. Walter de Gruyter & Co., Berlin, 1991.

- [CGT82] CHEEGER, J., GROMOV, M., and TAYLOR, M., Finite propagation speed, kernel estimates for functions of the Laplace operator, and the geometry of complete Riemannian manifolds. *J. Differential Geom.* **17** (1982), 15–53.
- [Dav85] DAVIES, E. B., L^1 properties of second order elliptic operators. *Bull. London Math. Soc.* **17** (1985), 417–436.
- [Dav92] ———, Heat kernel bounds, conservation of probability and the Feller property. *J. Anal. Math.* **58** (1992), 99–119. Festschrift on the occasion of the 70th birthday of Shmuel Agmon.
- [Ebe99] EBERLE, A., *Uniqueness and non-uniqueness of semigroups generated by singular diffusion operators*. Lect. Notes in Math. 1718. Springer-Verlag, Berlin etc., 1999.
- [ERSZ07] ELST, A. F. M. TER, ROBINSON, D. W., SIKORA, A., and ZHU, Y., Second-order operators with degenerate coefficients. *Proc. London Math. Soc.* **95** (2007), 299–328.
- [FOT94] FUKUSHIMA, M., OSHIMA, Y., and TAKEDA, M., *Dirichlet forms and symmetric Markov processes*, vol. 19 of de Gruyter Studies in Mathematics. Walter de Gruyter & Co., Berlin, 1994.
- [Gri99] GRIGOR’YAN, A., Estimates of heat kernels on Riemannian manifolds. In *Spectral theory and geometry (Edinburgh, 1998)*, vol. 273 of London Math. Soc. Lecture Note Ser., 140–225. Cambridge Univ. Press, Cambridge, 1999.
- [HKM93] HEINONEN, J., KILPELÄINEN, T., and MARTIO, O., *Nonlinear potential theory of degenerate elliptic operators*. Oxford Mathematical Monographs. Clarendon Press, Oxford, 1993.
- [Kat80] KATO, T., *Perturbation theory for linear operators*. Second edition, Grundlehren der mathematischen Wissenschaften 132. Springer-Verlag, Berlin etc., 1980.
- [MR92] MA, Z. M., and RÖCKNER, M., *Introduction to the theory of (non symmetric) Dirichlet Forms*. Universitext. Springer-Verlag, Berlin etc., 1992.
- [MZ97] MALY, J., and ZIEMER, P., *Fine regularity of solutions of elliptic partial differential equations*, vol. 51 of Math. Surveys and Monographs. Amer. Math. Soc., Providence, 1997.
- [RS06] ROBINSON, D. W., and SIKORA, A. Analysis of degenerate elliptic operators of Grušin type, 2006. arXiv math.AP/0607584.
- [RS08] ———, Analysis of degenerate elliptic operators of Grušin type. *Math. Z.* **260** (2008), 475–508.
- [RS10a] ———, Degenerate elliptic operators in one-dimension. *J. Evol. Equ.* (2010). arXiv:0909.0567.

- [RS10b] ———, Markov uniqueness of degenerate elliptic operators. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (2010). arXiv:0912.4536.
- [Sta99] STANNAT, W., (Non-symmetric) Dirichlet operators on L^1 : existence, uniqueness and associated Markov processes. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **28** (1999), 99–140.
- [Stu95] STURM, K.-T., Analysis on local Dirichlet spaces. II. Upper Gaussian estimates for the fundamental solutions of parabolic equations. *Osaka J. Math.* **32** (1995), 275–312.
- [Stu98] ———, The geometric aspect of Dirichlet forms. In *New directions in Dirichlet forms*, vol. 8 of AMS/IP Stud. Adv. Math., 233–277. Amer. Math. Soc., Providence, RI, 1998.