

Cliques in graphs with bounded minimum degree

Allan Siu Lun Lo*

September 28, 2010

Abstract

Let $k_r(n, \delta)$ be the minimum number of r -cliques in graphs with n vertices and minimum degree δ . We evaluate $k_r(n, \delta)$ for $\delta \leq 4n/5$ and some other cases. Moreover, we give a construction, which we conjecture to give all extremal graphs (subject to certain conditions on n , δ and r).

1 Introduction

Let $f_r(n, e)$ be the minimum number of r -cliques in graphs of order n and size e . Determining $f_r(n, e)$ has been a long studied problem. The case $r = 3$, that is counting triangles, has been studied by various people. Erdős [3], Lovasz and Simonovits [7] studied the case when $e = \binom{n}{2}/2 + l$ with $0 < l \leq n/2$. Fisher [4] considered the situation when $\binom{n}{2}/2 \leq e \leq 2\binom{n}{2}/3$, but it was not until nearly twenty years later that a dramatic breakthrough of Razborov [9] established the asymptotic value of $f_3(n, e)$ for a general e . The proof of this used the concept of flag algebra developed in [10]. Unfortunately, it seemed difficult to generalise Razborov's proof even for $f_4(n, e)$. Nikiforov [8] later gave a simple and elegant proof of the asymptotic values of both $f_3(n, e)$ and $f_4(n, e)$ for general e . However, the asymptotic value of $f_r(n, e)$ for $r \geq 5$ have not yet been determined, and the best known lower bounds was given Bollobás [2].

In this paper, we are interested in a variant of $f_r(n, e)$, where instead of considering the number of edges we consider the minimum degree. Define $k_r(n, \delta)$ to be the minimum number of r -cliques in

*DPMMS, University of Cambridge, Cambridge CB3 0WB, UK. Email: al-lan.lo@cantab.net This author is supported by EPSRC.

graphs of order n with minimum degree δ . In addition, $k_r^{reg}(n, \delta)$ is defined to be the minimum number of r -cliques in δ -regular graphs of order n . It should be noted that there exist n and δ such that $k_r(n, \delta) = 0$, but $k_r^{reg}(n, \delta) > 0$. For example, if $r = 3$, n odd and $2n/5 < \delta n < 2$, then it is easy to show that $k_3(n, \delta) = 0$. However, a theorem of Andrásfai, Erdős and Sós [1] states that every triangle-free graph of order n with minimal degree greater than $2n/5$ is bipartite. Since no regular graph with an odd number of vertices can be bipartite, $k_3^{reg}(n, \delta) > 0$ for n odd and $2n/5 < \delta < n/2$, whilst $k_3(n, \delta) = 0$. The author [5] evaluated $k_3^{reg}(n, \delta)$ for $n \geq 10^7$ odd and $2n/5 + \sqrt{n}/5 \leq \delta \leq n/2$.

Let $\delta = (1 - \beta)n$ with $0 < \beta \leq 1$ and $p = \lceil \beta^{-1} \rceil - 1$. Throughout this paper, β and βn are assumed to be a rational and an integer respectively. Note that p is defined so that by Turán's Theorem [11] $k_r(n, (1 - \beta)n) > 0$ for all n (such that βn is an integer) if and only if $r \leq p + 1$. Since the case $\beta = 1$ implies the trivial case $\delta = 0$, we may assume that $0 < \beta < 1$. Furthermore, we consider the cases $1/(p + 1) \leq \beta < 1/p$ separately for positive integers p . Hence, the condition $p = 2$ is equivalent to $1/3 \leq \beta < 1/2$, that is, $n/2 < \delta \leq 2n/3$.

Next, we define a family $\mathcal{G}(n, \beta)$ of graphs, which gives an upper bound on $k_r(n, \delta)$, where $\delta = (1 - \beta)n$ and integers $r \geq 3$.

Definition 1.1. *Let n and $(1 - \beta)n$ be positive integers not both odd with $0 < \beta < 1$. Define $\mathcal{G}(n, \beta)$ to be the family of graphs $G = (V, E)$ of order n satisfying the following properties. There is a partition of V into V_0, V_1, \dots, V_{p-1} with $|V_0| = (1 - (p - 1)\beta)n$ and $|V_i| = \beta n$ for $1 \leq i \leq p - 1$, where again $p = \lceil \beta^{-1} \rceil - 1$. For $0 \leq i < j \leq p - 1$, the bipartite graph $G[V_i, V_j]$ induced by the vertex classes V_i and V_j is complete. For $1 \leq i \leq p - 1$, the subgraph $G[V_i]$ induced by V_i is empty and $G[V_0]$ is a $(1 - p\beta)n$ -regular graph such that the number of triangles in $G[V_0]$ is minimal over all $(1 - p\beta)n$ -regular graphs of order $|V_0| = (1 - (p - 1)\beta)n$.*

Note that $\mathcal{G}(n, \beta)$ is only defined if n and $(1 - \beta)n$ are not both odd. Thus, whenever we mention $\mathcal{G}(n, \beta)$, we automatically assume that n or $(1 - \beta)n$ is even. Furthermore, we say (n, β) is *feasible* if $G[V_0]$ is triangle-free for $G \in \mathcal{G}(n, \beta)$. Note that $G[V_0]$ is regular of degree $(1 - p\beta)n \leq (1 - (p - 1)\beta)n/2 = |V_0|/2$. Thus, if $|V_0|$ is even, then $G[V_0]$ is triangle-free. Therefore, for a given β , there exist infinitely many choices of n such that (n, β) is a feasible pair. If (n, β) is not a feasible pair, then $|V_0|$ is odd. Moreover, it is easy to show that $k_3(G[V_0]) = k_3^{reg}(n_0, \delta_0) = o(n^3)$, where $n_0 = |V_0| = (1 - (p - 1)\beta)n$, $\delta_0 = (1 - p\beta)n$ and $k_r(H)$ is the number of r -cliques in a graph H .

By Definition 1.1, every $G \in \mathcal{G}(n, \beta)$ is $(1 - \beta)n$ -regular. In particular, for positive integers $r \geq 3$, the number of r -cliques in G is exactly

$$\begin{aligned} k_r(G) &= g_r(\beta)n^r + \binom{p-1}{r-3}(1-p\beta)^{r-3}n^{r-3}k_3(G[V_0]), \\ &= g_r(\beta)n^r + \binom{p-1}{r-3}(1-p\beta)^{r-3}n^{r-3}k_3^{reg}(n_0, \delta_0), \end{aligned} \quad (1)$$

where $n_0 = (1 - (p-1)\beta)n$, $\delta_0 = (1 - p\beta)n$ and

$$\begin{aligned} g_r(\beta) &= \binom{p-1}{r}\beta^r + \binom{p-1}{r-1}(1-(p-1)\beta)\beta^{r-1} \\ &\quad + \frac{1}{2}\binom{p-1}{r-2}(1-p\beta)(1-(p-1)\beta)\beta^{r-2} \end{aligned}$$

with $\binom{x}{y}$ defined to be 0 if $x < y$ or $y < 0$. Since $k_3^{reg}(n_0, \delta_0) = o(n^3)$, (1) becomes $k_r(G) = (g_r(\beta) + o(1))n^r$. In fact, most of the time, we consider the case when (n, β) is feasible, i.e. $k_3(G[V_0]) = 0$ and $k_r(G) = g_r(\beta)n^r$. We conjecture that if (n, β) is feasible then $\mathcal{G}(n, \beta)$ is the extremal family for $k_r(n, \delta)$ with $\delta = (1 - \beta)n$ and $3 \leq r \leq p+1 = \lceil \beta^{-1} \rceil$.

Conjecture 1.2. *Let n and δ be positive integers. Then*

$$k_r(n, \delta) \geq g_r(\beta)n^r,$$

where $\delta = (1 - \beta)n$ and $r \geq 3$. Moreover, for $3 \leq r \leq p+1 = \lceil \beta^{-1} \rceil$ equality holds if and only if (n, β) is feasible and the extremal graphs are members of $\mathcal{G}(n, \beta)$.

By Turán's Theorem [11], the above conjecture is true when $p = 1$ or $r > p+1$. If $\beta = 1/(p+1)$ and $(p+1)|n$, then $\mathcal{G}(n, 1/(p+1))$ only consists $T_{p+1}(n)$, the $(p+1)$ -partite Turán graph of order n . Bollobás [2] proved that if $(p+1)|n$ and $e = (1 - 1/(p+1))n^2/2$, then $f_r(n, e) = k_r(T_{p+1}(n))$. Moreover, $T_{p+1}(n)$ is the only graph of order n with e edges and $f_r(n, e)$ r -cliques. Hence, it is an easy exercise to show that Conjecture 1.2 is true when $\beta = 1/(p+1)$.

It should be noted that since $\mathcal{G}(n, \beta)$ defines a family of regular graphs, we also conjecture that $k_r^{reg}(n, \delta)$ is achieved by $G \in \mathcal{G}(n, \beta)$, where $\delta = (1 - \beta)n$. However, we do not address the problem $k_r^{reg}(n, \delta)$ here. For the remainder of the paper, all graphs are also assumed to be of order n with minimum degree $\delta = (1 - \beta)n$ unless stated otherwise.

2 Main results

By our previous observation, Conjecture 1.2 is true for the following three cases: $p = 1$, $r > p + 1$ and $\delta = (1 - 1/(p + 1))n$. That leaves the situation when $3 \leq r \leq p + 1$ and $\delta > n/2$. In Section 3, we prove Conjecture 1.2 for $n/2 < \delta \leq 2n/3$, as follows.

Theorem 2.1. *Let n and δ be positive integers with $n/2 < \delta \leq 2n/3$. Then*

$$k_3(n, \delta) \geq g_3(\beta)n^3,$$

where $\delta = (1 - \beta)n$. Moreover, equality holds if and only if (n, β) is feasible and the extremal graphs are members of $\mathcal{G}(n, \beta)$.

The ideas in the proof, which is short, form the framework for our other results. The next simplest case is that of K_{p+2} -free graphs. Notice that, by the definition of p , G must contain K_{p+1} 's but need not contain K_{p+2} . Conjecture 1.2 is proved for K_{p+2} -free graphs by the next theorem.

Theorem 2.2. *Let n and δ be positive integers. Let G be a K_{p+2} -free graph of order n with minimum degree δ , where $\delta = (1 - \beta)n$ and $p = \lceil \beta^{-1} \rceil - 1$. Then,*

$$k_r(G) \geq g_r(\beta)n^r$$

for positive integers r . Moreover, for $3 \leq r \leq p + 1$ equality holds if and only if (n, β) is feasible, and the extremal graphs are members of $\mathcal{G}(n, \beta)$.

Theorem 2.2 is proved in Section 5, after some notations and basic inequalities have been set up in Section 4. It shows that the difficult in proving Conjecture 1.2 is in handling $(p + 2)$ -cliques. We discuss this situation in Section 6 for the case $p = 3$, and by a detailed analysis of 5-cliques in Section 7, proving Conjecture 1.2 for $2n/3 < \delta \leq 3n/4$, as follows.

Theorem 2.3. *Let n and δ be positive integers with $2n/3 < \delta \leq 3n/4$. Then*

$$k_r(n, \delta) \geq g_r(\beta)n^r,$$

for positive integers r and $\delta = (1 - \beta)n$. Moreover, for $3 \leq r \leq 4$ equality holds if and only if (n, β) is feasible and the extremal graphs are members of $\mathcal{G}(n, \beta)$.

This theorem is the hardest in the paper. We have in fact proved Conjecture 1.2 for $3n/4 < \delta \leq 4n/5$ by a similar argument. It is too complicated to be included in this paper, but it can be found

in [6]. For each positive integer $p \geq 5$, it is likely that by following the arguments in the proof of Theorem 2.3 one could construct a proof for Conjecture 1.2 when $(1 - 1/p)n < \delta \leq (1 - 1/(p + 1))n$.

We give two more results in support of Conjecture 1.2 in Section 8 and Section 9. The first is that for every positive integer p , Conjecture 1.2 holds for a positive proportion of values of δ .

Theorem 2.4. *For every positive integer p , there exists a (calculable) constant $\epsilon_p > 0$ so that if n and δ are positive integers such that $(1 - 1/(p + 1) - \epsilon_p)n < \delta \leq (1 - 1/(p + 1))n$, then*

$$k_r(n, \delta) \geq g_r(\beta)n^r,$$

for positive integers r and $\delta = (1 - \beta)n$. Moreover, for $3 \leq r \leq p + 1$ equality holds if and only if (n, β) is feasible and the extremal graphs are members of $\mathcal{G}(n, \beta)$.

Finally, using a different argument, we can show that Conjecture 1.2 holds in the case $r = p + 1$ (the largest value of r for which r -cliques are guaranteed).

Theorem 2.5. *Let n and δ be positive integers. Then*

$$k_{p+1}(n, \delta) \geq g_{p+1}(\beta)n^{p+1},$$

where $\delta = (1 - \beta)n$ and $p = \lceil \beta^{-1} \rceil - 1$. Moreover, equality holds if and only if (n, β) is feasible and the extremal graphs are members of $\mathcal{G}(n, \beta)$.

3 Proof of Theorem 2.1

Here we prove Theorem 2.1, that is Conjecture 1.2 for $n/2 < \delta \leq 2n/3$, so $1/3 \leq \beta < 1/2$ and $p = 2$.

Proof of Theorem 2.1. Let G be a graph of order n with minimum degree δ . Since G has at least $\delta n/2 = (1 - \beta)n^2/2$ edges,

$$(1 - 2\beta)\beta n k_2(G) \geq (1 - 2\beta)(1 - \beta)\beta n^3/2 = g_3(\beta)n^3.$$

Thus, in proving the inequality in Theorem 2.1, it is enough to show that $k_3(G) \geq (1 - 2\beta)\beta n k_2(G)$.

For an edge e , define $d(e)$ to be the number of triangles containing e and write $D(e) = d(e)/n$. Clearly,

$$n \sum_{e \in E(G)} D(e) = \sum d(e) = 3k_3(G).$$

In addition, $D(e) \geq 1 - 2\beta$ for each edge e , because each vertex in G misses at most βn vertices. Since $\beta < 1/2$, $D(e) > 0$ for all $e \in E(G)$ and so every edge is contained in a triangle. Let T be a triangle in G . Similarly, define $d(T)$ to be the number of 4-cliques containing T and write $D(T) = d(T)/n$. We claim that

$$\sum_{e \in E(T)} D(e) \geq 2 - 3\beta + D(T). \quad (2)$$

Let n_i be the number vertices in G with exactly i neighbours in T for $i = 0, 1, 2, 3$. Clearly, $n = n_0 + n_1 + n_2 + n_3$. By counting the number of edges incident with T , we obtain

$$3(1 - \beta)n \leq \sum_{v \in V(T)} d(v) = 3n_3 + 2n_2 + n_1 \leq 2n_3 + n_2 + n. \quad (3)$$

On the other hand, $n_3 = d(T)$ and $n_2 + 3n_3 = \sum_{e \in E(G)} d(e)$. Hence, (2) holds. Notice that if equality holds in (2) then $d(v) = (1 - \beta)n$ for all $v \in T$.

For an edge e , define $D_-(e) = \min\{D(e), \beta\}$. We claim that

$$\sum_{e \in E(T)} D_-(e) \geq 2 - 3\beta \quad (4)$$

for every triangle T . If $D(e) = D_-(e)$ for each edge e in T , then (4) holds by (2). Otherwise, there exists $e_0 \in E(T)$ such that $D(e_0) \neq D_-(e_0)$. This means that $D_-(e_0) = \beta$. Recall that for the other two edges e in T , $D(e) \geq 1 - 2\beta$, so $\sum D_-(e) \geq \beta + 2(1 - 2\beta) = 2 - 3\beta$. Hence, (4) holds for every triangle T .

Next, by summing (4) over all triangles T in G , we obtain

$$n \sum_{e \in E(G)} D_-(e)D(e) = \sum_T \sum_{e \in E(T)} D_-(e) \geq (2 - 3\beta)k_3(G). \quad (5)$$

We are going to bound $\sum D_-(e)D(e)$ above in terms of $\sum D(e)$, which is equal to $3k_3(G)/n$, by the following proposition.

Proposition 3.1. *Let \mathcal{A} be a finite set. Suppose $f, g : \mathcal{A} \rightarrow \mathbb{R}$ with $f(a) \leq M$ and $g(a) \geq m$ for all $a \in \mathcal{A}$. Then*

$$\sum_{a \in \mathcal{A}} f(a)g(a) \leq m \sum_{a \in \mathcal{A}} f(a) + M \sum_{a \in \mathcal{A}} g(a) - mM|\mathcal{A}|,$$

with equality if and only if for each $a \in \mathcal{A}$, $f(a) = M$ or $g(a) = m$.

Proof. Observe that $\sum_{a \in \mathcal{A}} (M - f(a))(g(a) - m) \geq 0$. \square

Recall that $D(e) \geq 1 - 2\beta$ and $D_-(e) \leq \beta$. By Proposition 3.1 taking $\mathcal{A} = E(G)$, $f = D_-$, $g = D$, $M = \beta$ and $m = 1 - 2\beta$, we have

$$\begin{aligned} n \sum_{e \in E(G)} D(e)D_-(e) &\leq (1 - 2\beta)n \sum_{e \in E(G)} D_-(e) + \beta n \sum_{e \in E(G)} D(e) - (1 - 2\beta)\beta nk_2(G) \\ &\leq (1 - \beta)n \sum_{e \in E(G)} D(e) - (1 - 2\beta)\beta nk_2(G) \end{aligned} \quad (6)$$

$$n \sum_{e \in E(G)} D(e)D_-(e) \leq 3(1 - \beta)k_3(G) - (1 - 2\beta)\beta nk_2(G). \quad (7)$$

After substitution of (7) into (5) and rearrangement, we have

$$k_3(G) \geq (1 - 2\beta)\beta k_2(G)n.$$

Thus, we have proved the inequality in Theorem 2.1.

Now suppose equality holds, i.e. $k_3(G) = (1 - 2\beta)\beta k_2(G)n$. This means that equality holds in (6), so (since $\beta < 1/2$) $D(e) = D_-(e)$ for all $e \in E(G)$. Because equality holds in (4), $\sum_{e \in E(T)} D(e) = 2 - 3\beta$ for triangles T . Hence, $D(T) = 0$ for every triangle T by (2), so G is K_4 -free. In addition, by the remark following (2), G is $(1 - \beta)n$ -regular, because every vertex lies in a triangle as $D(e) > 0$ for all edges e . Since equality holds in Proposition 3.1, either $D(e) = 1 - 2\beta$ or $D(e) = \beta$ for each edge e . Recall that equality holds for (2), so every triangle T contains exactly one edge e_1 with $D(e_1) = \beta$ and two edges, e_2 and e_3 , with $D(e_2) = D(e_3) = 1 - \beta$. Pick an edge e with $D(e) = \beta$ and let W be the set of common neighbours of the end vertices of e , so $|W| = \beta n$. Clearly W is an independent set, otherwise G contains a K_4 . For each $w \in W$, $d(w) = (1 - \beta)n$ implies $N(w) = V(G) \setminus W$. Therefore, $G[V(G) \setminus W]$ is $(1 - 2\beta)n$ -regular. If there is a triangle T in $G[V(G) \setminus W]$, then $T \cup w$ forms a K_4 for $w \in W$. This contradicts the assumption that G is K_4 -free, so $G[V(G) \setminus W]$ is triangle-free. Hence, G is a member of $\mathcal{G}(n, \beta)$ and (n, β) is feasible. Therefore, the proof is complete. \square

4 Degree of a clique

Denote the set of t -cliques in $G[U]$ by $\mathcal{K}_t(U)$ and write $k_r(U)$ for $|\mathcal{K}_r(U)|$. If $U = V(G)$, we simply write \mathcal{K}_r and k_r .

Define the *degree* $d(T)$ of a t -clique T to be the number of $(t + 1)$ -cliques containing T . In other words, $d(T) = |\{S \in \mathcal{K}_{t+1} : T \subset S\}|$. If $t = 1$, then $d(v)$ coincides with the ordinary definition of the degree for a vertex v . If $t = 2$, then $d(uv)$ is the number of common neighbours

of the end vertices of the edge uv , that is the codegree of u and v . Clearly, $\sum_{T \in \mathcal{K}_t} d(T) = (t+1)k_{t+1}$ for $t \geq 1$. For convenience, we write $D(T)$ to denote $d(T)/n$.

Recall that $p = \lceil \beta^{-1} \rceil - 1$ and $1/(p+1) \leq \beta < 1/p$. Let $G_0 \in \mathcal{G}(n, \beta)$ with (n, β) feasible. Let T be a t -clique in G_0 . It is natural to see that there are three types of cliques according to $|T \cap V_0|$. However, if we consider $d(T)$, then there are only two types. To be precise

$$D(T) = \begin{cases} 1 - t\beta & \text{if } |V(T) \cap V_0| = 0, 1 \text{ and} \\ (p - t + 1)\beta & \text{if } |V(T) \cap V_0| = 2, \end{cases}$$

for $T \in \mathcal{K}_t(G_0)$ and $2 \leq t \leq p+1$. Next, define the functions D_+ and D_- as follows. For a graph G with minimum degree $\delta = (1 - \beta)n$, define

$$\begin{aligned} D_-(T) &= \min\{D(T), (p - t + 1)\beta\}, \text{ and} \\ D_+(T) &= D(T) - D_-(T) = \max\{0, D(T) - (p - t + 1)\beta\} \end{aligned}$$

for $T \in \mathcal{K}_t$ and $1 \leq t \leq p+1$. We say that a clique T is *heavy* if $D_+(T) > 0$. The graph G is said to be *heavy-free* if and only if G does not contain any heavy cliques. Now, we study some basic properties of $D(T)$, $D_-(T)$ and $D_+(T)$.

Lemma 4.1. *Let $0 < \beta < 1$ and $p = \lceil \beta^{-1} \rceil - 1$. Suppose G is a graph of order n with minimum degree $(1 - \beta)n$. Suppose $S \in \mathcal{K}_s$ and $T \in \mathcal{K}_t(S)$ for $1 \leq t < s$. Then*

- (i) $D(S) \geq 1 - s\beta$,
- (ii) $D(S) \geq D(T) - (s - t)\beta$,
- (iii) for $s \leq p+1$, $D_+(T) \leq D_+(S) \leq D_+(T) + (s - t)\beta$,
- (iv) if T is heavy and $s \leq p+1$ then S is heavy, and
- (v) if T is not heavy and $s \leq p+1$, then $D_+(S) \leq (s - t)\beta$. In particular, if $t = s - 1 \leq p$, then $D_+(S) \leq \beta$.

Moreover, G is K_{p+2} -free if and only if G is heavy-free.

Proof. For each $v \in S$, there are at most βn vertices not joined to v . Hence, $D(S) \geq 1 - s\beta$, so (i) is true. Similarly, consider the vertices in $S \setminus T$, so (ii) is also true. If $s \leq p+1$ and $D_+(T) > 0$, then we have

$$\begin{aligned} D_+(S) + (p - s + 1)\beta &\geq D(S) \\ &\geq D(T) - (s - t)\beta \\ &= D_+(T) + (p - t + 1)\beta - (s - t)\beta, \end{aligned}$$

so the left inequality of (iii) is true. Since $D(S) \leq D(T)$, the right inequality of (iii) is also true by the definition of $D_+(S)$ and $D_+(T)$. Hence, (iv) and (v) are true by the left and right inequality in (iii) respectively. Notice that $D(U) = D_+(U)$ for $U \in \mathcal{K}_{p+1}$. Hence, by (iv), G is K_{p+2} -free if and only if G is heavy-free. \square

Now we prove the generalised version of (2), that is, the sum of degrees of t -subcliques in a s -clique.

Lemma 4.2. *Let $0 < \beta < 1$. Let s and t be integers with $2 \leq t < s$. Suppose G is a graph of order n with minimum degree $(1 - \beta)n$. Then*

$$\sum_{T \in \mathcal{K}_t(S)} D(T) \geq (1 - \beta)s \binom{s-2}{t-1} - (t-1) \binom{s-1}{t} + \binom{s-2}{t-2} D(S)$$

for $S \in \mathcal{K}_s$. Moreover, if equality holds, then $d(v) = (1 - \beta)n$ for all $v \in S$.

Proof. Let n_i be the number of vertices with exactly i neighbours in S . The following three equations :

$$\sum_i n_i = n, \tag{8}$$

$$\sum_i i n_i = \sum_{v \in V(S)} d(v) \geq s(1 - \beta)n, \tag{9}$$

$$\sum_i \binom{i}{t} n_i = \sum_{T \in \mathcal{K}_t(S)} D(T)n, \tag{10}$$

follow from a count of the number of vertices, edges and $(t+1)$ -cliques respectively. Next, by considering $(t-1) \binom{s-1}{t} (8) - \binom{s-2}{t-1} (9) + (10)$, we have

$$\sum_{T \in \mathcal{K}_t(S)} D(T)n \geq \left((1 - \beta)s \binom{s-2}{t-1} - (t-1) \binom{s-1}{t} \right) n + \sum_{0 \leq i \leq s} x_i n_i,$$

where $x_i = \binom{i}{t} + (t-1) \binom{s-1}{t} - i \binom{s-2}{t-1}$. Notice that $x_i = x_{i+1} + \binom{s-2}{t-1} - \binom{i}{t-1} \geq x_{i+1}$ for $0 \leq i \leq s-2$. For $i = s-1$, we have

$$\begin{aligned} x_{s-1} &= \binom{s-1}{t} + (t-1) \binom{s-1}{t} - (s-1) \binom{s-2}{t-1} \\ &= t \binom{s-1}{t} - (s-1) \binom{s-2}{t-1} = 0. \end{aligned}$$

For $i = s$, $n_s = D(S)n$ and

$$\begin{aligned}
x_s &= \binom{s}{t} + (t-1)\binom{s-1}{t} - s\binom{s-2}{t-1} \\
&= t\binom{s-1}{t} + \binom{s-1}{t-1} - s\binom{s-2}{t-1} \\
&= (s-t+1)\binom{s-1}{t-1} - s\binom{s-2}{t-1} \\
&= (s-t+1)\binom{s-2}{t-2} - (t-1)\binom{s-2}{t-1} \\
&= \binom{s-2}{t-2}.
\end{aligned}$$

In particular, if equality holds in the lemma, then equality holds in (9). This means that $d(v) = (1 - \beta)n$ for all $v \in S$. \square

Most of the time, we are only interested in the case when $s = t + 1$. Hence, we state the following corollary.

Corollary 4.3. *Let $0 < \beta < 1$. Suppose G is a graph of order n with minimum degree $(1 - \beta)n$. Then*

$$\sum_{T \in \mathcal{K}_t(S)} D(T) \geq 2 - (t+1)\beta + (t-1)D(S)$$

for $S \in \mathcal{K}_{t+1}$ and integer $t \geq 2$. Moreover, if equality holds, then $d(v) = (1 - \beta)n$ for all $v \in S$. \square

In the next lemma, we show that the functions D in Lemma 4.2 can be replaced with D_- .

Lemma 4.4. *Let $0 < \beta < 1$ and $p = \lceil \beta^{-1} \rceil - 1$. Let s and t be integers with $2 \leq t < s \leq p + 1$. Suppose G is a graph of order n with minimum degree $(1 - \beta)n$. Then, for $S \in \mathcal{K}_s$*

$$\sum_{T \in \mathcal{K}_t(S)} D_-(T) \geq (1 - \beta)s \binom{s-2}{t-1} - (t-1) \binom{s-1}{t} + \binom{s-2}{t-2} D_-(S).$$

Proof. Since $D_+(S) \geq D_+(T)$ for every $T \in \mathcal{K}_t(S)$ by Lemma 4.1 (iii), there is nothing to prove by Lemma 4.2 if there are at most $\binom{s-2}{t-2}$ heavy t -cliques in S . Now suppose there are more than $\binom{s-2}{t-2}$ heavy t -cliques in S . In particular, S contains a heavy t -clique, so S is itself heavy with $D_-(S) = (p + 1 - s)\beta$ by Lemma 4.1 (iv). Thus, the right hand side of the inequality is $\binom{s}{t}(1 - t\beta) + \binom{s-2}{t-2}((p + 1)\beta - 1)$. By Lemma 4.1 (i) we have that $D_-(T) \geq (1 - t\beta)$ for $T \in \mathcal{K}_t(S)$.

Furthermore, by Lemma 4.1 (iv) $D_-(T) = (p - t + 1)\beta$ if T is heavy, so summing $D_-(T)$ over $T \in \mathcal{K}_t(S)$ gives

$$\begin{aligned} \sum_{T \in \mathcal{K}_t(S)} D_-(T) &\geq k_t^+(S)(p - t + 1)\beta + \left(\binom{s}{t} - k_t^+(S) \right) (1 - t\beta) \\ &= \binom{s}{t} (1 - t\beta) + k_t^+(S)((p + 1)\beta - 1). \end{aligned}$$

This completes the proof of the lemma. \square

Define the function $\tilde{D} : \mathcal{K}_{t+1} \rightarrow \mathbb{R}$ such that

$$\tilde{D}(S) = \sum_{T \in \mathcal{K}_t(S)} D_-(T) - \left(2 - (t + 1)\beta + (t - 1)D_-(S) \right)$$

for $S \in \mathcal{K}_{t+1}$ and $2 \leq t \leq p$. Hence, for $s = t + 1$, Lemma 4.4 gives the following corollary.

Corollary 4.5. *Let $0 < \beta < 1$ and $p = \lceil \beta^{-1} \rceil - 1$. Let t be integer with $2 \leq t \leq p$. Suppose G is a graph of order n with minimum degree $(1 - \beta)n$. Then $\tilde{D}(S) \geq 0$ for $S \in \mathcal{K}_{t+1}$. \square*

Next, we bound $\sum_{S \in \mathcal{K}_{t+1}} \tilde{D}(S)$ from above using Proposition 3.1.

Lemma 4.6. *Let $0 < \beta < 1$ and $p = \lceil \beta^{-1} \rceil - 1$. Let t be an integer with $2 \leq t \leq p$. Suppose G is a graph of order n with minimum degree $(1 - \beta)n$. Then*

$$\begin{aligned} \sum_{S \in \mathcal{K}_{t+1}} \tilde{D}(S) &\leq (t - 1 + (p - 2t + 2)(t + 1)\beta)k_{t+1} + (t - 1) \sum_{S \in \mathcal{K}_{t+1}} D_+(S) \\ &\quad - (1 - t\beta)(p - t + 1)\beta nk_t - (t - 1)(t + 2)\frac{k_{t+2}}{n} - (1 - t\beta)n \sum_{T \in \mathcal{K}_t} D_+(T). \end{aligned}$$

Moreover, equality holds if and only if for each $T \in \mathcal{K}_t$, either $D_-(T) = 1 - t\beta$ or $D_-(T) = (p - t + 1)\beta$.

Proof. Notice that the sum $\tilde{D}(S)$ over $S \in \mathcal{K}_{t+1}$ is equal to

$$\sum_{S \in \mathcal{K}_{t+1}} \sum_{T \in \mathcal{K}_t(S)} D_-(T) - (2 - (t + 1)\beta)k_{t+1} - (t - 1) \sum_{S \in \mathcal{K}_{t+1}} D_-(S). \quad (11)$$

Consider each each term separately. Since $D(S) = D_-(S) + D_+(S)$,

$$\sum_{S \in \mathcal{K}_{t+1}} D_-(S) = \sum_{S \in \mathcal{K}_{t+1}} D(S) - \sum_{S \in \mathcal{K}_{t+1}} D_+(S) = \frac{(t + 2)k_{t+2}}{n} - \sum_{S \in \mathcal{K}_{t+1}} D_+(S).$$

By interchanging the order of summations, we have

$$\sum_{S \in \mathcal{K}_{t+1}} \sum_{T \in \mathcal{K}_t(S)} D_-(T) = n \sum_{T \in \mathcal{K}_t} D_-(T) D(T),$$

and by Proposition 3.1 taking $\mathcal{A} = \mathcal{K}_t$, $f = D_-$, $g = D$, $M = (p - t + 1)\beta$ and $m = 1 - t\beta$

$$\begin{aligned} &\leq (1 - t\beta)n \sum_{T \in \mathcal{K}_t} D_-(T) + (p - t + 1)\beta n \sum_{T \in \mathcal{K}_t} D(T) - (1 - t\beta)(p - t + 1)\beta n k_t \\ &= (1 + (p - 2t + 1)\beta)n \sum_{T \in \mathcal{K}_t} D(T) - (1 - t\beta)n \sum_{T \in \mathcal{K}_t} D_+(T) - (1 - t\beta)(p - t + 1)\beta n k_t \\ &= (1 + (p - 2t + 1)\beta)(t + 1)k_{t+1} - (1 - t\beta)n \sum_{T \in \mathcal{K}_t} D_+(T) - (1 - t\beta)(p - t + 1)\beta n k_t. \end{aligned}$$

Hence, substituting these identities back into (11), we obtain the desired inequality in the lemma.

By Proposition 3.1, equality holds if and only if for each $T \in \mathcal{K}_t$, either $D(T) = 1 - t\beta$ or $D_-(T) = (p - t + 1)\beta$. \square

To keep our calculations simple, we are going to establish a few relationships between $g_t(\beta)$ and $g_{t+1}(\beta)$ in the next lemma.

Lemma 4.7. *Let $0 < \beta < 1$ and $p = \lceil \beta^{-1} \rceil - 1$. Let t be an integer with $2 \leq t \leq p$. Then*

$$\begin{aligned} (t + 1)g_{t+1}(\beta) &= (1 - t\beta)g_t(\beta) \\ &\quad + \frac{1}{2} \binom{p-1}{t-2} ((p+1)\beta - 1)(1 - (p-1)\beta)(1 - p\beta)\beta^{t-2}, \end{aligned} \tag{12}$$

$$g_{t+1}(\beta) = \frac{(1 - t\beta)(p - t + 1)\beta g_t(\beta) + (t - 1)(t + 2)g_{t+2}(\beta)}{t - 1 + (t + 1)(p - 2t + 2)\beta}. \tag{13}$$

Moreover

$$\frac{g_p(\beta)}{g_{p+1}(\beta)} = \frac{1}{\beta} \left(1 + \frac{\beta g_{p-1}(\beta')}{(1 - \beta)g_p(\beta')} \right), \tag{14}$$

where $\beta' = \beta/(1 - \beta)$.

Proof. We fix β (and p) and write g_t to denote $g_t(\beta)$. Pick n such that (n, β) is feasible and let $G \in \mathcal{G}(n, \beta)$ with partition classes V_0, V_1, \dots, V_{p-1} as described in Definition 1.1. Thus, for $T \in \mathcal{K}_t$, $D(T) = 1 - t\beta$ or $D(T) = (p - t + 1)\beta$. Since $D(T) = (p - t + 1)\beta$ if and only if $|V(T) \cap V_0| = 2$, there are exactly

$$\frac{1}{2} \binom{p-1}{t-2} (1 - (p-1)\beta)(1 - p\beta)\beta^{t-2} n^t$$

t -cliques T with $D(T) = (p - t + 1)\beta$. Also, we have

$$(t + 1)g_{t+1}n^{t+1} = (t + 1)k_{t+1} = n \sum_{T \in \mathcal{K}_t} D(T).$$

Hence, (12) is true, by expanding the right hand side of the above equation. For $2 \leq s < p$, let f_s and f_{s+1} be (12) with $t = s$ and $t = s + 1$ respectively. Then (13) follows by considering $(p - s + 1)f_s - (s - 1)\beta f_{s+1}$.

Now let $G' = G \setminus V_{p-1}$. Notice that G' is $(1 - 2\beta)n$ -regular with $(1 - \beta)n$ vertices. We observe that G' is a member of $\mathcal{G}(n', \beta')$, where $n' = (1 - \beta)n$ and $\beta' = \beta/(1 - \beta)$. Observe that $\lceil \beta'^{-1} \rceil - 1 = p - 1$, so $1/p \leq \beta' < 1/(p - 1)$. Recall that $k_t(G) = g_t(\beta)n^t$ for all $2 \leq t \leq p$, so $k_{p+1}(G)g_p(\beta) = k_p(G)g_{p+1}(\beta)n$. Similarly, $k_p(G')g_{p-1}(\beta') = k_{p-1}(G')g_p(\beta')n$. By considering $\mathcal{K}_p(G)$ and $\mathcal{K}_{p+1}(G)$, we obtain the following two equations :

$$k_{p+1}(G) = \beta n k_p(G'), \quad (15)$$

$$\begin{aligned} k_p(G) &= \beta n k_{p-1}(G') + k_p(G') = \beta n \frac{g_{p-1}(\beta')k_p(G')}{n'g_p(\beta')} + k_p(G') \\ &= \left(1 + \frac{\beta g_{p-1}(\beta')}{(1 - \beta)g_p(\beta')}\right) k_p(G'). \end{aligned} \quad (16)$$

By substituting (15) and (16) into $k_p(G)n/k_{p+1}(G) = g_p(\beta)/g_{p+1}(\beta)$, we obtain (14). The proof is complete. \square

5 K_{p+2} -free graphs

In this section, all graphs are assumed to be K_{p+2} -free. Lemma 4.1 implies that these graphs are also heavy-free. This means that $D_+(T) = 0$ and $D(T) \leq (p - t + 1)\beta$ for all $T \in \mathcal{K}_t$ and $t \leq p + 1$. We prove the theorem below, which easily implies Theorem 2.2 as $g_2(\beta)n^2 = (1 - \beta)n^2/2 \leq k_2(G)$.

Theorem 5.1. *Let $0 < \beta < 1$ and $p = \lceil \beta^{-1} \rceil - 1$. Suppose G is a K_{p+2} -free graph of order n with minimum degree $(1 - \beta)n$. Then*

$$\frac{k_s(G)}{g_s(\beta)n^s} \geq \frac{k_t(G)}{g_t(\beta)n^t} \quad (17)$$

holds for $2 \leq t < s \leq p + 1$. Moreover, the following three statements are equivalent:

- (i) *Equality holds for some $2 \leq t < s \leq p + 1$.*
- (ii) *Equality holds for all $2 \leq t < s \leq p + 1$.*

(iii) The pair (n, β) is feasible and G is a member of $\mathcal{G}(n, \beta)$.

Proof. Fix β and write g_t to denote $g_t(\beta)$. Recall that $D_+(T) = 0$ for cliques T . By Corollary 4.5 and Lemma 4.6, we have

$$k_{t+1} \geq \frac{(1 - t\beta)(p - t + 1)\beta nk_t + (t - 1)(t + 2)k_{t+2}/n}{t - 1 + (p - 2t + 2)(t + 1)\beta} \quad (18)$$

First, we are going to prove (17). It is sufficient to prove the case when $s = t + 1$. We proceed by induction on t from above. For $t = p$, $k_{p+2} = 0$ and so (18) becomes

$$(p - 1 - (p - 2)(p + 1)\beta)k_{p+1} \geq (1 - p\beta)\beta nk_p.$$

Since $g_{p+2} = 0$, we have $k_{p+1}/g_{p+1}n^{p+1} \geq k_p/g_p n^p$ by (13). Hence, (17) is true for $t = p$. For $t < p$, (18) becomes

$$\begin{aligned} & (t - 1 + (t + 1)(p - 2t + 2)\beta)k_{t+1} \\ & \geq (1 - t\beta)(p + 1 - t)\beta nk_t + (t - 1)(t + 2)k_{t+2}/n \end{aligned}$$

by the induction hypothesis

$$\geq (1 - t\beta)(p + 1 - t)\beta nk_t + (t - 1)(t + 2)g_{t+2}k_{t+1}/g_{t+1}. \quad (19)$$

Thus, (17) follows from (13).

It is clear that (iii) implies both (i) and (ii) by Definition 1.1 and the feasibility of (n, β) . Suppose (i) holds, so equality holds in (17) for $t = t_0$ and $s = s_0$ with $t_0 < s_0$. We claim that equality must also hold for $t = p$ and $s = p + 1$. Suppose the claim is false and equality holds for $t = t_0$ and $s = s_0$, where s_0 is maximal. Since equality holds for $t = t_0$, by (17), equality holds for $t = t_0, \dots, s_0 - 1$ with $s = s_0$. We may assume that $t = s_0 - 1$ and $s_0 \neq p + 1$ and $k_{s_0+1}/g_{s_0+1}n > k_{s_0}/g_{s_0}$. However, this would imply a strictly inequality in (19) contradicting the fact that equality holds for $s = s_0$ and $t = s_0 - 1$. Thus, the proof of the claim is complete, that is, if (i) holds then equality holds in (17) for $t = p$ and $s = p + 1$.

Therefore, in order to prove that (i) implies (iii), it is sufficient to show that if $k_{p+1}/g_{p+1}n^{p+1} = k_p/g_p n^p$, then (n, β) is feasible and G is a member of $\mathcal{G}(n, \beta)$. We proceed by induction on p . It is true for $p = 2$ by Theorem 3, so we may assume $p \geq 3$. Since equality holds in (17), we have equality in (18), Corollary 4.5 and Lemma 4.6. Since D_+ is a zero function, equality in Corollary 4.5 implies equality in Corollary 4.3 and so G is $(1 - \beta)n$ -regular as every vertex is a $(p + 1)$ -clique. In addition, for each $T \in \mathcal{K}_p$, either $D(T) = 1 - p\beta$ or $D(T) = \beta$ by equality in Lemma 4.6. Moreover, Corollary 4.3 implies

that $\sum_{T \in \mathcal{K}_p(S)} D(T) \geq 2 - (p+1)\beta$ for $S \in \mathcal{K}_{p+1}$. Thus, there exists $T \in \mathcal{K}_p(S)$ with $D(T) = \beta$. Pick $T \in \mathcal{K}_p$ with $D(T) = \beta$ and let $W = \bigcap \{N(v) : v \in V(S)\}$, so $|W| = \beta$. Since G is K_{p+2} -free, W is a set of independent vertices. For each $w \in W$, $d(w) = (1-\beta)n$, so $N(w) = V(G) \setminus W$. Thus, the graph $G' = G[V(G) \setminus W]$ is $(1-\beta)n'$ -regular, where $n' = (1-\beta)n$, $\beta'n' = (1-2\beta)n$ and $\beta' = \beta/(1-\beta)$. Note that $\lceil \beta'^{-1} \rceil - 1 = p-1$. Since G is K_{p+2} -free, G' is K_{p+1} -free. Also, $k_{p+1}(G) = \beta n k_p(G')$ and

$$\begin{aligned} k_p(G) &= \beta n k_{p-1}(G') + k_p(G') \stackrel{\text{by (17)}}{\leq} \beta \frac{g_{p-1}(\beta') k_p(G')}{g_p(\beta')} + k_p(G') \\ &= \left(1 + \beta \frac{g_{p-1}(\beta')}{(1-\beta)g_p(\beta')}\right) k_p(G') \stackrel{\text{by (14)}}{=} \frac{g_p(\beta)\beta}{g_{p+1}(\beta)} k_p(G'). \end{aligned} \quad (20)$$

Hence,

$$g_p(\beta)\beta n k_p(G') = g_p(\beta)k_{p+1}(G) \stackrel{\text{by (17)}}{=} g_{p+1}(\beta)n k_p(G) \stackrel{\text{by (20)}}{\leq} g_p(\beta)\beta n k_p(G').$$

Therefore, we have $k_p(G')/g_p(\beta')n^p = k_{p-1}(G')/g_{p-1}(\beta')n^{p-1}$. By the induction hypothesis, $G' \in \mathcal{G}(n', \beta')$, which implies $G \in \mathcal{G}(n, \beta)$. This completes the proof of the theorem. \square

6 $k_r(n, \delta)$ for $2n/3 < \delta \leq 3n/4$

By Theorem 2.2, in order to prove Conjecture 1.2 it remains to handle the heavy cliques. However, even though both Corollary 4.5 and Lemma 4.6 are sharp by considering $G \in \mathcal{G}(n, \beta)$, they are not sufficient to prove Conjecture 1.2 even for the case when $2n/3 < \delta \leq 3n/4$ by the observation below. Let $2n/3 < \delta \leq 3n/4$, $1/4 \leq \beta < 1/3$ and $p = 3$. By Corollary 4.5 and Lemma 4.6, we have

$$(1+3\beta)k_3 + \sum_{T \in \mathcal{K}_3} D_+(T) \geq 2(1-2\beta)\beta n k_2 + \frac{4}{n}k_4 + (1-2\beta)n \sum_{e \in \mathcal{K}_2} D_+(e), \quad (21)$$

$$(2-4\beta)k_4 + 2 \sum_{S \in \mathcal{K}_4} D_+(S) \geq (1-3\beta)\beta n k_3 + \frac{10}{n}k_5 + (1-3\beta)n \sum_{T \in \mathcal{K}_3} D_+(T), \quad (22)$$

for $t = 2$ and $t = 3$ respectively. Since D_- is a zero function on 4-cliques,

$$\sum_{S \in \mathcal{K}_4} D_+(S) = \sum_{S \in \mathcal{K}_4} D(S) = 5k_5/n.$$

Hence, the terms with k_5 and $\sum D_+(S)$ cancel in (22). Also, $(1-2\beta) > 0$, so we may ignore the term with $\sum D_+(e)$ in (21). Recall that $g_2(\beta) = (1-\beta)/2$ and $g_3(\beta) = (1-2\beta)^2\beta$. After substitution of (22) into (21) replacing the k_4 term and rearrangement, we get

$$k_3(G) \geq g_3(\beta)n^3 - \frac{4\beta-1}{1-\beta} \sum_{T \in \mathcal{K}_3} D_+(T).$$

However, $(4\beta-1) \geq 0$ only if $\beta = 1/4$. Hence, we are going to strengthen both (22) and (21). Recall that (21) is a consequence of Corollary 4.5 and Lemma 4.6 for $t = 2$. Therefore, the following lemma, which is a strengthening of Corollary 4.5 for $t = 2$, would lead to a strengthening of (21).

Lemma 6.1. *Let $1/4 \leq \beta < 1/3$. Suppose G is a graph of order n with minimum degree $(1-\beta)n$. Then, for $T \in \mathcal{K}_3$*

$$\tilde{D}(T) \geq \left(1 - \frac{2}{29-75\beta}\right) \frac{4\beta-1}{1-2\beta} D_+(T) - (1-2\beta) \sum_{e \in \mathcal{K}_2(T)} \frac{D_+(e)}{D_+(e)+\beta}. \quad (23)$$

Moreover, if equality holds then T is not heavy and $d(v) = (1-\beta)n$ for all $v \in T$.

Proof. Let c be $(1-2/(29-75\beta))(4\beta-1)/(1-2\beta)$. Corollary 4.5 gives $\tilde{D}(T) \geq 0$, so we may assume that T is heavy. In addition, Corollary 4.3 implies that

$$\tilde{D}(T) + \sum_{e \in \mathcal{K}_2(T)} D_+(e) \geq D_+(T). \quad (24)$$

Since $c < 1$, we may further assume that T contains at least one heavy edge or else (23) holds as (24) becomes $\tilde{D}(T) \geq D_+(T) > cD_+(T)$. Let $e_0 \in \mathcal{K}_2(T)$ with $D_+(e_0)$ maximal. By substituting (24) into (23), it is sufficient to show that the function

$$f = \left(1 - \frac{1-2\beta}{D_+(e_0)+2\beta}\right) \tilde{D}(T) - \left(c - \frac{1-2\beta}{D_+(e_0)+2\beta}\right) D_+(T)$$

is non-negative.

First consider the case when $D_+(T) \leq 1-3\beta$. Lemma 4.1 (iii) implies $D_+(e_0) \leq D_+(T) \leq 1-3\beta$. Hence,

$$\frac{1-2\beta}{D_+(e_0)+2\beta} - c \geq \frac{1-2\beta}{1-\beta} - c > 0.$$

Also, $1 - 2\beta \leq 2\beta < D_+(e_0) + 2\beta$. Therefore, $f > 0$ by considering the coefficients of $\tilde{D}(T)$ and $D(T)$. Hence, we may assume $D_+(T) > 1 - 3\beta$. Since T is heavy, $D_-(T) = \beta$. Therefore, by the definition of \tilde{D} , we have

$$\tilde{D}(T) = \sum_{e \in \mathcal{K}_2(T)} D_-(e) - 2(1 - \beta). \quad (25)$$

We split into different cases separately depending on the number of heavy edges in T .

Suppose all edges are heavy. Thus, $\tilde{D}(T) = 2(4\beta - 1)$ by (25), because $D_-(e) = 2\beta$ for all edges e in T . Clearly $D_+(T) = D(T) - \beta \leq 1 - \beta$. Hence, (23) is true as

$$\tilde{D}(T) = 2(4\beta - 1) \geq (4\beta - 1)(1 - \beta)/(1 - 2\beta) \geq cD_+(T).$$

Thus, there exists an edge in T that is not heavy and $D_+(T) \leq \beta$ by Lemma 4.1 (v).

Suppose T contains one or two heavy edges. We are going to show that in both cases

$$\tilde{D}(T) \geq 2(D_+(T) - (1 - 3\beta)).$$

First assume that there is exactly one heavy edge in T . Let e_1 and e_2 be the two non-heavy edges in T . Note that $D_-(e_i) = D(e_i) \geq D(T) = D_+(T) + \beta$ for $i = 1, 2$. Thus, (25) and Lemma 4.1 imply that $\tilde{D}(T) \geq 2(D_+(T) - (1 - 3\beta))$. Assume that T contains two heavy edges. Let e_1 be the non-heavy edge in T . Similarly, we have $D_-(e_1) \geq D_+(T) + \beta$. Recall that $D_+(T) \leq \beta$, so (25) and Lemma 4.1 imply

$$\begin{aligned} \tilde{D}(T) &\geq (4\beta + D_+(T) - (1 - 3\beta)) \\ &= 4\beta - 1 + D_+(T) - (1 - 3\beta) \geq 2(D_+(T) - (1 - 3\beta)). \end{aligned}$$

Since $\tilde{D}(T) \geq 2(D_+(T) - (1 - 3\beta))$, in proving (23), it is enough to show that

$$\begin{aligned} D(e_0)f &= (D_+(e_0) + 2\beta)f \\ &\geq 2(D_+(e_0) + 4\beta - 1)(D_+(T) - (1 - 3\beta)) \\ &\quad - ((D_+(e_0) + 2\beta)c - (1 - 2\beta))D_+(T) \end{aligned} \quad (26)$$

is non-negative for $0 < D_+(e_0) \leq D_+(T)$ and $1 - 3\beta \leq D_+(T) \leq \beta$. Notice that for a fixed $D_+(T)$ it is enough to check the boundary points of $D_+(e_0)$. For $D_+(e_0) = 0$, we have

$$\begin{aligned} D(e_0)f &\geq (2(3 - c)\beta - 1)D_+(T) - 2(4\beta - 1)(1 - 3\beta) \\ &\geq (4\beta - 1)(D_+(T) - (1 - 3\beta)) > 0. \end{aligned}$$

For $D_+(e_0) = D_+(T)$, the right hand side of (26) becomes a quadratic function in $D_+(T)$. Moreover, both coefficients of $D_+(T)^2$ and $D_+(T)$ are positive. Thus, it enough to check for $D_+(T) = 1 - 3\beta$. For $D_+(T) = D_+(e_0) = 1 - 3\beta$, (26) becomes

$$D(e_0)f \geq (1 - c - (2 - c)\beta)(1 - 3\beta) > 0.$$

Hence, we have proved the inequality in Lemma 6.1.

It is easy to check that if equality holds in (23) then $D_+(T) = 0$. Thus, for all edges e in T , $D_+(e) = 0$ by Lemma 4.1. Furthermore, equality holds in (24), so equality holds in Corollary 4.3 as $D_+(T) = 0 = D_+(e)$. Hence, $d(v) = (1 - \beta)n$ for $v \in S$. This completes the proof of the lemma. \square

Together with Lemma 4.6 with $t = 2$, we obtain the strengthening of (21).

Corollary 6.2. *Let $1/4 \leq \beta < 1/3$. Suppose G is a graph of order n with minimum degree $(1 - \beta)n$. Then*

$$(1 + 3\beta)k_3 + \frac{2}{1 - 2\beta} \left(1 - 3\beta + \frac{4\beta - 1}{29 - 75\beta} \right) \sum_{T \in \mathcal{K}_3} D_+(T) \geq 2(1 - 2\beta)\beta nk_2 + 4\frac{k_4}{n}$$

holds. Moreover, if equality holds, then G is $(1 - \beta)n$ -regular and for each edge e , either we have $D(e) = 1 - 2\beta$ or $D(e) = 2\beta$. \square

Note that by mimicking the proof of Lemma 6.1, we could obtain a strengthening of Corollary 4.5 for $t = 3$. It would lead to a strengthening of (22). However, it is still not sufficient to prove the Conjecture 1.2 when β is close to $1/3$. Instead, we prove the following statement. The proof requires a detailed analysis of \mathcal{K}_5 , so it is postponed to Section 7.

Lemma 6.3. *Let $1/4 \leq \beta < 1/3$. Suppose G is a graph order n with minimum degree $(1 - \beta)n$. Then*

$$(2 - 4\beta)k_4 \geq (1 - 3\beta)\beta nk_3 + \left(1 - 3\beta + \frac{4\beta - 1}{29 - 75\beta} \right) n \sum_{T \in \mathcal{K}_3} D_+(T). \quad (27)$$

Moreover, equality holds only if (n, β) is feasible, and $G \in \mathcal{G}(n, \beta)$.

By using the two strengthened versions of (21) and (22), that is, Corollary 6.2 and Lemma 6.3, we prove the theorem below, which implies Theorem 2.3.

Theorem 6.4. *Let $1/4 \leq \beta < 1/3$. Let s and t be integers with $2 \leq t < s \leq 4$. Suppose G is a graph of order n with minimum degree $(1 - \beta)n$. Then*

$$\frac{k_s(G)}{g_s(\beta)n^s} \geq \frac{k_t(G)}{g_t(\beta)n^t}.$$

Moreover, the following three statements are equivalent:

- (i) Equality holds for some $2 \leq t < s \leq 4$.
- (ii) Equality holds for all $2 \leq t < s \leq 4$.
- (iii) The pair (n, β) is feasible, and G is a member of $\mathcal{G}(n, \beta)$.

Proof. Recall that $p = 3$ as $1/4 \leq \beta < 1/3$, so

$$g_2(\beta) = (1 - \beta)/2, \quad g_3(\beta) = (1 - 2\beta)^2\beta \text{ and } g_4 = (1 - 2\beta)(1 - 3\beta)\beta^2/2.$$

Note that in proving the inequality, it is sufficient to prove the case when $s = t + 1$. Lemma 6.3 states that $(2 - 4\beta)k_4 \geq (1 - 3\beta)\beta nk_3$. This implies $k_4/g_4(\beta)n^4 \geq k_3/g_3(\beta)n^3$ by (13) with $t = 3$. Hence, the theorem is true for $t = 3$. For $t = 2$, by substituting Corollary 6.2 into Lemma 6.3, we obtain

$$\begin{aligned} (1 + 3\beta)k_3 + \frac{2}{1 - 2\beta} \left(1 - 3\beta + \frac{4\beta - 1}{29 - 75\beta} \right) \sum_{T \in \mathcal{K}_3} D_+(T) &\geq 2(1 - 2\beta)\beta nk_2 \\ &+ \frac{4}{(2 - 4\beta)n} \left((1 - 3\beta)\beta nk_3 + \left(1 - 3\beta + \frac{4\beta - 1}{29 - 75\beta} \right) n \sum D_+(T) \right). \end{aligned}$$

Observe that the $\sum D_+(T)$ terms on both sides cancel. Hence, after rearrangement, we have $(1 - \beta)k_3 \geq 2(1 - 2\beta)^2\beta nk_2$. Thus, $k_3/g_3(\beta)n^4 \geq k_2/g_2(\beta)n^3$ as required.

This is clear that (iii) implies (i) and (ii) by the construction of $\mathcal{G}(n, \beta)$ and the feasibility of (n, β) . Suppose (i) holds, so equality holds for some $2 \leq t < s \leq 4$. It is easy to deduce that equality also holds for $s = 4$ and $t = 3$. By Lemma 6.3, (n, β) is feasible, and $G \in \mathcal{G}(n, \beta)$. \square

7 Proof of Lemma 6.3

In this section, T , S and U always denote a 3-clique, 4-clique and 5-clique respectively. Before presenting the proof, we recall some basic facts about T , S and U . Observe that $D_-(S) = 0$ for $S \in \mathcal{K}_4$, so $D_+(S) = D(S)$. Recall that $\tilde{D}(S) = \sum_{T \in \mathcal{K}_3(S)} D_-(T) - (2 - 4\beta)$.

Let T_1, \dots, T_4 be triangles in S with $D(T_i) \leq D(T_{i+1})$ for $1 \leq i \leq 3$. Since $D_-(T) \leq \beta$, we have

$$\tilde{D}(S) = \begin{cases} 2(4\beta - 1) & \text{if } k_3^+(S) = 4, \\ 4\beta - 1 + (D(T_1) - (1 - 3\beta)) & \text{if } k_3^+(S) = 3, \\ D(T_1) + D(T_2) - 2(1 - 3\beta) & \text{if } k_3^+(S) = 2, \end{cases} \quad (28)$$

where $k_3^+(S)$ is the number of heavy triangles in S . Also recall that $D(T) \geq 1 - 3\beta$ by Lemma 4.1 (i). We will often make reference to these formulae throughout this section.

Proof of Lemma 6.3. Define the function $\eta : \mathcal{K}_4 \rightarrow \mathbb{R}$ to be

$$\eta(S) = \tilde{D}(S) - \frac{4\beta - 1}{29 - 75\beta} \sum_{T \in \mathcal{K}_3(S)} \frac{D_+(T)}{D_+(T) + \beta}$$

for $S \in \mathcal{K}_4$. Recall that for a heavy triangle T , $D(T) = D_+(T) + \beta$. Thus, only heavy 3-cliques in S contribute to $\sum D_+(T)/(D_+(T) + \beta)$. We now claim that it is enough to show that $\sum_{S \in \mathcal{K}_4} \eta(S) \geq 0$. If $\sum_{S \in \mathcal{K}_4} \eta(S) \geq 0$, then Lemma 4.6 with $t = 3$ implies that

$$\begin{aligned} 0 &\leq \sum_{S \in \mathcal{K}_4} \eta(S) = \sum_{S \in \mathcal{K}_4} \tilde{D}(S) - \frac{4\beta - 1}{29 - 75\beta} n \sum_{T \in \mathcal{K}_3} D_+(T) \\ &\leq (2 - 4\beta)k_4 - (1 - 3\beta)\beta nk_3 - \left(1 - 3\beta + \frac{4\beta - 1}{29 - 75\beta}\right) n \sum_{T \in \mathcal{K}_3} D_+(T) \\ &\quad + 2 \sum_{S \in \mathcal{K}_4} D_+(S) - 10k_5/n, \end{aligned}$$

where the last inequality is due to Lemma 4.6 with $t = 3$. Observe that $\sum_{S \in \mathcal{K}_4} D_+(S) = \sum_{S \in \mathcal{K}_4} D(S) = 5k_5/n$, so the terms with $\sum D_+(S)$ and k_5/n cancel. Rearranging the inequality, we obtain the inequality in Lemma 6.3.

Suppose $\sum_{S \in \mathcal{K}_4} \eta(S) < 0$. Then, there exists a 4-clique S with $\eta(S) < 0$. Such a 4-clique is called *bad*, otherwise it is called *good*. The sets of bad and good 4-cliques are denoted by \mathcal{K}_4^{bad} and \mathcal{K}_4^{good} respectively. In the next claim, we identify the structure of a bad 4-clique.

Claim 7.1. *Let S be a bad 4-clique. Let*

$$\Delta = (1 - 3\beta)(1 + \epsilon) \text{ and } \epsilon = (4\beta - 1)/(150\beta^2 - 137\beta + 30).$$

Then, the following hold

- (i) S contains exactly one heavy edge and two heavy triangles,
- (ii) $0 < D(S) < \Delta$,

(iii) $D(T) + D(T') < 2\Delta$, where T and T' are the two non-heavy triangles in S .

Proof. Let T_1, \dots, T_4 be triangles in S with $D(T_i) \leq D(T_{i+1})$ for $1 \leq i \leq 3$. We may assume that $D_+(T_4) > 0$, otherwise S is good by Corollary 4.5 as $\eta(S) = \tilde{D}(S) \geq 0$. Hence, S is also heavy by Lemma 4.1 (iv). We separate cases by the number of heavy triangles in S .

First, suppose all triangles are heavy. Hence, $\tilde{D}(S) = 2(4\beta - 1)$ by (28). Clearly, $D_+(T_i) \leq 1 - \beta$ for $1 \leq i \leq 4$, so

$$\begin{aligned} \eta(S) &\geq 2(4\beta - 1) - \frac{4\beta - 1}{29 - 75\beta} \sum_{T \in \mathcal{K}_3(S)} \frac{D_+(T)}{D_+(T) + \beta} \\ &\geq 2(4\beta - 1) \left(1 - \frac{2(1 - \beta)}{29 - 75\beta} \right) = \frac{2(4\beta - 1)(27 - 73\beta)}{29 - 75\beta} \geq 0. \end{aligned}$$

This contradicts the assumption that S is bad. Thus, not all triangles in S are heavy, so $0 < D(S) \leq \beta$ by Lemma 4.1 (v). Also, $D_+(T) \leq D_+(S) = D(S) \leq \beta$.

Suppose all but one triangles are heavy, so $\tilde{D}(S) \geq 4\beta - 1$ by (28). Hence,

$$\begin{aligned} \eta(S) &\geq 4\beta - 1 - \frac{4\beta - 1}{29 - 75\beta} \sum_{T \in \mathcal{K}_3(S)} \frac{D_+(T)}{D_+(T) + \beta} \\ &\geq (4\beta - 1) \left(1 - \frac{3}{29 - 75\beta} \frac{D(S)}{D(S) + \beta} \right) \\ &\geq (4\beta - 1) \left(1 - \frac{3}{2(29 - 75\beta)} \right) = \frac{5(4\beta - 1)(11 - 30\beta)}{2(29 - 75\beta)} \geq 0, \end{aligned}$$

which is a contradiction.

Suppose there is only one heavy triangle, T_4 , in S . Corollary 4.3 implies that $\tilde{D}(S) + D_+(T_4) \geq 2D_+(S) = 2D(S)$. Note that $D_+(T_4) \leq D_+(S) = D(S)$, so $\tilde{D}(S) \geq D(S)$. Thus,

$$\begin{aligned} \eta(S) &\geq D(S) - \frac{4\beta - 1}{29 - 75\beta} \frac{D_+(T_4)}{D_+(T_4) + \beta} \geq D(S) - \frac{4\beta - 1}{29 - 75\beta} \frac{D(S)}{D(S) + \beta} \\ &= \left(1 - \frac{4\beta - 1}{(29 - 75\beta)(D(S) + \beta)} \right) D(S) \geq \left(1 - \frac{4\beta - 1}{(29 - 75\beta)\beta} \right) D(S) > 0. \end{aligned}$$

Hence, S has exactly two heavy triangles, namely T_3 and T_4 .

If $D(S) \geq \Delta$, then

$$\begin{aligned}
\eta(S) &= D(T_1) + D(T_2) - 2(1 - 3\beta) - \frac{4\beta - 1}{29 - 75\beta} \left(\frac{D_+(T_3)}{D_+(T_3) + \beta} + \frac{D_+(T_4)}{D_+(T_4) + \beta} \right) \\
&\geq 2(D(S) - (1 - 3\beta)) - \frac{2(4\beta - 1)}{29 - 75\beta} \frac{D(S)}{D(S) + \beta} \\
&> 2(1 - 3\beta)\epsilon - \frac{2(4\beta - 1)}{29 - 75\beta} \frac{\Delta}{\Delta + \beta} \\
&\geq 2(1 - 3\beta)\epsilon - \frac{2(4\beta - 1)\Delta}{(29 - 75\beta)(1 - 2\beta)} = 0.
\end{aligned}$$

Thus, $D(S) < \Delta$. If $D(T_1) + D(T_2) \geq 2\Delta$, then $\tilde{D}(S) \geq 2(\Delta - (1 - 3\beta)) = 2(1 - 3\beta)\epsilon$ by (28). Moreover, since $D_+(T_i) \leq D(S) < \Delta$ for $i = 3, 4$,

$$\eta(S) > 2(1 - 3\beta)\epsilon - \frac{2(4\beta - 1)}{29 - 75\beta} \frac{\Delta}{\Delta + \beta} \geq 2(1 - 3\beta)\epsilon - \frac{2(4\beta - 1)\Delta}{(29 - 75\beta)(1 - 2\beta)} = 0.$$

Thus, (iii) is true.

We have shown that S contains exactly two heavy triangles. Therefore, to prove (i), it is sufficient to prove that S contains exactly one heavy edge. A triangle containing a heavy edge is heavy by Lemma 4.1 (iv). Since S contains two heavy triangle, there is at most one heavy edge in S . It is enough to show that if S does not contain any heavy edge and $D(S) < \Delta$, then S is good, which is a contradiction. Assume that S contains no heavy edge. Let $e_i = T_i \cap T_4$ be an edge of T_4 for $i = 1, 2, 3$. We claim that $\tilde{D}(S) \geq D_+(T_4)$. By Corollary 4.3 taking $S = T_4$ and $t = 2$, we obtain

$$\begin{aligned}
D(e_1) + D(e_2) + D(e_3) &\geq 2 - 3\beta + D(T_4) \\
D(e_1) + D(e_2) &\geq 2 - 4\beta + D_+(T_4).
\end{aligned}$$

as $D(e_3) \leq 2\beta$ and $D_-(T_4) = \beta$. By Lemma 4.1 (ii), we get

$$D(T_1) + D(T_2) \geq D(e_1) + D(e_2) - 2\beta \geq 2(1 - 3\beta) + D_+(T_4).$$

Hence, $\tilde{D}(S) \geq D_+(T_4)$ by (28). Therefore,

$$\begin{aligned}
\eta(S) &\geq D_+(T_4) - \frac{4\beta - 1}{29 - 75\beta} \sum_{T \in \mathcal{K}_4(S)} \frac{D_+(T)}{D_+(T) + \beta} \\
&\geq \left(1 - \frac{2(4\beta - 1)}{(29 - 75\beta)(D_+(T_4) + \beta)} \right) D_+(T_4) \\
&\geq \left(1 - \frac{2(4\beta - 1)}{(29 - 75\beta)\beta} \right) D_+(T_4) > 0
\end{aligned}$$

and so S is good, a contradiction. This completes the proof of the claim. \square

Since a bad 4-clique S must be heavy, that is, $D(S) > 0$, it is contained in some 5-clique. A 5-clique is called *bad* if it contains at least one bad 4-clique. We denote \mathcal{K}_5^{bad} to be the set of bad 5-cliques. Define $\tilde{\eta}(S)$ to be $\eta(S)/D(S)$ for $S \in \mathcal{K}_4$ with $D(S) > 0$. Clearly,

$$n \sum_{S \in \mathcal{K}_4} \eta(S) = \sum_{U \in \mathcal{K}_5} \sum_{S \in \mathcal{K}_4(U)} \tilde{\eta}(S) + n \sum_{S \in \mathcal{K}_4: D(S)=0} \eta(S). \quad (29)$$

Recall that our aim is to show that $\sum_{S \in \mathcal{K}_4} \eta(S) \geq 0$. Since $D(S) = 0$ implies that S is good, we have $\eta(S) \geq 0$. Hence, it is enough to show that $\sum_{S \in \mathcal{K}_4(U)} \tilde{\eta}(S) \geq 0$ for each bad 5-clique U .

Now, we give a lower bound on $\tilde{\eta}(S)$ for bad 4-cliques S . By Claim 7.1,

$$\eta(S) \geq -\frac{4\beta - 1}{29 - 75\beta} \sum_{T \in \mathcal{K}_3(S)} \frac{D_+(T)}{D_+(T) + \beta} \geq -\frac{2(4\beta - 1)}{29 - 75\beta} \frac{D(S)}{D(S) + \beta}.$$

Hence,

$$\tilde{\eta}(S) \geq -\frac{2(4\beta - 1)}{(29 - 75\beta)(D(S) + \beta)} > -\frac{2(4\beta - 1)}{(29 - 75\beta)\beta}. \quad (30)$$

Next, we are going to bound $D(S)$ above for $S \in \mathcal{K}_4(U) \setminus \mathcal{K}_4^{bad}$ and $U \in \mathcal{K}_5^{bad}$. Let $S^b \in \mathcal{K}_4^{bad}(U)$. Observe that $S \cap S^b$ is a 3-clique. Then, by Lemma 4.1 and Claim 7.1, we have

$$D(S) \leq D(S \cap S^b) = D_+(S \cap S^b) + \beta \leq D(S^b) + \beta < \Delta + \beta. \quad (31)$$

Recall that a bad 4-clique S contains a heavy edge by Claim 7.1 and hence so does a bad 5-clique U . We split \mathcal{K}_5^{bad} into subcases depending on the number of heavy edges in U . The next claim studies the relationship between the number of heavy edges and bad 4-cliques in a bad 5-clique U .

Claim 7.2. *Let $U \in \mathcal{K}_5^{bad}$ with $h \geq 2$ heavy edges and b bad 4-cliques. Then $b \leq 2h/(h - 1) = 2 + 2/(h - 1)$. Moreover, if there exist two heavy edges sharing a common vertex, $b \leq 3$.*

Proof. Define H to be the graph induced by the heavy edges in U . Write u_S for the vertex in U not in $S \in \mathcal{K}_4(U)$. This defines a bijection between $V(U)$ and $\mathcal{K}_4(U)$. If S is bad, u_S is adjacent to all but one heavy edges by Claim 7.1. By summing the degrees of H , $2h = \sum_{S \in \mathcal{K}_4(U)} d(u_S) \geq b(h - 1)$. Thus, $b \leq 2h/(h - 1)$.

If there exist two heavy edges sharing a common vertex in H , then every bad 4-clique must miss one of the vertices of these two heavy edges. Hence, $b \leq 3$. \square

Claim 7.3. Let $U \in \mathcal{K}_5^{bad}$ with two heavy edges. Then $\sum_{S \in \mathcal{K}_4(U)} \tilde{\eta}(S) > 0$.

Proof. Let e and e' be two heavy edges in U , and let b be the number of bad 4-cliques in U . We consider the cases whether e and e' are vertex disjoint or not separately. First, assume that e and e' are vertex disjoint. Notice that $\sum_{S \in \mathcal{K}_4^{bad}(U)} \tilde{\eta}(S) > -b\gamma$ by (30), where $\gamma = 2(4\beta - 1)/(29 - 75\beta)\beta$ and $b \leq 4$ by Claim 7.2. Also, there is exactly one heavy 4-clique S containing both e and e' . Therefore, it is sufficient to prove that $\eta(S) \geq bD(S)\gamma$. Since S contains two disjoint heavy edges, all triangles in S are heavy by Lemma 4.1 (iv). Thus, $\tilde{D}(S) = 2(4\beta - 1)$ by (28). Observe that $T = S \cap S'$ is a triangle for $S' \in \mathcal{K}_4(U) \setminus S$. Moreover, $D_+(T) \leq D_+(S') = D(S')$ by Lemma 4.1 (iii). Hence,

$$\begin{aligned} \eta(S) &\geq 2(4\beta - 1) - \frac{4\beta - 1}{29 - 75\beta} \sum_{S' \in \mathcal{K}_4(U) \setminus S} \frac{D(S')}{D(S') + \beta} \\ &> (4\beta - 1) \left(2 - \frac{1}{29 - 75\beta} \left(\frac{b\Delta}{\Delta + \beta} + \frac{(4-b)(\Delta + \beta)}{\Delta + 2\beta} \right) \right) \end{aligned}$$

by Claim 7.1 (ii) and (31). Therefore, $\eta(S) - bD(S)\gamma$ is at least

$$\begin{aligned} &(4\beta - 1) \left(2 - \frac{1}{29 - 75\beta} \left(\frac{b\Delta}{\Delta + \beta} + \frac{(4-b)(\Delta + \beta)}{\Delta + 2\beta} \right) \right) - b(\Delta + \beta)\gamma \\ &\geq (4\beta - 1) \left(2 - \frac{4\Delta}{(29 - 75\beta)(\Delta + \beta)} \right) - 4(\Delta + \beta)\gamma > 0. \end{aligned}$$

Thus, if U contains two vertex disjoint heavy edges, $\sum_{S \in \mathcal{K}_4(U)} \tilde{\eta}(S) > 0$. Similar argument also holds for the case when e and e' share a common vertex. \square

Recall that a bad 5-clique contains at least one heavy edge. Thus, we are left with the case $U \in \mathcal{K}_5^{bad}$ containing exactly one heavy edge.

Claim 7.4. Suppose $U \in \mathcal{K}_5^{bad}$ with exactly one heavy edge e . Then, $\sum_{S \in \mathcal{K}_4(U)} \tilde{\eta}(S) > 0$.

Proof. Let u_1, \dots, u_5 be the vertices of U with u_4u_5 is the heavy edge. Write S_i and η_i to be $U - u_i$ and $\eta(S_i)$ respectively for $1 \leq i \leq 5$. Similarly write $T_{i,j}$ to be $U - u_i - u_j$ for $1 \leq i < j \leq 5$. Recall that a bad 4-clique contains a heavy edge by Claim 7.1 (i). Hence, S_i is a bad 4-clique only if $i \leq 3$. Without loss of generality, S_1, \dots, S_b are the bad 4-cliques in U .

Since S_3 contains a heavy edge, it contains at least 2 heavy triangles by Lemma 4.1 (iv). If S_3 contains either three or four heavy

triangles, then S_3 is not bad by Claim 7.1 (i). By a similar argument as in the proof of Claim 7.3, we can deduce that $\eta_3 \geq 2\gamma D(S_3)$, where as before $\gamma = 2(4\beta - 1)/(29 - 75\beta)\beta$. Therefore, $\sum_{S \in \mathcal{K}_4(U)} \tilde{\eta}(S) > 0$ as $b \leq 2$. Thus, we may assume that there are exactly two heavy triangles in S_i for $1 \leq i \leq 3$. By Lemma 4.1 (v), $D(S_i) < \beta$ for $1 \leq i \leq 3$. For $1 \leq i \leq b$,

$$D(T_{i,4}) + D(T_{i,5}) < 2\Delta = 2(1 - 3\beta)(1 + \epsilon)$$

by Claim 7.1 (iii). For $b < i \leq 3$, $\tilde{D}(S_i) = D(T_{i,4}) + D(T_{i,5}) - 2(1 - 3\beta)$ by (28). Thus,

$$\begin{aligned} D(T_{i,4}) + D(T_{i,5}) &= \eta_i + 2(1 - 3\beta) + \frac{4\beta - 1}{29 - 75\beta} \sum_{T \in \mathcal{K}_3(S_i)} \frac{D_+(T)}{D_+(T) + \beta} \\ &\leq \eta_i + 2(1 - 3\beta) + \frac{\gamma\beta D(S_i)}{D(S_i) + \beta} \\ &\leq \eta_i + 2(1 - 3\beta) + \gamma\beta/2. \end{aligned}$$

After applying Corollary 4.5 to S_4 and S_5 taking $t = 3$, and adding the two inequalities together, we obtain

$$\begin{aligned} 2(2 - 4\beta) &\leq \sum_{1 \leq i \leq 3} (D_-(T_{i,4}) + D_-(T_{i,5})) + 2D_-(T_{4,5}) \\ 2(2 - 5\beta) &\leq \sum_{1 \leq i \leq b} (D(T_{i,4}) + D(T_{i,5})) + \sum_{b < i \leq 3} (D(T_{i,4}) + D(T_{i,5})) \\ &< 2b(1 - 3\beta)(1 + \epsilon) + \sum_{b < i \leq 3} \eta_i + (3 - b)(2(1 - 3\beta) + \gamma\beta/2) \\ 2(4\beta - 1) &< 2b(1 - 3\beta)\epsilon + \sum_{b < i \leq 3} \eta_i + (3 - b)\gamma\beta/2 \end{aligned} \quad (32)$$

If $b = 3$, the above inequality becomes $2(4\beta - 1) < 6(1 - 3\beta)\epsilon < 2(4\beta - 1)$, which is a contradiction. Thus, $b \leq 2$. Notice that $\eta_i > -D(S_i)\gamma > -\gamma$ for $1 \leq i \leq b$. Hence, $\sum_{S \in \mathcal{K}_4^{bad}(U)} \tilde{\eta}(S) > -b\gamma$. Also, recall that $D(S_i) \leq \beta$ for $1 \leq i \leq 3$. It is enough to show that $\sum_{b < i \leq 3} \eta_i \geq b\gamma\beta$. Suppose the contrary, so $\sum_{b < i \leq 3} \eta_i < b\gamma\beta$. Then, (32) becomes

$$2(4\beta - 1) < 2b(1 - 3\beta)\epsilon + (3 + b)\gamma\beta/2 \leq 4(1 - 3\beta)\epsilon + 5\gamma\beta/2 < 2(4\beta - 1),$$

which is a contradiction. The proof of the claim is complete. \square

Hence, by Claim 7.3 and Claim 7.4, (29) becomes $\sum_{S \in \mathcal{K}_4} \eta(S) \geq 0$, so the inequality in Lemma 6.3 holds. Now suppose equality holds in Lemma 6.3. Claim 7.3 and Claim 7.4 imply that no 5 clique is bad, so no 4-clique is bad. Furthermore, we must have $\eta(S) = 0$ for

all $S \in \mathcal{K}_4$. It can be checked that if the definition of a bad 4-clique includes heavy 4-cliques S with $\eta(S) = 0$, then all arguments still hold. Thus, we can deduce that G is K_5 -free. Hence, G is also K_5 -free. By Theorem 4 taking $s = 4$ and $t = 3$, we obtain that (n, β) is feasible and $G \in \mathcal{G}(n, \beta)$. \square

8 Proof of Theorem 2.4

Here, we prove Theorem 2.4. Since the proof of theorem uses similar arguments in the proof of Theorem 5.1 and Lemma 6.1, we only give a sketch of the proof.

Sketch of Proof of Theorem 2.4. For $2 \leq t \leq p$ and $1/(p+1) \leq \beta < 1/p$, define

$$\begin{aligned} A_t^p(\beta) &= (t-1)((p+1)\beta - 1)C_t^p(\beta), \text{ and} \\ B_t^p(\beta) &= ((p+1)\beta - 1)C_t^p(\beta), \end{aligned}$$

where $C_j(\beta)$ satisfies the recurrence

$$C_t(\beta) + 1 = (p-t+1)\beta C_{t-1}(\beta)$$

with the initial condition $C_p(\beta) = 0$ for $1/(p+1) \leq \beta < 1/p$. Explicitly, $C_{p-j}^p(\beta) = \sum_{0 \leq i < j} i! \beta^{i-j} / j!$ for $0 \leq j \leq p-2$. These functions will be used as coefficients in corresponding statements of Lemma 6.1 for $2 \leq t < p$. Define the integer $r(\beta)$ to be the smallest integer at least 2 such that for $r \leq t \leq p$, $A_t^p(\beta) < 1$ and $B_t^p(\beta) < (p-t)\beta$. Let

$$\beta_p = \sup\{\beta_0 : r(\beta) = 2 \text{ for all } 1/(p+1) \leq \beta < \beta_0\}$$

and $\epsilon_p = \beta_p - 1/(p+1)$. Observe that $A_t(\beta)$, $B_t(\beta)$ and $C_t(\beta)$ are right continuous functions of β . Moreover, both $A_t(\beta)$ and $B_t(\beta)$ tend to zero as β tends $1/(p+1)$ from above, so $\beta_p > 1/(p+1)$ and $\epsilon_p > 0$. By mimicking the poof of Lemma 6.1, we have

$$\tilde{D}(S) \geq A_{t+1}^p(\beta)D_+(S) - B_t^p(\beta) \sum_{T \in \mathcal{K}_t(S)} \frac{D_+(T)}{D(T)}$$

for $S \in \mathcal{K}_{t+1}$, $1/(p+1) \leq \beta < \beta_p$ and $2 \leq t \leq p$. Then, following the arguments in the proof of Theorem 5.1, we can deduce that

$$\frac{k_s(G)}{g_s(\beta)n^s} \geq \frac{k_t(G)}{g_t(\beta)n^t} + \frac{1-t\beta - B_t^p(\beta)}{(1-t\beta)(p-t+1)\beta g_t(\beta)n^t} \sum_{T \in \mathcal{K}_t} D_+(T)$$

for $2 \leq t < s \leq p+1$ and $1/(p+1) < \beta \leq \beta_p$. Since $1-t\beta - B_t^p(\beta) \geq 0$, the proof of theorem is completed. \square

Clearly, ϵ_p defined in the proof is not optimal. Generalising the proof of Lemma 6.3 would lead to an improvement on ϵ_p .

9 Counting $(p + 1)$ -cliques

In this section, we are going to prove the below theorem, which implies Theorem 2.5.

Theorem 9.1. *Let $0 < \beta < 1$ and $p = \lceil \beta^{-1} \rceil - 1$. Suppose G is a graph of order n with minimum degree $(1 - \beta)n$. Then, for any integer $2 \leq t \leq p$,*

$$\frac{k_{p+1}(G)}{g_{p+1}(\beta)n^{p+1}} \geq \frac{k_t(G)}{g_t(\beta)n^t}.$$

Moreover, for $t = 2$, equality holds if and only if (n, β) is feasible, and G is a member of $\mathcal{G}(n, \beta)$.

For positive integers $2 \leq t \leq s \leq p + 1$, define the function $\phi_t^s : \mathcal{K}_s \rightarrow \mathbb{R}$ such that

$$\phi_t^s(S) = \begin{cases} D_-(S) & \text{if } t = s, \text{ and} \\ \sum_{U \in \mathcal{K}_{s-1}(S)} \phi_t^{s-1}(U) & \text{if } t < s \end{cases}$$

for $S \in \mathcal{K}_s$. Observe that for $G_0 \in \mathcal{G}(n, \beta)$ with (n, β) feasible,

$$\phi_t^s(S) = \begin{cases} (s-t)!(1-t\beta) & \text{if } |V(S) \cap V_0| = 0, 1 \\ (1-t\beta)s!/t! + ((p+1)\beta-1)(s-2)!/(t-2)! & \text{if } |V(S) \cap V_0| = 2 \end{cases}$$

for s -cliques S in G_0 . Let $\Phi_t^s(S) = \min\{\phi_t^s(S), \varphi_t^s\}$ for $S \in \mathcal{K}_s$ and $2 \leq t \leq s \leq p + 1$, where

$$\varphi_t^s = (1-t\beta)s!/t! + ((p+1)\beta-1)(s-2)!/(t-2)!$$

to be the analogue of D_- for ϕ_t^s . The next lemma gives a lower bound on $\Phi_t^s(S)$ for $S \in \mathcal{K}_s$.

Lemma 9.2. *Let $0 < \beta < 1$ and $p = \lceil \beta^{-1} \rceil - 1$. Let G be a graph of order n with minimum degree $(1 - \beta)n$. Then,*

$$\Phi_t^s(S) \geq (1-t\beta)s!/t! + (D_-(S) - (1-s\beta))(s-2)!/(t-2)!$$

for $S \in \mathcal{K}_s$ and $2 \leq t < s \leq p + 1$. In particular, for $s = p + 1$ and $t = p$

$$\sum_{S \in \mathcal{K}_{p+1}} \Phi_t^{p+1}(S) \geq \left((1-t\beta) \frac{(p+1)!}{t!} - (1-(p+1)\beta) \frac{(p-1)!}{(t-2)!} \right) k_{p+1}. \quad (33)$$

Proof. Fix β and t and we proceed by induction on s . The inequality holds for $s = t + 1$ by Corollary 4.5. Suppose $s \geq t + 2$ and that the lemma is true for $t, \dots, s - 1$. Hence

$$\phi_t^s(S) = \sum_{T \in \mathcal{K}_{s-1}(S)} \phi_t^{s-1}(T) \geq \sum_{T \in \mathcal{K}_{s-1}(S)} \Phi_t^{s-1}(T)$$

and by the induction hypothesis,

$$\begin{aligned} &\geq \sum_{T \in \mathcal{K}_{s-1}(S)} \left((1-t\beta) \frac{(s-1)!}{t!} + (D_-(T) - (1-(s-1)\beta)) \frac{(s-3)!}{(t-2)!} \right) \\ &= (1-t\beta) \frac{s!}{t!} + \frac{(s-3)!}{(t-2)!} \left(\sum_{T \in \mathcal{K}_{s-1}(S)} D_-(T) - s(1-(s-1)\beta) \right) \\ &\geq (1-t\beta) \frac{s!}{t!} + (D_-(S) - (1-s\beta)) \frac{(s-2)!}{(t-2)!}, \end{aligned}$$

where the last inequality comes from Corollary 4.5 with $t = s - 1$. The right hand side is increasing in $D_-(S)$. In addition, the right hand side equals to φ_t^s only if $D_-(S) = (p - s + 1)\beta$. Thus, the proof of the lemma is complete. \square

Now, we bound $\sum_{S \in \mathcal{K}_s} \Phi_t^s(S)$ from above using Proposition 3.1 to obtain the next lemma. The proof is essentially a straightforward application of Proposition 3.1 with an algebraic check.

Lemma 9.3. *Let $0 < \beta < 1$ and $p = \lceil \beta^{-1} \rceil - 1$. Let G be a graph of order n with minimum degree $(1 - \beta)n$. Then, for $2 \leq t \leq s \leq p + 1$*

$$\begin{aligned} \sum_{S \in \mathcal{K}_s} \Phi_t^s(S) &\leq \varphi_t^{s-1} s k_s + 2((p+1)\beta - 1) \sum_{i=t+1}^{s-1} \left(\frac{(i-3)!}{(t-2)!} k_i n^{s-i} \prod_{j=i}^{s-1} (1-j\beta) \right) \\ &\quad + ((t+1)k_{t+1} - (p-t+1)\beta k_t n) n^{s-t-1} \prod_{j=t}^{s-1} (1-j\beta). \end{aligned}$$

Proof. Fix β and t . We proceed by induction on s . Suppose $s = t + 1$. Note that $\Phi_t^{t+1}(S) \leq \sum_{T \in \mathcal{K}_t(S)} D_-(T)$. By Proposition 3.1, taking $\mathcal{A} = \mathcal{K}_t$, $f = D_-$, $g = D$, $M = (p - t + 1)\beta$ and $m = 1 - t\beta$,

$$\begin{aligned} \sum_{S \in \mathcal{K}_{t+1}} \Phi_t^{t+1}(S) &\leq \sum_{S \in \mathcal{K}_{t+1}} \sum_{T \in \mathcal{K}_t(S)} D_-(T) = n \sum_{T \in \mathcal{K}_t} D(T) D_-(T) \\ &\leq (p-t+1)\beta n \sum_{T \in \mathcal{K}_t} D(T) + (1-t\beta)n \sum_{T \in \mathcal{K}_t} D_-(T) - (1-t\beta)(p-t+1)\beta n k_t \\ &\leq (t+1)(1 - (p-2t+1)\beta) k_{t+1} - (1-t\beta)(p-t+1)\beta n k_t. \end{aligned}$$

Hence, the lemma is true for $s = t + 1$. Now assume that $s \geq t + 2$ and the lemma is true up to $s - 1$. By Proposition 3.1 taking $\mathcal{A} = \mathcal{K}_t$, $f = \Phi_t^{s-1}$, $g = D$, $M = \varphi_t^{s-1}$ and $m = 1 - (s - 1)\beta$, we have

$$\begin{aligned} \sum_{S \in \mathcal{K}_s} \Phi_t^s(S) &= n \sum_{T \in \mathcal{K}_{s-1}} D(T) \Phi_t^{s-1}(T) \\ &\leq \varphi_t^{s-1} \sum_{T \in \mathcal{K}_{s-1}} nD(T) + (1 - (s - 1)\beta)n \sum_{T \in \mathcal{K}_{s-1}} \Phi_t^{s-1}(T) - \varphi_t^{s-1}(1 - (s - 1)\beta)nk_{s-1} \\ &= \varphi_t^{s-1}sk_s + (1 - (s - 1)\beta)n \sum_{T \in \mathcal{K}_{s-1}} \Phi_t^{s-1}(T) - \varphi_t^{s-1}(1 - (s - 1)\beta)nk_{s-1}. \end{aligned}$$

Next, we apply induction hypothesis on $\sum \Phi_t^{s-1}(T)$. Note that

$$(s - 1)\varphi_t^{s-2} - \varphi_t^{s-1} = 2((p + 1)\beta - 1)(s - 4)!/(t - 2)!.$$

After collecting the terms, we obtain the desire inequality. \square

Now we are ready to prove Theorem 9.1. The proof is very similar to the proof of Theorem 5.1.

Proof of Theorem 9.1. We fix β and write g_t to be $g_t(\beta)$. We proceed by induction on t from above. The theorem is true for $t = p$ by Lemma 9.2 and Lemma 9.3. Hence, we may assume $t < p$. By Lemma 9.3,

$$\begin{aligned} \sum \Phi_t^{p+1}(S) &\leq (p + 1)\varphi_t^p k_{p+1} + 2((p + 1)\beta - 1) \sum_{i=t+1}^p \left(\frac{(i - 3)!}{(t - 2)!} k_i n^{p+1-i} \prod_{j=i}^p (1 - j\beta) \right) \\ &\quad + ((t + 1)k_{t+1} - (p - t + 1)\beta nk_t) n^{p-t} \prod_{j=t}^p (1 - j\beta), \end{aligned}$$

and by the induction hypothesis

$$\begin{aligned} &\leq (p + 1)\varphi_t^p k_{p+1} + 2((p + 1)\beta - 1) \sum_{i=t+1}^p \left(\frac{k_{p+1} g_i (i - 3)!}{g_{p+1} (t - 2)!} \prod_{j=i}^p (1 - j\beta) \right) \\ &\quad + \left((t + 1) \frac{k_{p+1}}{g_{p+1}} g_{t+1} - (p - t + 1)\beta nk_t \right) n^{p-t} \prod_{j=t}^p (1 - j\beta). \end{aligned}$$

Substitute the above inequality into (33) and rearranging to obtain the desire inequality.

Now suppose that equality holds, so equality holds in (33). Therefore, $D(S) = D_-(S) = 0$ for all $S \in \mathcal{K}_{p+1}$. Thus, G is K_{p+2} -free. By Theorem 5.1 (n, β) is feasible, and $G \in \mathcal{G}(n, \beta)$. This completes the proof of the theorem. \square

Acknowledgements

The author is greatly indebted to Andrew Thomason for his comments and his help in making the proof clearer.

References

- [1] B. Andrásfai, P. Erdős, and V. T. Sós, *On the connection between chromatic number, maximal clique and minimal degree of a graph*, Discrete Math. **8** (1974), 205–218.
- [2] B. Bollobás, *On complete subgraphs of different orders*, Math. Proc. Cambridge Philos. Soc. **79** (1976), 19–24.
- [3] P. Erdős, *On the number of complete subgraphs and circuits contained in graphs.*, Časopis Pěst. Mat. (1969), 290–296.
- [4] D. Fisher, *Lower bounds on the number of triangles in a graph*, Journal of Graph Theory **13** (1989), 505–512.
- [5] A. S. L. Lo, *Triangles in regular graphs with density below one half*, Combinatorics Probability and Computing **18** (2009), 435–440.
- [6] A. S. L. Lo, *Cliques in graphs*, Ph.D. thesis, University of Cambridge, 2010.
- [7] L. Lovász and M. Simonovits, *On complete subgraphs of a graph ii*, Studies Pure Math. (1983), 459–496.
- [8] V. Nikiforov, *The number of cliques in graphs of given order and size*, <https://umdrive.memphis.edu/vnikifrv/public/pdfs/Ni07n.pdf>, Preprint.
- [9] A. Razborov, *On the minimal density of triangles in graphs*, Combinatorics, Probability and Computing (2008), 603–618.
- [10] A. A. Razborov, *Flag algebras*, J. Symbolic Logic **72** (2007), 1239–1282.
- [11] P. Turán, *Eine Extremalaufgabe aus der Graphentheorie*, Mat. Fiz. Lapok **48** (1941), 436–452.