

OPERATOR ALGEBRA QUANTUM GROUPS OF UNIVERSAL GAUGE GROUPS

SNIGDHAYAN MAHANTA AND VARGHESE MATHAI

ABSTRACT. In this paper, we quantize universal gauge groups such as $SU(\infty)$, in the σ - C^* -algebra setting. More precisely, we propose a concise definition of σ - C^* -quantum groups and explain the concept here. At the same time, we put this definition in the mathematical context of countably compactly generated groups as well as C^* -compact quantum groups.

If H is a compact and Hausdorff topological group, then the C^* -algebra of all continuous functions $C(H)$ admits a comultiplication map $\Delta : C(H) \rightarrow C(H) \hat{\otimes} C(H)$ arising from the multiplication in H . This observation motivated Woronowicz (see, for instance, [12]), amongst others such as Soibelman [10], to introduce the notion of a C^* -compact quantum group in the setting of operator algebras as a unital C^* -algebra with a coassociative comultiplication, satisfying a few other conditions. If the group H is only locally compact then the situation becomes significantly more difficult. One of the reasons is that the multiplication map $m : H \times H \rightarrow H$ is no longer a proper map and one needs to introduce multiplier algebras of C^* -algebras to obtain a comultiplication, see for instance, Kustermans-Vaes [5]. For an excellent and thorough introduction to this theory the readers are referred to, for instance, [6]. In the sequel we show that if $H = \varinjlim_n H_n$ is a countably compactly generated group, i.e., if $H_n \subset H_{n+1}$ are compact and Hausdorff topological groups for all $n \in \mathbb{N}$ and if H is the direct limit, then a story similar to the compact group case goes through using the general framework of σ - C^* -algebras as systematically developed by Phillips [8, 9], motivated by some earlier work by Arveson, Mallios, Voiculescu, amongst others. There is a clean formulation of, what we call, σ - C^* -quantum groups, which are noncommutative generalizations of $C(H)$. Examples of countably compactly generated groups are $U(\infty) = \varinjlim_n U(n)$, $SU(\infty) = \varinjlim_n SU(n)$, where $U(n)$ (resp. $SU(n)$) are the unitary (resp. special unitary) groups. They are also known in the physics literature as *universal gauge groups*, see Harvey-Moore [4] and Carey-Mickelsson [3]. Such spaces are not locally compact and hence the existing literature on quantum groups cannot handle them. Moreover, locally compact groups that are not compact, are also not countably compactly generated. We also discuss in detail the interesting example of the quantum version of the universal special unitary group, $C(SU_q(\infty))$.

A *pro C^* -algebra* is an inverse limit of C^* -algebras and $*$ -homomorphisms, where the inverse limit is constructed inside the category of all topological $*$ -algebras and continuous $*$ -homomorphisms. For the general theory of topological $*$ -algebras one may refer to, for instance, [7]. The underlying topological $*$ -algebra of a pro C^* -algebra is necessarily complete

2010 *Mathematics Subject Classification.* 58B32, 58B34, 46L65.

Key words and phrases. σ - C^* -quantum groups, universal gauge groups, operator algebras.

Acknowledgements. Both authors gratefully acknowledge support under the Australian Research Council's *Discovery Projects* funding scheme.

and Hausdorff. It is not a C^* -algebra in general; it would be so if, for instance, the directed set is finite. If the directed set is countable, then the inverse limit is called a σ - C^* -algebra. One can choose a linearly directed cofinal subset inside any countable directed set and the passage to a cofinal subset does not change the inverse limit. Therefore, we shall always identify a σ - C^* -algebra $A \cong \varprojlim_n A_n$, where $n \in \mathbb{N}$. The inverse limit could have also been constructed inside the category of C^* -algebras; however, the two results will not agree. For instance, if $H = \varinjlim_n H_n$ as above, then the inverse limit $\varprojlim_n C(H_n)$ inside the category of topological $*$ -algebras is $C(H)$, whereas that inside the category of C^* -algebras is $C_b(H)$, i.e., the norm bounded functions on H . It is known that $C_b(H) \cong C(\beta H)$, where βH is the Stone–Čech compactification of H . Therefore, if one wants to model a space via its algebra of all continuous functions then the former inverse limit is the appropriate one. *Henceforth, the inverse limits are always constructed inside the category of topological $*$ -algebras.* It is known that any $*$ -homomorphism between two pro C^* -algebras is automatically continuous, provided the domain is a σ - C^* -algebra (see Theorem 5.2. of [8]). Furthermore, the category of commutative and unital σ - C^* -algebras with unital $*$ -homomorphisms (automatically continuous) is contravariantly equivalent to the category of countably compactly generated and Hausdorff spaces with continuous maps via the functor $X \mapsto C(X)$ (see Proposition 5.7. of [8]). If $A \cong \varprojlim_n A_n$, $B \cong \varprojlim_n B_n$ are two σ - C^* -algebras, then the minimal tensor product is defined to be $A \hat{\otimes}_{\min} B = \varprojlim_n A_n \hat{\otimes}_{\min} B_n$. *Henceforth, $A \hat{\otimes} B$ will always denote the minimal or spatial tensor product between σ - C^* -algebras.*

If H is a countably compactly generated and Hausdorff topological group, although the multiplication map $m : H \times H \rightarrow H$ is not proper, we get an induced comultiplication map $m^* : C(H) \rightarrow C(H \times H) \cong C(H) \hat{\otimes} C(H)$, which will be coassociative owing to the associativity of m . Motivated by the definition of Woronowicz (see also Definition 1 of [5]), we propose:

Definition. *A unital σ - C^* -algebra A is called a σ - C^* -quantum group if there is a unital $*$ -homomorphism $\Delta : A \rightarrow A \hat{\otimes} A$ which satisfies coassociativity, i.e., $(\Delta \hat{\otimes} \text{id})\Delta = (\text{id} \hat{\otimes} \Delta)\Delta$ and such that the linear spaces $\Delta(A)(A \hat{\otimes} 1)$ and $\Delta(A)(1 \hat{\otimes} A)$ are dense in $A \hat{\otimes} A$.*

Lemma. *Let $\{A_n, \theta_n : A_n \rightarrow A_{n-1}\}_{n \in \mathbb{N}}$ be a countable inverse system of C^* -algebras and let $B_n \subset A_n$ be dense subsets for all n such that $\theta_n(B_n) \subset B_{n-1}$. Then $\varprojlim_n B_n$ is a dense subset of the σ - C^* -algebra $\varprojlim_n A_n$.*

Proof. The assertion follows from the Corollary to Proposition 9 in §4-4 of [2]. □

Example. *Obviously, any C^* -compact quantum group is a σ - C^* -quantum group. Let $\{A_n, \theta_n : A_n \rightarrow A_{n-1}\}_{n \in \mathbb{N}}$ be a countable inverse system of C^* -compact quantum groups with θ_n surjective and unital for all n . Furthermore, let us assume that the comultiplication homomorphisms Δ_n form a morphism of inverse systems of C^* -algebras $\{\Delta_n\} : \{A_n\} \rightarrow \{A_n \hat{\otimes} A_n\}$. Then $(A, \Delta) = (\varprojlim_n A_n, \varprojlim_n \Delta_n)$ is a σ - C^* -quantum group. Indeed, the density of the linear spaces $\Delta(A)(A \hat{\otimes} 1)$ and $\Delta(A)(1 \hat{\otimes} A)$ inside $A \hat{\otimes} A$ follow from the above Lemma.*

Our next goal is to outline the construction of the quantum universal special unitary group, $C(SU_q(\infty))$. Recall that for $q \in (0, 1)$, the C^* -algebra $C(SU_q(n))$ is the universal C^* -algebra generated by $n^2 + 2$ elements $G_n := \{u_{ij}^n : i, j = 1, \dots, n\} \cup \{0, 1\}$, which satisfy the following relations

$$(1) \quad 0^* = 0^2 = 0, \quad 1^* = 1^2 = 1, \quad 01 = 0 = 10, \quad 1u_{ij}^n = u_{ij}^n 1 = u_{ij}^n, \quad 0u_{ij}^n = u_{ij}^n 0 = 0 \text{ for all } i, j$$

$$(2) \quad \sum_{k=1}^n u_{ik}^n (u_{jk}^n)^* = \delta_{ij} 1, \quad \sum_{k=1}^n (u_{ki}^n)^* u_{kj}^n = \delta_{ij} 1$$

$$(3) \quad \sum_{i_1=1}^n \sum_{i_2=1}^n \cdots \sum_{i_n=1}^n E_{i_1 i_2 \cdots i_n} u_{j_1 i_1}^n \cdots u_{j_n i_n}^n = E_{j_1 j_2 \cdots j_n} 1$$

where

$$E_{i_1 i_2 \cdots i_n} := \begin{cases} 0 & \text{whenever } i_1, i_2, \dots, i_n \text{ are not distinct;} \\ (-q)^{\ell(i_1, i_2, \dots, i_n)}. & \end{cases}$$

Here $\delta_{ij} 1 = 0$ if $i \neq j$, where 0 denotes the element in the generating set of $C(SU_q(n))$. Moreover, $\ell(i_1, i_2, \dots, i_n)$ denotes the number of inversed pairs in the permutation (i_1, i_2, \dots, i_n) . The C^* -algebra $C(SU_q(n))$ has a C^* -compact quantum group structure with the comultiplication Δ given by

$$\Delta(0) := 0 \otimes 0, \quad \Delta(1) := 1 \otimes 1 \text{ and } \Delta(u_{ij}^n) := \sum_k u_{ik}^n \otimes u_{kj}^n.$$

There is a surjective $*$ -homomorphism $\theta_n : C(SU_q(n)) \rightarrow C(SU_q(n-1))$ defined on the generators by

$$\begin{aligned} \theta_n(x) &:= x \text{ if } x = 0, 1 \\ \theta_n(u_{ij}^n) &:= \begin{cases} u_{ij}^{n-1} & \text{if } 1 \leq i, j \leq n-1, \\ \delta_{ij} 1 & \text{otherwise,} \end{cases} \end{aligned}$$

such that the following diagram commutes for all $n \geq 2$

$$\begin{array}{ccc} C(SU_q(n)) & \xrightarrow{\Delta_n} & C(SU_q(n)) \hat{\otimes} C(SU_q(n)) \\ \theta_n \downarrow & & \downarrow \theta_n \hat{\otimes} \theta_n \\ C(SU_q(n-1)) & \xrightarrow{\Delta_{n-1}} & C(SU_q(n-1)) \hat{\otimes} C(SU_q(n-1)). \end{array}$$

One can verify this assertion by a routine calculation on the generators. Consequently, for $n \geq 2$ the families $\{C(SU_q(n)), \theta_n\}$ and $\{C(SU_q(n)) \hat{\otimes} C(SU_q(n)), \theta_n \hat{\otimes} \theta_n\}$ form countable inverse systems of C^* -algebras and $\{\Delta_n\} : \{C(SU_q(n))\} \rightarrow \{C(SU_q(n)) \hat{\otimes} C(SU_q(n))\}$ becomes a morphism of inverse systems of C^* -algebras. We construct the underlying σ - C^* -algebra of the universal quantum gauge group as the inverse limit

$$C(SU_q(\infty)) = \varprojlim_n C(SU_q(n)).$$

In fact, $C(SU_q(\infty))$ is a σ - C^* -quantum group, since it is the inverse limit of C^* -compact quantum groups, where the comultiplication Δ on $C(SU_q(\infty))$ is defined to be $\Delta = \varprojlim_n \Delta_n$ (see the Example above).

If G is a set of generators and R a set of relations, such that the pair (G, R) is *admissible* (see Definition 1.1. of [1]), then one can always construct a universal C^* -algebra $C^*(G, R)$. For instance, the universal C^* -algebra generated by the set $\{1, x\}$, subject to the relations $\{1^* = 1^2 = 1, 1x = x1 = x, x^*x = 1 = xx^*\}$, is isomorphic to $C(S^1)$. The generators and relations of $C(SU_q(n))$ described above are also admissible.

Remark. *All matrix C^* -compact quantum groups considered, for instance, in [11, 12], such that the relations put a bound on the norm of each generator, are of the form $C^*(G, R)$, where (G, R) is an admissible pair of generators and relations.*

Let $\{(G_i, R_i)\}_{i \in \mathbb{N}}$ be a countable family of admissible pairs of generators and relations, so that $C^*(G_i, R_i)$ exist for all i . Let $F(G)$ denote the associative nonunital complex $*$ -algebra (freely) generated by the concatenation of the elements of $G \amalg G^*$ and finite \mathbb{C} -linear combinations thereof, where \amalg denotes disjoint union and $G^* = \{g^* \mid g \in G\}$ (formal adjoints). We call a relation in R *algebraic* if it is of the form $f = 0$ (or can be brought to that form), where $f \in F(G)$. For instance, if $G = \{1, x\}$, then $x^*x = 1$ is algebraic, whereas $\|x\| \leq 1$ is not. If (G, R) is a pair of generators and relations, then a *representation* ρ of (G, R) in a (pro) C^* -algebra B is a set map $\rho : G \rightarrow B$, such that $\rho(G)$ satisfies the relations R inside B . If (G, R) is a *weakly admissible* pair of generators and relations (see Definition 1.3.4. of [9]), then one can construct the universal pro C^* -algebra $C^*(G, R)$ (see Proposition 1.3.6. of *ibid.*). It is known that any combination (even the empty set) of algebraic relations is weakly admissible (see Example 1.3.5.(1) of *ibid.*).

We further make the following hypotheses:

- (a) There are surjective maps $\theta_i : G_i \rightarrow G_{i-1}$, so that one may form the inverse limit in the category of sets $G = \varprojlim_i G_i$, with canonical projection maps $p_i : G \rightarrow G_i$. We also require the surjections θ_i to admit sections $s_{i-1} : G_{i-1} \rightarrow G_i$ satisfying $\theta_i \circ s_{i-1} = \text{id}_{G_{i-1}}$, so that we get canonical splittings $\gamma_i : G_i \rightarrow G$ satisfying $p_i \circ \gamma_i = \text{id}_{G_i}$. The map γ_i sends $g_i \rightarrow \{h_j\}$, where

$$(4) \quad h_j = \begin{cases} g_i & \text{if } j = i, \\ \theta_{i-n+1} \circ \cdots \circ \theta_i(g_i) & \text{if } j = i - n, n > 0, \\ s_{i+m-1} \circ \cdots \circ s_i(g_i) & \text{if } j = i + m, m > 0. \end{cases}$$

- (b) We require that for all i the iterated applications of θ_j 's and s_k 's on G_i satisfy R_i for all $j \leq i$ and $k \geq i$.

The surjective maps θ_i induce surjective $*$ -homomorphisms $\theta_i : C^*(G_i, R_i) \rightarrow C^*(G_{i-1}, R_{i-1})$; consequently, $\{C^*(G_i, R_i), \theta_i\}_{i \in \mathbb{N}}$ forms a countable inverse system of C^* -algebras. We may form the inverse limit $\varprojlim_i C^*(G_i, R_i)$, which is by construction a σ - C^* -algebra. Let (G, R) be a pair of generators and relations, where $G = \varprojlim_i G_i$ and R denotes the set of relations $\{\gamma_i(G_i) \text{ satisfies } R_i \text{ for all } i\}$. A *representation* ρ of (G, R) in a (pro) C^* -algebra B is a set map $\rho : G \rightarrow B$, such that $\rho \circ \gamma_i(G_i)$ satisfies R_i inside B for all i . We assume that (G, R) is a weakly admissible pair, so that one can construct the universal pro C^* -algebra $C^*(G, R)$.

Theorem. *There is an isomorphism of pro C^* -algebras $C^*(G, R) \cong \varprojlim_i C^*(G_i, R_i)$.*

Proof. It suffices to show that $\varprojlim_i C^*(G_i, R_i)$ is a universal representation of (G, R) , i.e., there is a map $\iota : G \rightarrow \varprojlim_i C^*(G_i, R_i)$ such that $\iota \circ \gamma_i(G_i)$ satisfies R_i inside $\varprojlim_i C^*(G_i, R_i)$

for all i and given any representation ρ of the pair (G, R) in a pro C^* -algebra B , there is a unique continuous $*$ -homomorphism $\kappa : \varprojlim_i C^*(G_i, R_i) \rightarrow B$ making the following diagram commute:

$$\begin{array}{ccc} G = \varprojlim_i G_i & \xrightarrow{\iota} & \varprojlim_i C^*(G_i, R_i) \\ & \searrow \rho & \downarrow \kappa \\ & & B. \end{array}$$

The map $\iota : G \rightarrow \varprojlim_i C^*(G_i, R_i)$ is defined as $g \mapsto \{p_i(g)\}$, which is a representation of (G, R) due to the Hypothesis (b) above. The construction of the universal pro C^* -algebra $C^*(G, R)$ (resp. C^* -algebra $C^*(G_i, R_i)$) is defined via a certain Hausdorff completion of $F(G)$ (resp. $F(G_i)$) with respect to representations in pro C^* -algebras (resp. C^* -algebras) satisfying R (resp. R_i). The surjective maps θ_i induce $*$ -homomorphisms $\theta_i : F(G_i) \rightarrow F(G_{i-1})$, whence we may construct the $*$ -algebra $\varprojlim_i F(G_i)$ (purely algebraic inverse limit). By the above Lemma it suffices to define κ on coherent sequences of the form $\{w_i\} \in \varprojlim_i F(G_i)$, which then extends uniquely to a $*$ -homomorphism on the entire $\varprojlim_i C^*(G_i, R_i)$. Thanks to the maps $\rho \circ \gamma_i : G_i \rightarrow B$, ρ extends uniquely to a $*$ -homomorphism $\varprojlim_i F(G_i) \rightarrow B$. Now there is a unique choice for $\kappa(\{w_i\})$ forced by the compatibility requirement, i.e., $\kappa(\{w_i\}) = \rho(\{w_i\})$. By construction κ is a $*$ -homomorphism and it is automatically continuous, since $\varprojlim_i C^*(G_i, R_i)$ is a σ - C^* -algebra. \square

In the example of $C(SU_q(\infty))$, one could try to define the section maps $s_{n-1} : G_{n-1} \rightarrow G_n$ as

$$0 \mapsto 0, \quad 1 \mapsto 1, \quad u_{ij}^{n-1} \mapsto u_{ij}^n.$$

But the Hypothesis (b) will not be satisfied and hence the above Theorem is unfortunately not applicable. However, the Theorem could be of independent interest as it can be applied to inverse systems, where the structure $*$ -homomorphisms admit sections (also $*$ -homomorphisms).

Let $G_n := \{w_{ij}^n : i, j = 1, \dots, n\} \cup \{0, 1\}$ be a set of generators satisfying the relations R_n

$$0^* = 0^2 = 0, \quad 1^* = 1^2 = 1, \quad 01 = 0 = 10, \quad 1w_{ij}^n = w_{ij}^n 1 = w_{ij}^n, \quad 0w_{ij}^n = w_{ij}^n 0 = 0, \quad \|w_{ij}^n\| \leq 1$$

for all i, j . The pair (G_n, R_n) is an admissible pair for all n , so that there is a universal C^* -algebra $C^*(G_n, R_n)$. There are surjective maps $\theta_n : G_n \rightarrow G_{n-1}$ given by

$$\begin{aligned} \theta_n(x) &:= x \text{ if } x = 0, 1 \\ \theta_n(w_{ij}^n) &:= \begin{cases} w_{ij}^{n-1} & \text{if } 1 \leq i, j \leq n-1, \\ \delta_{ij} 1 & \text{otherwise.} \end{cases} \end{aligned}$$

making $\{C^*(G_n, R_n), \theta_n\}$ an inverse system of C^* -algebras and surjective $*$ -homomorphisms. There are obvious sections $s_{n-1} : G_{n-1} \rightarrow G_n$ sending $0 \mapsto 0$, $1 \mapsto 1$ and $w_{ij}^{n-1} \mapsto w_{ij}^n$ giving rise to maps $\gamma_n : G_n \rightarrow G = \varprojlim_n G_n$ as described above (see Equation (4)). There are surjective $*$ -homomorphisms $\pi_n : C^*(G_n, R_n) \rightarrow C(SU_q(n))$ for all $n \geq 2$ given on the generators by $\pi_n(x) = x$ for $x = 0, 1$ and $\pi_n(w_{ij}^n) = u_{ij}^n$, which produce a morphism of inverse systems $\{\pi_n\} : \{C^*(G_n, R_n)\} \rightarrow \{C(SU_q(n))\}$. Indeed, it follows from the Relations (1), (2)

and (3) that the norms of the generators of $C(SU_q(n))$ do not exceed 1 in any representation. Consequently, there is a surjective $*$ -homomorphism of σ - C^* -algebras

$$\varprojlim_n \pi_n : \varprojlim_n C^*(G_n, R_n) \rightarrow C(SU_q(\infty)).$$

However, the authors cannot provide a good description of the kernel at the moment. Let us set $G = \varprojlim_n G_n$ and let R denote the set of relations $\{\gamma_n(G_n) \text{ satisfies } R_n \text{ for all } n\}$. Note that $\|x\| \leq 1$ viewed as a relation for a representation in a pro C^* -algebra B means that $p(x) \leq 1$ for all C^* -seminorms p on B . The family of pairs (G_n, R_n) satisfy Hypotheses (b) and the pair (G, R) is weakly admissible (see Example 1.3.5.(2) of [9]), so that the above Theorem applies, i.e., $\varprojlim_n C^*(G_n, R_n) \cong C^*(G, R)$. As a corollary, we deduce that the elements of $(\varprojlim_n \pi_n)(G)$ provide explicit generators of $C(SU_q(\infty))$.

REFERENCES

- [1] B. Blackadar. Shape theory for C^* -algebras. *Math. Scand.* 56 (1985), no. 2, 249–275.
- [2] N. Bourbaki. General Topology, Chapters 1-4. Elements of Mathematics, Springer-Verlag 1998.
- [3] A. L. Carey and J. Mickelsson. The universal gerbe, Dixmier-Douady class, and gauge theory. *Lett. Math. Phys.* 59 (2002), no. 1, 47–60.
- [4] J. A. Harvey and G. Moore. Noncommutative tachyons and K-theory. Strings, branes, and M-theory. *J. Math. Phys.* 42 (2001), no. 7, 2765–2780.
- [5] J. Kustermans and S. Vaes. The operator algebra approach to quantum groups. *Proc. Natl. Acad. Sci. USA* 97 (2000), no. 2, 547–552.
- [6] J. Kustermans, S. Vaes, L. Vainerman, A. Van Daele and S. Woronowicz. Lecture Notes School/Conference on Noncommutative Geometry and Quantum Groups, Warsaw 2001. <https://perswww.kuleuven.be/u0018768/artikels/lecture-notes.pdf>
- [7] A. Mallios. Topological algebras. Selected topics. North-Holland Mathematics Studies, 124, 1984.
- [8] N. C. Phillips. Inverse limits of C^* -algebras. *J. Operator Theory* 19 (1988), no. 1, 159–195.
- [9] N. C. Phillips. Inverse limits of C^* -algebras and applications. Operator algebras and applications, Vol. 1, 127–185, London. Math. Soc. Lecture Note Ser., 135, 1988.
- [10] Ya. S. Soibelman. Algebra of functions on a compact quantum group and its representations. *Algebra i Analiz*, 2(1):190–212, 1990.
- [11] S. L. Woronowicz. A remark on compact matrix quantum groups. *Lett. Math. Phys.*, 21 (1991), no. 1, 35–39.
- [12] S. L. Woronowicz. Compact quantum groups. Symétries quantiques (Les Houches, 1995), 845–884, North-Holland, Amsterdam, 1998.

E-mail address: `snigdhayan.mahanta@adelaide.edu.au`

E-mail address: `mathai.varghese@adelaide.edu.au`

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF ADELAIDE, ADELAIDE, SA 5005, AUSTRALIA