

JUCYS–MURPHY ELEMENTS FOR PARTITION ALGEBRAS

JOHN ENYANG

ABSTRACT. An inductive formula is given for a family of elements which are shown to play a role in the partition algebras which is analogous to that played by classical Jucys–Murphy elements in the group algebra of the symmetric group. Using Schur–Weyl duality it is shown that the aforementioned inductive definition is equivalent to the combinatorial definition given by Halverson and Ram for Jucys–Murphy elements of partition algebras. As a consequence of the inductive formula for Jucys–Murphy elements, a new presentation for partition algebras in terms of certain involutions is also derived.

CONTENTS

1. Introduction	1
2. The Partition Algebras	2
3. Jucys–Murphy Elements	4
4. A Presentation for the Partition Algebras	24
5. Schur–Weyl Duality	33
References	37

1. INTRODUCTION

Partition algebras arose in the work of Martin [Ma] and Jones [Jo] as a generalisation of the Temperley–Lieb algebra and the Potts model in statistical mechanics. From the point of view of Schur–Weyl duality, the partition algebras occur as centralisers for the action of the symmetric group on tensor space when the the symmetric group acts diagonally as the subgroup of permutation matrices in the complex general linear group [Jo]. Given their role as centraliser algebras arising from Schur–Weyl duality, several authors have worked to extend classical results from the representation theory of the symmetric group to the partition algebras ([Ma],[Jo],[Xi],[HR],[GG]). Using Schur–Weyl duality to prove conjectures by Owens [Ow], Halverson and Ram [HR] provided a combinatorial definition of Jucys–Murphy elements for the partition algebras which play an exactly analogous role to the classical Jucys–Murphy elements in the symmetric group. The purpose of this note is to revisit the construction of [HR], by providing a recursion for the Jucys–Murphy elements for partition algebras and establishing directly the basic properties of the Jucys–Murphy elements using only the presentation for the partition algebras given in [HR]. We also derive a new presentation for the partition algebras in terms of certain involutions that arise naturally from our recursive definition of Jucys–Murphy elements. The contents of this note are presented as follows:

- (1) In §2 we recall the presentation of the partition algebras given by [HR].
- (2) In §3 we state a recursion defining a family of operators

$$(1.1) \quad (L_i, L_{i+\frac{1}{2}} : i = 0, 1, \dots)$$

2010 *Mathematics Subject Classification.* Primary 20C08; Secondary 16G99, 20C32.

Key words and phrases. Partition algebras; Jucys–Murphy elements; central elements; presentation.

in terms of the presentation for the partition algebras given by [HR]. By direct calculation with generators and relations, we establish that the operators (1.1) form a commuting family with properties analogous to the Jucys–Murphy elements that arise in the representation theory of the symmetric group. Simultaneously, we establish some basic commutativity results for certain operators denoted

$$(1.2) \quad (\sigma_i, \sigma_{i+\frac{1}{2}} : i = 1, 2, \dots)$$

which arose in the recursive definition of the Jucys–Murphy elements (1.1).

- (3) In §4 we show that the elements of (1.2) are involutions which are related to the Coxeter generators for the symmetric group in a precise way. Using the relation between the involutions (1.2) and the Coxeter generators for the symmetric group, and the properties established in §3, we derive a new presentation for the partition algebras.
- (4) In §5 we describe precisely the action of the Jucys–Murphy elements (1.1) and the involutions (1.2) on tensor space. The action of the elements (1.1) on tensor space is then used to show that the recursive definition of Jucys–Murphy elements given in §2 is equivalent to the definition of Jucys–Murphy elements in [HR].

Given that the partition algebras are cellular [Xi] and have branching rules for induction and restriction that are described by a multiplicity-free Bratteli diagram, it is natural to expect that the partition algebras also have an integral Murphy–type basis [MS] with respect to which the Jucys–Murphy elements act triangularly. Using a framework for cellularity of algebras related to the Jones basic construction, Goodman and Graber [GG], [GG1] have demonstrated the existence of Murphy bases for the partition algebras, indexed by paths in the Bratteli diagram and compatible with restriction and induction. However, the approach of Goodman and Graber describes the action of Jucys–Murphy elements on the cell modules in terms of the action of certain central elements whose eigenvalues are not explicitly derived. In a separate note in preparation [En] we will provide an explicit construction of Murphy bases for the partition algebras, compute the action of the Jucys–Murphy elements on the cell modules, and use Nazarov’s method [Na] of computing generating functions for central elements to write down a seminormal form for the partition algebras.

Acknowledgements. The author is indebted to Arun Ram for numerous stimulating questions, comments and suggestions, and for the invitation to Melbourne where this research was completed. The author is also grateful to Fred Goodman and Susanna Fishel for several conversations during the course of this work.

2. THE PARTITION ALGEBRAS

The material of this section follows the exposition given by Halverson and Ram in [HR]. For $k = 1, 2, \dots$, let

$$A_k = \{\text{set partitions of } \{1, 2, \dots, k, 1', 2', \dots, k'\}\},$$


and

$$A_{k-\frac{1}{2}} = \{d \in A_k \mid k \text{ and } k' \text{ are in the same block of } d\}.$$

For convenience, any element $\rho \in A_k$ may be represented as a graph with k vertices in the top row, labelled from left to right, by $1, 2, \dots, k$ and k vertices in the bottom row, labelled, from left to right by $1', 2', \dots, k'$, with vertex i joined to vertex j if i and j belong to the same block of ρ . The representation of a partition by a diagram is not unique; for example the partition

$$\rho = \{\{1, 1', 3, 4', 5', 6\}, \{2, 2', 3', 4, 5, 6'\}\}$$

can be represented by the diagrams:



If $\rho_1, \rho_2 \in A_k$, then the composition $\rho_1 \circ \rho_2$ is the partition obtained by placing ρ_1 above ρ_2 and identifying each vertex in the bottom row of ρ_1 with the corresponding vertex in the top row of ρ_2 and deleting any components of the resulting diagram which contains only elements from the middle row. The composition product makes A_k into an associative monoid with identity

$$1 = \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \quad \dots \quad \begin{array}{c} \bullet \\ | \\ \bullet \end{array}$$

Let z be an indeterminate and $R = \mathbb{Z}[z]$. The partition algebra $\mathcal{A}_k(z)$ is the R -module freely generated by A_k , equipped with the product

$$\rho_1 \rho_2 = z^\ell \rho_1 \circ \rho_2, \quad \text{for } \rho_1, \rho_2 \in A_k,$$

where ℓ is the number of blocks removed from the middle row in constructing the composition $\rho_1 \circ \rho_2$. Let $\mathcal{A}_{k-\frac{1}{2}}(z)$ denote the subalgebra of $\mathcal{A}_k(z)$ generated by $A_{k-\frac{1}{2}}$. A presentation for $\mathcal{A}_k(z)$ has been given by Halverson and Ram.

Theorem 2.1 (Theorem 1.11 of [HR]). *If $k = 1, 2, \dots$, then the partition algebra $\mathcal{A}_k(z)$ is the unital associative R -algebra presented by the generators*

$$p_1, p_2, \dots, p_k, p_{1+\frac{1}{2}}, p_{2+\frac{1}{2}}, \dots, p_{k-\frac{1}{2}}, s_1, s_2, \dots, s_{k-1},$$


and the relations

- (1) (Coxeter relations)
 - (i) $s_i^2 = 1$, for $i = 1, \dots, k-1$.
 - (ii) $s_i s_j = s_j s_i$, if $j \neq i+1$.
 - (iii) $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$, for $i = 1, \dots, k-2$.
- (2) (Idempotent relations)
 - (i) $p_i^2 = z p_i$, for $i = 1, \dots, k$.
 - (ii) $p_{i+\frac{1}{2}}^2 = p_{i+\frac{1}{2}}$, for $i = 1, \dots, k-1$.
 - (iii) $s_i p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}} s_i = p_{i+\frac{1}{2}}$, for $i = 1, \dots, k-1$.
 - (iv) $s_i p_i p_{i+1} = p_i p_{i+1} s_i = p_i p_{i+1}$, for $i = 1, \dots, k-1$.
- (3) (Commutation relations)
 - (i) $p_i p_j = p_j p_i$, for $i = 1, \dots, k$ and $j = 1, \dots, k$.
 - (ii) $p_{i+\frac{1}{2}} p_{j+\frac{1}{2}} = p_{j+\frac{1}{2}} p_{i+\frac{1}{2}}$, for $i = 1, \dots, k-1$ and $j = 1, \dots, k-1$.
 - (iii) $p_i p_{j+\frac{1}{2}} = p_{j+\frac{1}{2}} p_i$, for $j \neq i, i+1$.
 - (iv) $s_i p_j = p_j s_i$, for $j \neq i, i+1$.
 - (v) $s_i p_{j+\frac{1}{2}} = p_{j+\frac{1}{2}} s_i$, for $j \neq i-1, i+1$.
 - (vi) $s_i p_i s_i = p_{i+1}$, for $i = 1, \dots, k-1$.
 - (vii) $s_i p_{i-\frac{1}{2}} s_i = s_{i-1} p_{i+\frac{1}{2}} s_{i-1}$, for $i = 2, \dots, k-1$.
- (4) (Contraction relations)
 - (i) $p_{i+\frac{1}{2}} p_j p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}$, for $j = i, i+1$.
 - (ii) $p_i p_{j-\frac{1}{2}} p_i = p_i$, for $j = i, i+1$.


The above relations also imply that:

$$\begin{aligned} p_{i+\frac{1}{2}} s_{i \pm 1} p_{i+\frac{1}{2}} &= p_{i+\frac{1}{2}} p_{i \pm \frac{1}{2}}, \\ p_i s_i p_i &= p_i p_{i+1} = p_{i+1} s_i p_{i+1}, \\ p_i p_{i+\frac{1}{2}} p_{i+1} &= p_i s_i, \\ p_{i+1} p_{i+\frac{1}{2}} p_i &= p_{i+1} s_i. \end{aligned}$$

In Theorem 2.1, Halverson and Ram have made the following identifications:



and



We also let \mathfrak{S}_k denote the symmetric group on k letters which is generated by s_1, \dots, s_{k-1} . If $u \in \mathfrak{S}_k \subset \mathcal{A}_k(z)$, and $\rho \in \mathcal{A}_k(z)$, we will sometimes write $\rho^u = u\rho u^{-1}$. It will be useful to note that R -linear map $* : \mathcal{A}_k(z) \rightarrow \mathcal{A}_k(z)$ defined on generators by

$$u^* = u^{-1} \quad (\text{for } u \in \mathfrak{S}_k)$$

and

$$p_i^* = p_i \quad (\text{for } i = 1, \dots, k) \quad \text{and} \quad p_{j+\frac{1}{2}}^* = p_{j+\frac{1}{2}} \quad (\text{for } j = 1, \dots, k-1),$$

is an algebra anti-involution of $\mathcal{A}_k(z)$. Restricting the map $*$ from $\mathcal{A}_k(z)$ to $\mathcal{A}_{k-\frac{1}{2}}(z)$, gives an algebra anti involution of $\mathcal{A}_{k-\frac{1}{2}}(z)$ which we also denote by $*$.

3. JUCYS–MURPHY ELEMENTS

The purpose of this section is to give a recursion for Jucys–Murphy elements in $\mathcal{A}_k(z)$ and $\mathcal{A}_{k+\frac{1}{2}}(z)$. It will be shown in §4 that this recursion is equivalent to the combinatorial definition of Jucys–Murphy elements given [HR]. Let $(\sigma_i : i = 1, 2, \dots)$ and $(L_i : i = 0, 1, \dots)$ be given by

$$L_0 = 0, \quad L_1 = p_1, \quad \sigma_1 = 1, \quad \text{and,} \quad \sigma_2 = s_1,$$

and, for $i = 1, 2, \dots$,

$$(3.1) \quad L_{i+1} = -s_i L_i p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}} L_i s_i + p_{i+\frac{1}{2}} L_i p_{i+1} p_{i+\frac{1}{2}} + s_i L_i s_i + \sigma_{i+1},$$

where, for $i = 2, 3, \dots$,

$$(3.2) \quad \begin{aligned} \sigma_{i+1} &= s_{i-1} s_i \sigma_i s_i s_{i-1} + s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i + p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} \\ &\quad - s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} - p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i. \end{aligned}$$

Define $(\sigma_{i+\frac{1}{2}} : i = 1, 2, \dots)$ and $(L_{i+\frac{1}{2}} : i = 0, 1, \dots)$ by

$$L_{\frac{1}{2}} = 0, \quad \sigma_{\frac{1}{2}} = 1, \quad \text{and,} \quad \sigma_{1+\frac{1}{2}} = 1,$$

and, for $i = 1, 2, \dots$,

$$(3.3) \quad L_{i+\frac{1}{2}} = -L_i p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}} L_i + p_{i+\frac{1}{2}} L_i p_i p_{i+\frac{1}{2}} + s_i L_{i-\frac{1}{2}} s_i + \sigma_{i+\frac{1}{2}},$$

where, for $i = 2, 3, \dots$,

$$(3.4) \quad \begin{aligned} \sigma_{i+\frac{1}{2}} &= s_{i-1}s_i\sigma_{i-\frac{1}{2}}s_is_{i-1} + p_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}s_i + s_ip_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}} \\ &\quad - p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} - s_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_i. \end{aligned}$$

Rewriting the last summand in (3.4) as

$$\begin{aligned} s_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_i &= s_{i-1}p_{i+\frac{1}{2}}s_{i-1}s_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}s_is_{i-1}p_{i+\frac{1}{2}}s_{i-1} \\ &= s_{i-1}p_{i+\frac{1}{2}}s_{i-1}s_ip_{i+1}p_{i+\frac{1}{2}}s_{i-1}s_iL_{i-1}s_{i-1}p_{i+\frac{1}{2}}s_{i-1} \\ &= s_{i-1}p_{i+\frac{1}{2}}s_{i-1}p_{i+1}s_{i-1}s_ip_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}s_{i-1} \\ &= s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}s_{i-1}, \end{aligned}$$

the expression (3.4) becomes

$$(3.5) \quad \begin{aligned} \sigma_{i+\frac{1}{2}} &= s_{i-1}s_i\sigma_{i-\frac{1}{2}}s_is_{i-1} + p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}s_{i-1} + s_{i-1}p_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}} \\ &\quad - p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} - s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}s_{i-1}. \end{aligned}$$

Using induction, it follows that if $i = 0, 1, \dots$, then $\sigma_{i+\frac{1}{2}} \in \mathcal{A}_{i+\frac{1}{2}}(z)$, and $L_{i+\frac{1}{2}} \in \mathcal{A}_{i+\frac{1}{2}}(z)$. Observe that if $i = 0, 1, \dots$, then $(L_i)^* = L_i$ and $(\sigma_{i+1})^* = \sigma_{i+1}$. The fact that $(L_{i+\frac{1}{2}})^* = L_{i+\frac{1}{2}}$ and $(\sigma_{i+\frac{1}{2}})^* = \sigma_{i+\frac{1}{2}}$ will be shown in Proposition 3.2.

Example 3.1. In terms of the diagram presentation for $\mathcal{A}_3(z)$, we have expressions for L_2 and σ_3 as:

$$L_2 = - \begin{array}{c} \bullet \\ | \\ \bullet \end{array} - \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} + \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} + \begin{array}{c} \bullet \\ | \\ \bullet \end{array} + \begin{array}{c} \bullet \\ | \\ \bullet \end{array} + \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} + \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array}$$

and

$$\sigma_3 = \begin{array}{c} \bullet \\ | \\ \bullet \end{array} + \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} + \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} - \begin{array}{c} \bullet \\ | \\ \bullet \end{array} - \begin{array}{c} \bullet \\ | \\ \bullet \end{array} - \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array}.$$

Using the above expressions for L_2 and σ_3 , we can write L_3 as

$$L_3 = \begin{array}{ccccccccccccc} & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & + & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} \\ & + & & + & & - & & + & & - & & + & \\ & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & + & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} \\ & - & & - & & + & & + & & + & & + & \\ & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & + & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & + & \begin{array}{c} \bullet \\ | \\ \bullet \end{array} & + & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & + & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} \\ & - & & - & & + & & + & & + & & + & \\ & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & + & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} & - & \begin{array}{c} \bullet \\ | \\ \bullet \\ | \\ \bullet \end{array} \end{array}$$

The relations given in the next proposition are fundamental for subsequent calculations.

Proposition 3.1. For $i = 1, 2, \dots$, the following statements hold:

- (1) $\sigma_{i+1}p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}$,
- (2) $s_{i+1}\sigma_{i+1}p_{i+\frac{3}{2}} = p_{i+\frac{1}{2}}s_{i+1}\sigma_{i+1}$,
- (3) $\sigma_{i+1}p_ip_{i+\frac{1}{2}} = s_iL_ip_{i+\frac{1}{2}}$,
- (4) $\sigma_{i+1}p_{i+1}p_{i+\frac{1}{2}} = L_ip_{i+\frac{1}{2}}$,

$$(5) \quad p_{i+\frac{1}{2}} L_i p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}.$$

Proof. (1) We consider each of the terms on the right hand side of the definition

$$\begin{aligned} \sigma_{i+1} p_{i+\frac{1}{2}} &= s_{i-1} s_i \sigma_i s_i s_{i-1} p_{i+\frac{1}{2}} + s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i+\frac{1}{2}} + p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \\ &\quad - s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} - p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+\frac{1}{2}}, \end{aligned}$$

beginning with the argument by induction:

$$s_{i-1} s_i \sigma_i s_i s_{i-1} p_{i+\frac{1}{2}} = s_{i-1} s_i \sigma_i p_{i-\frac{1}{2}} s_i s_{i-1} = s_{i-1} s_i p_{i-\frac{1}{2}} s_i s_{i-1} = p_{i+\frac{1}{2}}.$$

From the fact that $p_{i+\frac{1}{2}}$ commutes with L_{i-1} , $s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i+\frac{1}{2}} = p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}}$ and $p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} = p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}}$, while

$$s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} = s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}}$$

and, from the relation $p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} = p_{i-\frac{1}{2}} p_i s_{i-1} s_i p_{i-\frac{1}{2}} s_i$, we obtain

$$\begin{aligned} p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+\frac{1}{2}} &= p_{i-\frac{1}{2}} p_i s_{i-1} s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+\frac{1}{2}} \\ &= p_{i-\frac{1}{2}} s_{i-1} p_{i-1} s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+\frac{1}{2}} \\ &= p_{i-\frac{1}{2}} p_{i-1} s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+\frac{1}{2}} \\ &= p_{i-\frac{1}{2}} p_{i-1} p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \\ &= p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}}. \end{aligned}$$

Substituting the terms obtained above into the definition of $\sigma_{i+1} p_{i+\frac{1}{2}}$, we observe that all terms vanish except for $p_{i+\frac{1}{2}}$, which completes the proof of (1).

(2) The definition (3.2) gives

$$\begin{aligned} s_{i+1} \sigma_{i+1} p_{i+\frac{3}{2}} &= s_{i+1} s_{i-1} s_i \sigma_i s_i s_{i-1} p_{i+\frac{3}{2}} + s_{i+1} s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i+\frac{3}{2}} \\ &\quad + s_{i+1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_{i+\frac{3}{2}} - s_{i+1} s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i+\frac{3}{2}} \\ &\quad - s_{i+1} p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+\frac{3}{2}}. \end{aligned}$$

Now consider each of the terms on the right hand side of the above equality. Firstly,

$$\begin{aligned} s_{i+1} s_{i-1} s_i \sigma_i s_i s_{i-1} p_{i+\frac{3}{2}} &= s_{i+1} s_{i-1} s_i s_{i+1} \sigma_i s_{i+1} s_i s_{i-1} p_{i+\frac{3}{2}} \\ &= s_{i-1} s_{i+1} s_i s_{i+1} \sigma_i s_{i+1} s_i p_{i+\frac{3}{2}} s_{i-1} \\ &= s_{i-1} s_i s_{i+1} s_i \sigma_i p_{i+\frac{1}{2}} s_{i+1} s_i s_{i-1} \\ &= s_{i-1} s_i s_{i+1} p_{i-\frac{1}{2}} s_i \sigma_i s_{i+1} s_i s_{i-1} \quad (\text{by induction}) \\ &= s_{i-1} s_i p_{i-\frac{1}{2}} s_{i+1} s_i \sigma_i s_{i+1} s_i s_{i-1} \\ &= p_{i+\frac{1}{2}} s_{i-1} s_i s_{i+1} s_i \sigma_i s_{i+1} s_i s_{i-1} \\ &= p_{i+\frac{1}{2}} s_{i-1} s_{i+1} s_i s_{i+1} \sigma_i s_{i+1} s_i s_{i-1} \\ &= p_{i+\frac{1}{2}} s_{i+1} s_i s_{i-1} s_i \sigma_i s_i s_{i-1}. \end{aligned}$$

Next, use the fact that $s_i p_{i+\frac{3}{2}} s_i$ and $p_{i-\frac{1}{2}}$ commute to observe that

$$\begin{aligned}
s_{i+1} s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i+\frac{3}{2}} &= s_{i+1} s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} (s_i p_{i+\frac{3}{2}} s_i) s_i \\
&= s_{i+1} s_i p_{i-\frac{1}{2}} L_{i-1} s_i (s_i p_{i+\frac{3}{2}} s_i) p_{i-\frac{1}{2}} s_i \\
&= s_{i+1} s_i p_{i-\frac{1}{2}} L_{i-1} p_{i+\frac{3}{2}} s_i p_{i-\frac{1}{2}} s_i \\
&= s_{i+1} s_i p_{i+\frac{3}{2}} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i \\
&= p_{i+\frac{1}{2}} s_{i+1} s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i,
\end{aligned}$$

and that

$$\begin{aligned}
s_{i+1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_{i+\frac{3}{2}} &= s_{i+1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i+\frac{3}{2}} p_{i-\frac{1}{2}} \\
&= s_{i+1} p_{i-\frac{1}{2}} L_{i-1} (s_i p_{i+\frac{3}{2}} s_i) s_i p_{i-\frac{1}{2}} \\
&= s_{i+1} p_{i-\frac{1}{2}} (s_i p_{i+\frac{3}{2}} s_i) L_{i-1} s_i p_{i-\frac{1}{2}} \\
&= s_{i+1} (s_i p_{i+\frac{3}{2}} s_i) p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} \\
&= p_{i+\frac{1}{2}} s_{i+1} s_i s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} \\
&= p_{i+\frac{1}{2}} s_{i+1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}}.
\end{aligned}$$

Since $p_{i+\frac{3}{2}}$ commutes with $\mathcal{A}_{i+\frac{1}{2}}(z)$, we see that

$$\begin{aligned}
s_{i+1} s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i+\frac{3}{2}} &= s_{i+1} s_i p_{i+\frac{3}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} \\
&= p_{i+\frac{1}{2}} s_{i+1} s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}}.
\end{aligned}$$

For the last term, we use the fact that $(s_i p_{i+\frac{3}{2}} s_i)$ and $p_{i-\frac{1}{2}}$ commute to see that

$$\begin{aligned}
s_{i+1} p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+\frac{3}{2}} &= s_{i+1} p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} (s_i p_{i+\frac{3}{2}} s_i) s_i \\
&= s_{i+1} p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} (s_i p_{i+\frac{3}{2}} s_i) p_{i-\frac{1}{2}} s_i \\
&= s_{i+1} p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} (s_i p_{i+\frac{3}{2}} s_i) L_{i-1} p_{i-\frac{1}{2}} s_i \\
&= s_{i+1} p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} (s_{i-1} s_i p_{i+\frac{3}{2}} s_i s_{i-1}) L_{i-1} p_{i-\frac{1}{2}} s_i \\
&= s_{i+1} p_{i-\frac{1}{2}} (s_{i-1} s_i p_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{3}{2}} s_i s_{i-1}) L_{i-1} p_{i-\frac{1}{2}} s_i \\
&= s_{i+1} p_{i-\frac{1}{2}} (s_{i-1} s_i p_{i+\frac{3}{2}} s_i s_{i-1}) p_i p_{i+\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} s_i \\
&= s_{i+1} p_{i-\frac{1}{2}} s_i p_{i+\frac{3}{2}} s_i s_{i-1} p_i p_{i+\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} s_{i+1} s_i p_{i+\frac{3}{2}} s_i s_{i-1} p_i p_{i+\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} s_{i+1} s_i^2 s_{i-1} p_i p_{i+\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} s_i \\
&= p_{i+\frac{1}{2}} s_{i+1} p_{i-\frac{1}{2}} s_{i-1} p_i p_{i+\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} s_i.
\end{aligned}$$

Putting the above together, we have

$$\begin{aligned}
s_{i+1} \sigma_{i+1} p_{i+\frac{3}{2}} &= p_{i+\frac{1}{2}} s_{i+1} s_{i-1} s_i \sigma_i s_i s_{i-1} + p_{i+\frac{1}{2}} s_{i+1} s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i \\
&\quad + p_{i+\frac{1}{2}} s_{i+1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} - p_{i+\frac{1}{2}} s_{i+1} s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} \\
&\quad - p_{i+\frac{1}{2}} s_{i+1} p_{i-\frac{1}{2}} s_{i-1} p_i p_{i+\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} s_i,
\end{aligned}$$

which completes the proof of (2).

(3) We consider the terms on the right hand side of the equality

$$\begin{aligned}\sigma_{i+1}p_ip_{i+\frac{1}{2}} &= s_{i-1}s_i\sigma_is_i s_{i-1}p_ip_{i+\frac{1}{2}} + s_ip_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}s_ip_ip_{i+\frac{1}{2}} \\ &\quad + p_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}} - s_ip_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}} \\ &\quad - p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_ip_ip_{i+\frac{1}{2}},\end{aligned}$$

beginning with

$$\begin{aligned}s_{i-1}s_i\sigma_is_i s_{i-1}p_ip_{i+\frac{1}{2}} &= s_{i-1}s_i\sigma_ip_{i-1}s_is_{i-1}p_{i+\frac{1}{2}} \\ &= s_{i-1}s_i\sigma_ip_{i-1}p_{i+\frac{1}{2}}s_is_{i-1} \\ &= s_{i-1}s_is_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_is_{i-1} \\ &= s_is_{i-1}s_iL_{i-1}s_is_{i-1}p_{i+\frac{1}{2}} \\ &= s_is_{i-1}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}.\end{aligned}$$

For the second term, we have

$$\begin{aligned}s_ip_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}s_ip_ip_{i+\frac{1}{2}} &= s_ip_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}p_{i+1}s_ip_{i+\frac{1}{2}} \\ &= s_ip_{i-\frac{1}{2}}L_{i-1}p_is_ip_{i-\frac{1}{2}}s_ip_{i+\frac{1}{2}} \\ &= s_ip_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}}p_{i+\frac{1}{2}},\end{aligned}$$

and for the third,

$$\begin{aligned}p_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}p_{i+1}p_{i+\frac{1}{2}} &= p_{i-\frac{1}{2}}L_{i-1}p_is_ip_{i-\frac{1}{2}}p_{i+\frac{1}{2}} \\ &= p_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}}p_{i+\frac{1}{2}}.\end{aligned}$$

Using the relation $p_ip_{i-\frac{1}{2}}p_i = p_i$, we see that the fourth term satisfies

$$s_ip_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}} = s_ip_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}},$$

while, for the fifth term,

$$\begin{aligned}p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_ip_ip_{i+\frac{1}{2}} &= p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}p_{i+1}s_ip_{i+\frac{1}{2}} \\ &= p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}p_{i+1}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_ip_{i+\frac{1}{2}} \\ &= p_{i-\frac{1}{2}}s_ip_{i+1}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}} \\ &= p_{i-\frac{1}{2}}s_is_{i-1}L_{i-1}p_{i-\frac{1}{2}}p_{i+1}p_{i+\frac{1}{2}} \\ &= s_is_{i-1}p_{i+\frac{1}{2}}L_{i-1}p_{i-\frac{1}{2}}p_{i+1}p_{i+\frac{1}{2}} \\ &= s_is_{i-1}L_{i-1}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}}.\end{aligned}$$

Combining the above terms, we obtain

$$\begin{aligned}\sigma_{i+1}p_ip_{i+\frac{1}{2}} &= s_is_{i-1}L_{i-1}s_{i-1}p_{i+\frac{1}{2}} + s_ip_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}}p_{i+\frac{1}{2}} + p_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}}p_{i+\frac{1}{2}} \\ &\quad - s_ip_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}} - s_is_{i-1}L_{i-1}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}},\end{aligned}$$

which completes the proof (3).

(4) We consider each of the terms on the right hand side of the expression

$$\begin{aligned}\sigma_{i+1}p_{i+1}p_{i+\frac{1}{2}} &= s_{i-1}s_i\sigma_is_i s_{i-1}p_{i+1}p_{i+\frac{1}{2}} + s_ip_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}s_ip_{i+1}p_{i+\frac{1}{2}} \\ &\quad + p_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}p_{i+1}p_{i+\frac{1}{2}} - s_ip_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_{i+1}p_{i+\frac{1}{2}} \\ &\quad - p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_ip_{i+1}p_{i+\frac{1}{2}}.\end{aligned}$$

Firstly,

$$\begin{aligned}
s_{i-1}s_i\sigma_i s_i s_{i-1} p_{i+1} p_{i+\frac{1}{2}} &= s_{i-1}s_i\sigma_i p_i s_i s_{i-1} p_{i+\frac{1}{2}} \\
&= s_{i-1}s_i\sigma_i p_i p_{i-\frac{1}{2}} s_i s_{i-1} \\
&= s_{i-1}s_i L_{i-1} p_{i-\frac{1}{2}} s_i s_{i-1} \quad (\text{by induction}) \\
&= s_{i-1}s_i L_{i-1} s_i s_{i-1} p_{i+\frac{1}{2}} \\
&= s_{i-1} L_{i-1} s_{i-1} p_{i+\frac{1}{2}}.
\end{aligned}$$

For the second term,

$$\begin{aligned}
s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+1} p_{i+\frac{1}{2}} &= s_i p_{i-\frac{1}{2}} s_i \sigma_i p_i p_{i-\frac{1}{2}} s_i p_{i+1} p_{i+\frac{1}{2}} \quad (\text{by induction}) \\
&= s_i s_i \sigma_i p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} s_i p_{i+1} p_{i+\frac{1}{2}} \quad (\text{by item (2)}) \\
&= \sigma_i p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_i s_i p_{i+\frac{1}{2}} \\
&= \sigma_i p_{i+\frac{1}{2}} p_i s_i p_{i+\frac{1}{2}} \\
&= \sigma_i p_{i+\frac{1}{2}}.
\end{aligned}$$

For the third term,

$$\begin{aligned}
p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} &= p_{i-\frac{1}{2}} L_{i-1} p_i s_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}}.
\end{aligned}$$

For the fourth term, we use the relation $s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} = s_i p_{i-\frac{1}{2}} p_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}}$ in

$$\begin{aligned}
s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} &= s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \\
&= s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} p_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \\
&= s_i p_{i-\frac{1}{2}} s_i \sigma_i p_i p_{i-\frac{1}{2}} p_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \quad (\text{by induction}) \\
&= s_i p_{i-\frac{1}{2}} s_i \sigma_i p_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \\
&= s_i s_i \sigma_i p_{i+\frac{1}{2}} p_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \quad (\text{by item (2)}) \\
&= \sigma_i p_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \\
&= s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \quad (\text{by item (3)}).
\end{aligned}$$

For the final term, we use the relation $p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} = p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{1}{2}} s_i$ in

$$\begin{aligned}
p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+1} p_{i+\frac{1}{2}} &= p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+1} p_{i+\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{1}{2}} s_i \sigma_i p_i p_{i-\frac{1}{2}} s_i p_{i+1} p_{i+\frac{1}{2}} \quad (\text{by induction}) \\
&= p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{1}{2}} s_i \sigma_i p_i p_{i-\frac{1}{2}} p_i s_i p_{i+\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{1}{2}} s_i \sigma_i p_i p_{i+\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} s_i p_{i-1} s_i \sigma_i p_{i+\frac{1}{2}} p_i p_{i+\frac{1}{2}} \quad (\text{by item (2)}) \\
&= p_{i-\frac{1}{2}} s_i p_{i-1} s_i \sigma_i p_{i+\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} p_{i-1} \sigma_i p_{i+\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} \quad (\text{by item (3)}).
\end{aligned}$$

Putting the above together,

$$\begin{aligned}\sigma_{i+1}p_{i+1}p_{i+\frac{1}{2}} &= s_{i-1}L_{i-1}s_{i-1}p_{i+\frac{1}{2}} + \sigma_i p_{i+\frac{1}{2}} + p_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}}p_{i+\frac{1}{2}} \\ &\quad - s_{i-1}L_{i-1}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}} - p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}},\end{aligned}$$

which proves (4).

(5) Parts (1) and (3) give

$$p_{i+\frac{1}{2}}L_ip_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}s_iL_ip_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}\sigma_{i+1}p_ip_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}p_ip_{i+\frac{1}{2}} = p_{i+\frac{1}{2}},$$

as required. \square

Proposition 3.2. *If $i = 1, 2, \dots$, then*

- (1) $(\sigma_{i+\frac{1}{2}})^* = \sigma_{i+\frac{1}{2}}$,
- (2) $(L_{i+\frac{1}{2}})^* = L_{i+\frac{1}{2}}$.

Proof. (1) We show that the summand $p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}$ in (3.4) is fixed under the $*$ anti-involution on $\mathcal{A}_{i+\frac{1}{2}}(z)$, using Proposition 3.1, as follows:

$$\begin{aligned}p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} &= p_{i-\frac{1}{2}}p_i\sigma_is_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}p_i\sigma_is_ip_{i-\frac{1}{2}}s_is_{i-1}p_ip_{i-\frac{1}{2}} \\ &= p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}\sigma_is_{i-1}p_ip_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}\sigma_ip_{i-1}p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}.\end{aligned}$$

(2) Given that $(\sigma_{i+\frac{1}{2}})^* = \sigma_{i+\frac{1}{2}}$, it suffices to show that the summand $p_{i+\frac{1}{2}}L_ip_ip_{i+\frac{1}{2}}$ in (3.3) is fixed under the $*$ anti-involution on $\mathcal{A}_{i+\frac{1}{2}}(z)$. Using Proposition 3.1,

$$p_{i+\frac{1}{2}}L_ip_ip_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}p_{i+1}\sigma_{i+1}p_ip_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}p_{i+1}s_iL_ip_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}s_ip_iL_ip_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}p_iL_ip_{i+\frac{1}{2}},$$

gives the required result. \square

Proposition 3.3. *If $i = 1, 2, \dots$, then $\sigma_{i+\frac{1}{2}}s_i = s_i\sigma_{i+\frac{1}{2}} = \sigma_{i+1}$.*

Proof. After noting that σ_2 commutes with s_1 and that $\sigma_{1+\frac{1}{2}} = \sigma_2s_1$, the statement follows from induction and the fact that $p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}$ which was demonstrated in the proof of Proposition 3.2. \square

The following observation is made for later reference.

Lemma 3.4. *If $i = 1, 2, \dots$, then*

$$s_is_{i+1}\sigma_{i+1}s_{i+1}s_ip_{i+\frac{1}{2}} = s_{i+1}p_{i+\frac{1}{2}}L_is_ip_{i+\frac{3}{2}}p_{i+1}p_{i+\frac{1}{2}} = \sigma_{i+\frac{1}{2}}s_{i+1}p_{i+\frac{1}{2}}.$$

Proof. On the one hand,

$$\begin{aligned}s_is_{i+1}\sigma_{i+1}s_{i+1}s_ip_{i+\frac{1}{2}} &= s_is_{i+1}\sigma_{i+1}s_{i+1}p_{i+\frac{1}{2}} = s_is_{i+1}p_{i+\frac{3}{2}}\sigma_{i+1}s_{i+1} \\ &= s_ip_{i+\frac{3}{2}}\sigma_{i+1}s_{i+1} = s_i\sigma_{i+1}s_{i+1}p_{i+\frac{1}{2}} = \sigma_{i+\frac{1}{2}}s_{i+1}p_{i+\frac{1}{2}},\end{aligned}$$

and on the other,

$$\begin{aligned}s_{i+1}p_{i+\frac{1}{2}}L_is_ip_{i+\frac{3}{2}}p_{i+1}p_{i+\frac{1}{2}} &= s_{i+1}p_{i+\frac{1}{2}}p_{i+1}\sigma_{i+1}s_ip_{i+\frac{3}{2}}p_{i+1}p_{i+\frac{1}{2}} \\ &= s_{i+1}p_{i+\frac{1}{2}}p_{i+1}\sigma_{i+\frac{1}{2}}p_{i+\frac{3}{2}}p_{i+1}p_{i+\frac{1}{2}} = s_{i+1}p_{i+\frac{1}{2}}p_{i+1}p_{i+\frac{3}{2}}\sigma_{i+\frac{1}{2}}p_{i+1}p_{i+\frac{1}{2}} \\ &= s_ip_{i+\frac{3}{2}}s_is_{i+1}p_{i+1}p_{i+\frac{3}{2}}\sigma_{i+\frac{1}{2}}p_{i+1}p_{i+\frac{1}{2}} = s_ip_{i+\frac{3}{2}}s_ip_{i+2}p_{i+\frac{3}{2}}\sigma_{i+\frac{1}{2}}p_{i+1}p_{i+\frac{1}{2}} \\ &= s_ip_{i+\frac{3}{2}}s_i\sigma_{i+\frac{1}{2}}p_{i+2}p_{i+\frac{3}{2}}p_{i+1}p_{i+\frac{1}{2}} = s_ip_{i+\frac{3}{2}}\sigma_{i+1}p_{i+2}p_{i+\frac{3}{2}}p_{i+1}p_{i+\frac{1}{2}} \\ &= s_ip_{i+\frac{3}{2}}\sigma_{i+1}s_{i+1}p_{i+1}p_{i+\frac{1}{2}} = s_i\sigma_{i+1}s_{i+1}p_{i+\frac{1}{2}}p_{i+1}p_{i+\frac{1}{2}} \\ &= s_i\sigma_{i+1}s_{i+1}p_{i+\frac{1}{2}} = \sigma_{i+\frac{1}{2}}s_{i+1}p_{i+\frac{1}{2}},\end{aligned}$$

as required. \square

We are now in a position to state the first of the main commutativity results of this note.

Theorem 3.5. *The elements σ_{i+1} and $\sigma_{i+\frac{1}{2}}$ satisfy the following commutativity relations:*

- (1) $\sigma_{i+1}p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}\sigma_{i+1} = p_{i-\frac{1}{2}}L_{i-1}s_i p_{i-\frac{1}{2}}$ for $i = 2, 3, \dots$
- (2) $\sigma_{i+1}p_{i-1} = p_{i-1}\sigma_{i+1} = s_{i-1}\sigma_i p_{i+1}s_i s_{i-1}$ for $i = 2, 3, \dots$
- (3) $\sigma_{i+1}p_{i-\frac{3}{2}} = p_{i-\frac{3}{2}}\sigma_{i+1}$ for $i = 3, 4, \dots$
- (4) $\sigma_{i+1}s_{i-2} = s_{i-2}\sigma_{i+1}$ for $i = 3, 4, \dots$
- (5) $\sigma_{i+1}p_{i-2} = p_{i-2}\sigma_{i+1}$ for $i = 3, 4, \dots$
- (6) $\sigma_{i+\frac{1}{2}}p_{i-1} = p_{i-1}\sigma_{i+\frac{1}{2}}$ for $i = 2, 3, \dots$
- (7) $\sigma_{i+\frac{1}{2}}p_{i-\frac{3}{2}} = p_{i-\frac{3}{2}}\sigma_{i+\frac{1}{2}}$ for $i = 3, 4, \dots$
- (8) $\sigma_{i+\frac{1}{2}}s_{i-2} = s_{i-2}\sigma_{i+\frac{1}{2}}$ for $i = 3, 4, \dots$
- (9) $\sigma_{i+\frac{1}{2}}p_{i-2} = p_{i-2}\sigma_{i+\frac{1}{2}}$ for $i = 3, 4, \dots$

Proof. (1) Using Lemma 3.4,

$$\begin{aligned}
\sigma_{i+1}p_{i-\frac{1}{2}} &= -s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} - p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i-\frac{1}{2}} \\
&\quad + p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} + s_{i-1} s_i \sigma_i s_i s_{i-1} p_{i-\frac{1}{2}} + s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i-\frac{1}{2}} \\
&= -p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i-\frac{1}{2}} + p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} + s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i-\frac{1}{2}} \\
&= -p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} + p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} + s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \\
&= -p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} p_{i+\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} + p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} + s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \\
&= -p_{i-\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} + p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} + s_i p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \\
&= -p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} + p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} + p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}} \sigma_{i+1},
\end{aligned}$$

as required.

(2) We first show that

- (i) $s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i-1} = s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i-1}$,
- (ii) $p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i-1} = p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_{i-1}$,
- (iii) $s_{i-1} s_i \sigma_i s_i s_{i-1} p_{i-1} = s_{i-1} s_i \sigma_i p_{i+1} s_i s_{i-1} = p_{i-1} s_{i-1} s_i \sigma_i s_i s_{i-1}$.

The left hand side of (i) gives

$$s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i-1} = s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i+\frac{1}{2}} s_{i-1} p_{i-1} = s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i-1},$$

which is the right hand side of (i). Using the relation $p_{i+\frac{1}{2}} s_{i-1} s_i = s_{i-1} s_i p_{i-\frac{1}{2}}$, the left hand side of (ii) gives

$$\begin{aligned}
p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i-1} &= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i-1} s_i \\
&= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i s_{i-1} s_i \\
&= p_{i-\frac{1}{2}} p_i \sigma_{i-\frac{1}{2}} p_{i+\frac{1}{2}} s_{i-1} s_i p_{i-1} \\
&= p_{i-\frac{1}{2}} p_i \sigma_{i-\frac{1}{2}} s_{i-1} s_i p_{i-\frac{1}{2}} p_{i-1} \\
&= p_{i-\frac{1}{2}} p_i \sigma_i s_i p_{i-\frac{1}{2}} p_{i-1} \\
&= p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_{i-1},
\end{aligned}$$

as required. The item (iii) follows from the relation $s_{i-1}s_ip_{i+1} = p_{i-1}s_{i-1}s_i$. Next, using the definition (3.2),

$$\begin{aligned}\sigma_{i+1}p_{i-1} &= s_{i-1}s_i\sigma_is_is_{i-1}p_{i-1} + s_ip_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}s_ip_{i-1} + p_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}} \\ &\quad - s_ip_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_{i-1} - p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_ip_{i-1} \\ &= s_{i-1}s_i\sigma_is_is_{i-1}p_{i-1},\end{aligned}$$

the right hand side of which is fixed under the $*$ anti-involution on $\mathcal{A}_{i+1}(z)$.

(3) We first show that

- (i) $s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} = s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}s_{i-2}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}$,
- (ii) $s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{3}{2}}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} = s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}$,
- (iii) $s_ip_{i-\frac{1}{2}}\sigma_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} = s_{i-1}s_ip_{i-\frac{3}{2}}L_{i-2}s_{i-1}p_{i-\frac{3}{2}}s_is_{i-1}p_{i-\frac{3}{2}}$,
- (iv) $p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}p_{i-\frac{1}{2}}s_ip_{i-\frac{3}{2}} = p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}\sigma_{i-1}p_{i-\frac{1}{2}}s_ip_{i-\frac{3}{2}}$,
- (v) $p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}s_{i-2}L_{i-2}p_{i-\frac{3}{2}}p_{i-\frac{1}{2}}s_ip_{i-\frac{3}{2}} = p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}s_{i-2}L_{i-2}s_{i-2}p_{i-\frac{1}{2}}s_ip_{i-\frac{3}{2}}$,
- (vi) $p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}}p_{i-\frac{1}{2}}s_ip_{i-\frac{3}{2}} = s_{i-1}s_is_{i-1}p_{i-\frac{3}{2}}L_{i-2}s_{i-1}p_{i-\frac{3}{2}}s_is_{i-1}p_{i-\frac{3}{2}}$,
- (vii) $p_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} = p_{i-\frac{1}{2}}s_{i-2}L_{i-2}s_{i-2}s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}$,
- (viii) $p_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} = p_{i-\frac{1}{2}}\sigma_{i-1}s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}$,
- (ix) $s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}s_{i-2}s_ip_{i-\frac{1}{2}}s_ip_{i-\frac{3}{2}} = s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}}s_ip_{i-\frac{3}{2}}$,
- (x) $s_ip_{i-\frac{1}{2}}\sigma_{i-1}s_ip_{i-\frac{1}{2}}s_ip_{i-\frac{3}{2}} = s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}s_ip_{i-\frac{1}{2}}s_ip_{i-\frac{3}{2}}$.

For the left hand side of (i),

$$\begin{aligned}s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} &= s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}s_{i-1}p_{i-\frac{3}{2}}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} \\ &= s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-\frac{1}{2}}s_{i-2}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} = s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}},\end{aligned}$$

and for the right hand side of (i),

$$\begin{aligned}s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}s_{i-2}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} &= s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}s_{i-2}s_{i-1}p_{i-\frac{3}{2}}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} \\ &= s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{1}{2}}s_{i-2}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} = s_ip_{i-\frac{1}{2}}s_{i-2}p_{i-\frac{1}{2}}L_{i-2}s_{i-2}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} \\ &= s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}.\end{aligned}$$

For the left hand side of (ii),

$$\begin{aligned}s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{3}{2}}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} &= s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{3}{2}}s_{i-1}p_{i-\frac{3}{2}}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} \\ &= s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{1}{2}}p_{i-\frac{3}{2}}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} = s_ip_{i-\frac{1}{2}}s_{i-2}p_{i-\frac{1}{2}}L_{i-2}p_{i-\frac{3}{2}}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} \\ &= s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-\frac{3}{2}}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} = s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} = p_{i-\frac{3}{2}}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}},\end{aligned}$$

and for the right hand side of (ii),

$$\begin{aligned}s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} &= s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}}s_{i-1}p_{i-\frac{3}{2}}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} \\ &= s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} = s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}} \\ &= s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}}p_{i+\frac{1}{2}}p_{i-\frac{1}{2}} = s_ip_{i-\frac{3}{2}}L_{i-2}p_{i-\frac{1}{2}}p_{i-1}p_{i-\frac{3}{2}}p_{i+\frac{1}{2}} \\ &= s_ip_{i-\frac{3}{2}}L_{i-2}p_{i-\frac{1}{2}}p_{i-\frac{3}{2}}p_{i+\frac{1}{2}} = p_{i-\frac{3}{2}}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}}.\end{aligned}$$

For the left hand side of (iii),

$$\begin{aligned}
s_i p_{i-\frac{1}{2}} \sigma_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} &= s_i \sigma_{i-1} s_{i-1} p_{i-\frac{3}{2}} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} \\
&= s_i \sigma_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} = \sigma_{i-1} s_i s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} \\
&= \sigma_{i-1} p_{i-\frac{1}{2}} s_i s_{i-1} p_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} = s_{i-1} p_{i-\frac{3}{2}} s_{i-1} \sigma_{i-1} s_i p_{i-1} p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} \\
&= s_{i-1} p_{i-\frac{3}{2}} s_{i-1} \sigma_{i-1} p_{i-1} p_{i-\frac{3}{2}} s_i p_{i-\frac{1}{2}} = s_{i-1} p_{i-\frac{3}{2}} s_{i-1} L_{i-2} p_{i-\frac{3}{2}} s_i p_{i-\frac{1}{2}},
\end{aligned}$$

and for the right hand side of (iii),

$$\begin{aligned}
s_{i-1} s_i p_{i-\frac{3}{2}} L_{i-2} s_{i-1} p_{i-\frac{3}{2}} s_i s_{i-1} p_{i-\frac{3}{2}} &= s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_i s_{i-1} s_i p_{i-\frac{3}{2}} s_{i-1} p_{i-\frac{3}{2}} \\
&= s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-1} s_i p_{i-\frac{3}{2}} s_{i-1} p_{i-\frac{3}{2}} = s_{i-1} p_{i-\frac{3}{2}} s_{i-1} L_{i-2} s_i p_{i-\frac{3}{2}} s_{i-1} p_{i-\frac{3}{2}}.
\end{aligned}$$

For the left hand side of (iv),

$$\begin{aligned}
p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-2} p_{i-\frac{1}{2}} s_i p_{i-\frac{3}{2}} &= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-2} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i = p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}},
\end{aligned}$$

and, for the right hand side of (iv),

$$\begin{aligned}
p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} \sigma_{i-1} p_{i-\frac{1}{2}} s_i p_{i-\frac{3}{2}} &= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} \sigma_{i-1} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i = p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}}.
\end{aligned}$$

For the left hand side of (v),

$$p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} s_{i-2} L_{i-2} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i p_{i-\frac{3}{2}} = p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} s_{i-2} L_{i-2} p_{i-\frac{3}{2}} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i,$$

and, for the right hand side of (v),

$$\begin{aligned}
p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} s_{i-2} L_{i-2} s_{i-2} p_{i-\frac{1}{2}} s_i p_{i-\frac{3}{2}} &= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} s_{i-2} L_{i-2} s_{i-2} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} s_{i-2} L_{i-2} s_{i-2} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i.
\end{aligned}$$

For the left hand side of (vi),

$$\begin{aligned}
p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} p_{i-\frac{3}{2}} L_{i-2} p_{i-1} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i p_{i-\frac{3}{2}} &= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} p_{i-\frac{3}{2}} L_{i-2} p_{i-1} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} s_i s_i p_{i-\frac{3}{2}} L_{i-2} p_{i-1} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} s_{i-1} s_i p_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{3}{2}} p_{i-1} s_i L_{i-2} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} s_{i-1} s_i s_{i-1} p_{i-\frac{3}{2}} p_{i-1} s_i L_{i-2} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i = p_{i-\frac{1}{2}} s_i s_{i-1} p_{i-\frac{3}{2}} p_{i-1} s_i p_{i-\frac{1}{2}} L_{i-2} p_{i-\frac{3}{2}} s_i \\
&= s_i s_{i-1} p_{i-\frac{3}{2}} p_{i-1} s_i p_{i+\frac{1}{2}} s_{i-1} s_i L_{i-2} p_{i-\frac{3}{2}} s_i = s_i s_{i-1} p_{i-\frac{3}{2}} p_{i-1} s_i p_{i+\frac{1}{2}} s_{i-1} s_i L_{i-2} p_{i-\frac{3}{2}},
\end{aligned}$$

and for the right hand side of (vi),

$$\begin{aligned}
s_{i-1} s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-1} p_{i-\frac{3}{2}} s_{i-1} s_i s_{i-1} p_{i-\frac{3}{2}} &= s_{i-1} s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-1} p_{i-\frac{3}{2}} s_i s_{i-1} s_i p_{i-\frac{3}{2}} \\
&= s_{i-1} s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-1} s_i p_{i-\frac{3}{2}} s_{i-1} p_{i-\frac{3}{2}} s_i = s_{i-1} s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-1} s_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} s_i \\
&= s_{i-1} s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-1} s_i p_{i-\frac{3}{2}} s_i = s_i s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{3}{2}} L_{i-2} s_{i-1} p_{i-\frac{3}{2}}.
\end{aligned}$$

For the left hand side of (vii)

$$p_{i-\frac{1}{2}} s_{i-2} L_{i-2} p_{i-\frac{3}{2}} s_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} = p_{i-\frac{1}{2}} s_{i-2} L_{i-2} p_{i-\frac{3}{2}} s_i p_{i-\frac{1}{2}}$$

and for the right hand side of (vii)

$$p_{i-\frac{1}{2}} s_{i-2} L_{i-2} s_{i-2} s_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} = p_{i-\frac{1}{2}} s_{i-2} L_{i-2} s_{i-2} p_{i-\frac{3}{2}} s_i p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}} s_{i-2} L_{i-2} p_{i-\frac{3}{2}} s_i p_{i-\frac{1}{2}}.$$

For the left hand side of (viii),

$$\begin{aligned} p_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} &= p_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}} \\ &= p_{i-\frac{1}{2}}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}} \\ &= p_{i-\frac{3}{2}}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}}, \end{aligned}$$

and for the right hand side of (viii),

$$p_{i-\frac{1}{2}}\sigma_{i-1}s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} = p_{i-\frac{1}{2}}\sigma_{i-1}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}} = p_{i-\frac{3}{2}}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}}.$$

For the left hand side of (ix),

$$\begin{aligned} s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}s_{i-2}s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} &= s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}s_{i-2}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}}s_i \\ &= s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}}s_i \end{aligned}$$

and for the right hand side of (ix),

$$s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}}s_ip_{i-\frac{3}{2}} = s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}}s_i.$$

For the left hand side of (x),

$$s_ip_{i-\frac{1}{2}}\sigma_{i-1}s_ip_{i-\frac{1}{2}}s_ip_{i-\frac{3}{2}} = s_ip_{i-\frac{1}{2}}\sigma_{i-1}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}}s_i = p_{i-\frac{3}{2}}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}},$$

and for the right hand side of (x),

$$\begin{aligned} s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}} &= s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}}s_i \\ &= s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}}s_i = p_{i-\frac{3}{2}}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}}. \end{aligned}$$

Now, after substituting the expression

$$L_{i-1} = -p_{i-\frac{3}{2}}L_{i-2}s_{i-2} - s_{i-2}L_{i-2}p_{i-\frac{3}{2}} + p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}} - s_{i-1}L_{i-2}s_{i-1} + \sigma_{i-1}$$

into the definition (3.2), items (i)–(x) above imply that

$$\begin{aligned} \sigma_{i+1}p_{i-\frac{3}{2}} &= s_{i-1}s_is_{i-2}s_{i-1}\sigma_{i-1}s_{i-1}s_{i-2}s_is_{i-1}p_{i-\frac{3}{2}} + p_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}} \\ &\quad + s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}}s_i - s_{i-1}s_is_{i-1}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}p_{i-\frac{1}{2}}p_{i-1}p_{i-\frac{3}{2}}s_is_{i-1}p_{i-\frac{3}{2}} \\ &\quad - s_{i-1}s_ip_{i-\frac{3}{2}}p_{i-1}p_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{3}{2}}s_{i-1}s_is_{i-1}p_{i-\frac{3}{2}}. \end{aligned}$$

Since the right hand side of the above expression is fixed under the $*$ anti-involution on $\mathcal{A}_{i+1}(z)$, it follows that $\sigma_{i+1}p_{i-\frac{3}{2}} = p_{i-\frac{3}{2}}\sigma_{i+1}$.

(4) We show that after substituting the expression

$$L_{i-1} = -p_{i-\frac{3}{2}}L_{i-2}s_{i-2} - s_{i-2}L_{i-2}p_{i-\frac{3}{2}} + p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}} + s_{i-2}L_{i-2}s_{i-2} + \sigma_{i-1}$$

into the definition (3.2), conjugation by s_{i-2} permutes the summands of σ_{i+1} as follows:

- (i) $(s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}})^{s_{i-2}} = s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}$,
- (ii) $(s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{3}{2}}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}})^{s_{i-2}} = p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}p_{i-\frac{1}{2}}s_i$,
- (iii) $(s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}})^{s_{i-2}} = s_ip_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}s_ip_{i-\frac{1}{2}}s_i$,
- (iv) $(s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}s_{i-2}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}})^{s_{i-2}} = s_{i-1}s_is_{i-1}p_{i-\frac{3}{2}}L_{i-2}s_{i-2}p_{i-\frac{1}{2}}p_{i-1}p_{i-\frac{3}{2}}s_is_{i-1}$,
- (v) $(s_ip_{i-\frac{1}{2}}\sigma_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}})^{s_{i-2}} = p_{i-\frac{1}{2}}s_{i-2}L_{i-2}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}}$,
- (vi) $(p_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}})^{s_{i-2}} = p_{i-\frac{1}{2}}p_{i-\frac{3}{2}}L_{i-2}p_{i-1}p_{i-\frac{3}{2}}s_ip_{i-\frac{1}{2}}$,
- (vii) $(s_ip_{i-\frac{1}{2}}s_{i-2}L_{i-2}s_{i-2}s_ip_{i-\frac{1}{2}}s_i)^{s_{i-2}} = s_{i-1}s_is_{i-1}p_{i-\frac{3}{2}}L_{i-2}s_{i-1}p_{i-\frac{3}{2}}s_is_{i-1}$,
- (viii) $(s_ip_{i-\frac{1}{2}}\sigma_{i-1}s_ip_{i-\frac{1}{2}}s_i)^{s_{i-2}} = p_{i-\frac{1}{2}}\sigma_{i-1}s_ip_{i-\frac{1}{2}}$,
- (ix) $(s_{i-1}s_ip_{i-\frac{3}{2}}L_{i-2}s_{i-1}p_{i-\frac{3}{2}}s_is_{i-1})^{s_{i-2}} = p_{i-\frac{1}{2}}s_{i-2}L_{i-2}s_{i-2}s_ip_{i-\frac{1}{2}}$,
- (x) $(s_{i-1}s_is_{i-2}s_{i-1}\sigma_{i-1}s_{i-1}s_{i-2}s_is_{i-1})^{s_{i-2}} = s_{i-1}s_is_{i-2}s_{i-1}\sigma_{i-1}s_{i-1}s_{i-2}s_is_{i-1}$.

The item (i) follows from the relation $s_{i-2}p_{i-\frac{3}{2}} = p_{i-\frac{3}{2}}s_{i-2} = p_{i-\frac{3}{2}}$.

For the left hand side of (ii),

$$\begin{aligned}
(s_i p_{i-\frac{1}{2}} s_{i-2} L_{i-2} p_{i-\frac{3}{2}} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}})^{s_{i-2}} &= s_{i-2} s_i p_{i-\frac{1}{2}} s_{i-2} L_{i-2} p_{i-\frac{3}{2}} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} p_{i-1} p_{i-\frac{1}{2}} s_{i-2} \\
&= s_{i-2} s_i p_{i-\frac{1}{2}} s_{i-2} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} L_{i-2} p_{i-\frac{3}{2}} p_{i-1} p_{i-\frac{1}{2}} s_{i-2} \\
&= s_{i-2} s_i s_{i-2} s_{i-1} p_{i-\frac{3}{2}} p_{i+\frac{1}{2}} s_{i-1} L_{i-2} p_{i-\frac{3}{2}} p_{i-1} p_{i-\frac{1}{2}} s_{i-2} \\
&= s_i s_{i-1} p_{i-\frac{3}{2}} p_{i+\frac{1}{2}} s_{i-1} L_{i-2} p_{i-\frac{3}{2}} p_{i-1} p_{i-\frac{1}{2}} s_{i-2} \\
&= p_{i-\frac{1}{2}} s_i s_{i-1} p_{i-\frac{3}{2}} s_{i-1} L_{i-2} p_{i-\frac{3}{2}} p_{i-1} p_{i-\frac{1}{2}} s_{i-2} \\
&= p_{i-\frac{1}{2}} s_i s_{i-1} p_{i-\frac{3}{2}} s_{i-1} \sigma_{i-1} p_{i-1} p_{i-\frac{3}{2}} p_{i-1} p_{i-\frac{1}{2}} s_{i-2} \\
&= p_{i-\frac{1}{2}} s_i s_{i-1} p_{i-\frac{3}{2}} s_{i-1} \sigma_{i-1} p_{i-1} p_{i-\frac{1}{2}} s_{i-2} \\
&= p_{i-\frac{1}{2}} s_i s_{i-1} s_{i-1} \sigma_{i-1} p_{i-\frac{1}{2}} p_{i-1} p_{i-\frac{1}{2}} s_{i-2} \\
&= p_{i-\frac{1}{2}} s_i \sigma_{i-1} p_{i-\frac{1}{2}} s_{i-2},
\end{aligned}$$

and, for the right hand side of (ii),

$$\begin{aligned}
p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-2} p_{i-\frac{1}{2}} s_i &= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} p_{i-\frac{3}{2}} p_{i-2} \sigma_{i-1} p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} p_{i-1} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} p_{i-\frac{3}{2}} p_{i-2} \sigma_{i-1} p_{i-\frac{1}{2}} s_i = p_{i-\frac{1}{2}} p_{i-1} p_{i-\frac{3}{2}} p_{i-2} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \sigma_{i-1} p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} p_{i-1} p_{i-\frac{3}{2}} p_{i-2} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \sigma_{i-1} p_{i-\frac{1}{2}} s_i = p_{i-\frac{1}{2}} p_{i-1} s_{i-2} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \sigma_{i-1} p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} p_{i-1} s_{i-2} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \sigma_{i-1} p_{i-\frac{1}{2}} s_i = p_{i-\frac{1}{2}} p_{i-1} s_{i-2} s_{i-1} p_{i+\frac{1}{2}} p_{i+\frac{1}{2}} s_{i-1} \sigma_{i-1} s_i \\
&= p_{i-\frac{1}{2}} p_{i-1} p_{i-\frac{1}{2}} s_{i-2} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \sigma_{i-1} s_i = p_{i-\frac{1}{2}} s_{i-2} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \sigma_{i-1} s_i \\
&= p_{i-\frac{1}{2}} s_{i-2} s_{i-1} s_i p_{i-\frac{1}{2}} \sigma_{i-1} = p_{i-\frac{1}{2}} s_{i-2} s_i p_{i-\frac{1}{2}} \sigma_{i-1} = p_{i-\frac{1}{2}} s_i \sigma_{i-1} p_{i-\frac{1}{2}} s_{i-2},
\end{aligned}$$

where the last equality follows from $s_{i-2}p_{i-\frac{1}{2}}\sigma_{i-1} = \sigma_{i-1}p_{i-\frac{1}{2}}s_{i-2}$.

For the left hand side of (iii),

$$\begin{aligned}
(s_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} L_{i-2} p_{i-1} p_{i-\frac{3}{2}} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}})^{s_{i-2}} &= s_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} L_{i-2} p_{i-1} p_{i-\frac{3}{2}} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} s_{i-2} \\
&= s_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} L_{i-2} p_{i-1} p_{i-\frac{3}{2}} s_i p_{i-\frac{1}{2}} s_i s_{i-1} p_i p_{i-\frac{1}{2}} s_{i-2} \\
&= s_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} L_{i-2} p_{i-1} p_{i-\frac{3}{2}} s_i p_{i-\frac{1}{2}} p_{i-1} s_i p_{i-\frac{1}{2}} s_{i-2} \\
&= s_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} L_{i-2} s_i p_{i-1} p_{i-\frac{3}{2}} p_{i-\frac{1}{2}} p_{i-1} s_i p_{i-\frac{1}{2}} s_{i-2} \\
&= s_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} L_{i-2} s_i s_{i-1} p_i p_{i-\frac{3}{2}} s_{i-1} s_i p_{i-\frac{1}{2}} s_{i-2} \\
&= s_i p_{i-\frac{3}{2}} L_{i-2} p_{i-\frac{1}{2}} s_i s_{i-1} p_i p_{i-\frac{3}{2}} s_{i-1} s_i p_{i-\frac{1}{2}} s_{i-2} \\
&= s_i p_{i-\frac{3}{2}} L_{i-2} s_i s_{i-1} p_{i+\frac{1}{2}} p_i p_{i+\frac{1}{2}} p_{i-\frac{3}{2}} s_{i-1} s_i s_{i-2} \\
&= p_{i-\frac{3}{2}} L_{i-2} s_i s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{3}{2}} s_{i-1} s_i s_{i-2} \\
&= p_{i-\frac{3}{2}} L_{i-2} s_i s_{i-1} p_{i-\frac{3}{2}} s_{i-1} s_i p_{i-\frac{1}{2}} s_{i-2} \\
&= p_{i-\frac{3}{2}} p_{i-1} \sigma_{i-1} s_{i-1} p_{i-\frac{3}{2}} s_{i-1} s_i p_{i-\frac{1}{2}} s_{i-2} \\
&= p_{i-\frac{3}{2}} p_{i-1} p_{i-\frac{1}{2}} \sigma_{i-1} s_{i-1} s_{i-1} s_i p_{i-\frac{1}{2}} s_{i-2} \\
&= p_{i-\frac{3}{2}} p_{i-1} p_{i-\frac{1}{2}} \sigma_{i-1} s_i p_{i-\frac{1}{2}} s_{i-2},
\end{aligned}$$

and for the right hand side of (iii),

$$\begin{aligned}
& s_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} L_{i-2} s_{i-2} s_i p_{i-\frac{1}{2}} s_i = s_i p_{i-\frac{1}{2}} s_i p_{i-\frac{3}{2}} L_{i-2} s_{i-2} p_{i-\frac{1}{2}} s_i \\
& = s_{i-1} p_{i+\frac{1}{2}} s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-2} p_{i-\frac{1}{2}} s_i = p_{i-\frac{3}{2}} L_{i-2} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} s_{i-2} p_{i-\frac{1}{2}} s_i \\
& = p_{i-\frac{3}{2}} L_{i-2} s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{3}{2}} s_{i-1} s_{i-2} s_i = p_{i-\frac{3}{2}} p_{i-1} \sigma_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{3}{2}} s_{i-1} s_{i-2} s_i \\
& = p_{i-\frac{3}{2}} p_{i-1} \sigma_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{3}{2}} s_{i-1} s_{i-2} s_i = p_{i-\frac{3}{2}} p_{i-1} p_{i-\frac{1}{2}} \sigma_{i-1} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} s_{i-2} s_i \\
& = p_{i-\frac{3}{2}} p_{i-1} p_{i-\frac{1}{2}} \sigma_{i-1} s_i p_{i-\frac{1}{2}} s_i s_{i-2} s_i = p_{i-\frac{3}{2}} p_{i-1} p_{i-\frac{1}{2}} \sigma_{i-1} s_i p_{i-\frac{1}{2}} s_{i-2}.
\end{aligned}$$

From the left hand side of (iv), we obtain

$$\begin{aligned}
(s_i p_{i-\frac{1}{2}} s_{i-2} L_{i-2} s_{i-2} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}})^{s_{i-2}} &= (s_i p_{i-\frac{1}{2}} s_{i-2} s_{i-1} L_{i-2} s_{i-1} s_{i-2} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}})^{s_{i-2}} \\
&= (s_i s_{i-2} s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-1} s_{i-2} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}})^{s_{i-2}} \\
&= s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-1} s_{i-2} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} s_{i-2} \\
&= s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-2} s_{i-1} s_{i-2} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} s_{i-2} \\
&= s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-2} s_{i-1} p_{i+\frac{1}{2}} p_i s_{i-2} p_{i-\frac{1}{2}} s_{i-2} \\
&= s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-2} s_{i-1} p_{i+\frac{1}{2}} p_i s_{i-1} p_{i-\frac{3}{2}} s_{i-1} \\
&= s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-2} s_i p_{i-\frac{1}{2}} s_i s_{i-1} p_i s_{i-1} p_{i-\frac{3}{2}} s_{i-1} \\
&= s_i s_{i-1} s_i p_{i-\frac{3}{2}} L_{i-2} s_{i-2} p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{3}{2}} s_{i-1} \\
&= s_{i-1} s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-2} p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{3}{2}} s_{i-1} \\
&= s_{i-1} s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_{i-2} p_{i-\frac{1}{2}} p_i-1 p_{i-\frac{3}{2}} s_i s_{i-1}
\end{aligned}$$

which is identical to the right hand side of (iv).

From the left hand side of (v), we obtain

$$\begin{aligned}
(s_i p_{i-\frac{1}{2}} \sigma_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}})^{s_{i-2}} &= s_{i-2} s_i \sigma_{i-1} s_{i-1} p_{i-\frac{3}{2}} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} s_{i-2} \\
&= s_i \sigma_{i-\frac{3}{2}} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} = p_{i-\frac{1}{2}} \sigma_{i-\frac{3}{2}} s_i s_{i-1} p_i p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} \\
&= p_{i-\frac{1}{2}} \sigma_{i-\frac{3}{2}} s_i p_{i-1} p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} = p_{i-\frac{1}{2}} \sigma_{i-1} s_i p_{i-2} p_{i-\frac{1}{2}} p_{i-\frac{3}{2}} \\
&= p_{i-\frac{1}{2}} \sigma_{i-1} p_{i-2} p_{i-\frac{3}{2}} s_i p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}} s_{i-2} L_{i-2} p_{i-\frac{3}{2}} s_i p_{i-\frac{1}{2}},
\end{aligned}$$

which is identical to the right hand side of (v).

The item (vi) follows immediately from the relation $s_{i-2} p_{i-\frac{3}{2}} = p_{i-\frac{3}{2}} s_{i-2} = p_{i-\frac{3}{2}}$.

From the left hand side of (vii), we obtain

$$\begin{aligned}
(s_i p_{i-\frac{1}{2}} s_{i-2} L_{i-2} s_{i-2} s_i p_{i-\frac{1}{2}} s_i)^{s_{i-2}} &= s_{i-2} s_i s_{i-2} s_{i-1} p_{i-\frac{3}{2}} s_i-1 L_{i-2} s_{i-2} s_i p_{i-\frac{1}{2}} s_i s_{i-2} \\
&= s_i s_{i-1} p_{i-\frac{3}{2}} s_{i-1} L_{i-2} s_{i-2} s_i p_{i-\frac{1}{2}} s_i s_{i-2} = s_i s_{i-1} p_{i-\frac{3}{2}} s_i-1 L_{i-2} s_{i-2} s_i-1 p_{i+\frac{1}{2}} s_i-1 s_{i-2} \\
&= s_i s_{i-1} p_{i-\frac{3}{2}} s_{i-1} L_{i-2} s_i s_{i-1} p_{i-\frac{3}{2}} s_i-1 s_i = s_i s_{i-1} p_{i-\frac{3}{2}} s_i L_{i-2} s_i-1 s_i p_{i-\frac{3}{2}} s_i-1 s_i \\
&= s_i s_{i-1} s_i p_{i-\frac{3}{2}} L_{i-2} s_i-1 p_{i-\frac{3}{2}} s_i s_{i-1} s_i = s_{i-1} s_i s_{i-1} p_{i-\frac{3}{2}} L_{i-2} s_i-1 p_{i-\frac{3}{2}} s_i-1 s_i s_{i-1},
\end{aligned}$$

which is identical to the right hand side of (vii).

From the left hand side of (viii),

$$\begin{aligned}
(s_i p_{i-\frac{1}{2}} \sigma_{i-1} s_i p_{i-\frac{1}{2}} s_i)^{s_{i-2}} &= s_{i-2} s_i p_{i-\frac{1}{2}} \sigma_{i-1} s_i-1 p_{i+\frac{1}{2}} s_i-1 s_{i-2} \\
&= s_{i-2} s_i \sigma_{i-1} s_i-1 p_{i-\frac{3}{2}} p_{i+\frac{1}{2}} s_i-1 s_{i-2} = \sigma_{i-\frac{3}{2}} s_i s_{i-1} p_{i-\frac{3}{2}} p_{i+\frac{1}{2}} s_i-1 s_{i-2} \\
&= \sigma_{i-\frac{3}{2}} s_i s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{3}{2}} s_i-1 s_{i-2} = \sigma_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i s_{i-1} s_i-1 s_{i-2} p_{i-\frac{1}{2}} \\
&= \sigma_{i-\frac{3}{2}} p_{i-\frac{1}{2}} s_i s_{i-2} p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}} \sigma_{i-\frac{3}{2}} s_i-2 s_i p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}} \sigma_{i-1} s_i p_{i-\frac{1}{2}},
\end{aligned}$$

which is identical to the right hand side of (viii).

From the left hand side of (ix),

$$\begin{aligned} & (s_{i-1}s_ip_{i-\frac{3}{2}}L_{i-2}s_{i-1}p_{i-\frac{3}{2}}s_is_{i-1})^{s_{i-2}} = s_{i-2}s_{i-1}p_{i-\frac{3}{2}}L_{i-2}s_is_{i-1}s_ip_{i-\frac{3}{2}}s_{i-1}s_{i-2} \\ &= p_{i-\frac{1}{2}}s_{i-2}s_{i-1}L_{i-2}s_is_{i-1}s_is_{i-1}s_{i-2}p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}s_{i-2}L_{i-2}s_is_{i-1}s_is_{i-1}s_{i-2}p_{i-\frac{1}{2}} \\ &= p_{i-\frac{1}{2}}s_{i-2}L_{i-2}s_is_{i-2}p_{i-\frac{1}{2}}, \end{aligned}$$

which is identical to the right hand side of (ix).

The equality (x) follows from the fact that $s_{i-2}s_{i-1}s_is_{i-2}s_{i-1} = s_{i-1}s_is_{i-2}s_{i-1}s_i$.

(5) By (2) and (4), σ_{i+1} commutes with $\langle s_{i-2}, p_{i-1} \rangle$, and so with $p_{i-2} = s_{i-2}p_{i-1}s_{i-2}$.

(6) By Proposition 3.3, and (2),

$$\sigma_{i+\frac{1}{2}}p_{i-1} = s_i\sigma_{i+1}p_{i-1} = p_{i-1}s_i\sigma_{i+1} = p_{i-1}\sigma_{i+\frac{1}{2}}.$$

(7)–(9) Can be proved using the same argument as part (6). \square

Theorem 3.6. *The elements L_{i+1} and $L_{i+\frac{1}{2}}$ satisfy the following commutation relations:*

- (1) $L_{i+1}p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}L_{i+1} = p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}}$, for $i = 1, 2, \dots$
- (2) $L_{i+1}p_i = p_iL_{i+1}$, for $i = 1, 2, \dots$
- (3) $L_{i+1}p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}L_{i+1}$, for $i = 2, 3, \dots$
- (4) $L_{i+1}s_{i-1} = s_{i-1}L_{i+1}$, for $i = 2, 3, \dots$
- (5) $L_{i+1}p_{i-1} = p_{i-1}L_{i+1}$, for $i = 2, 3, \dots$
- (6) $L_{i+\frac{1}{2}}p_i = p_iL_{i+\frac{1}{2}}$, for $i = 1, 2, \dots$
- (7) $L_{i+\frac{1}{2}}p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}L_{i+\frac{1}{2}}$, for $i = 2, 3, \dots$
- (8) $L_{i+\frac{1}{2}}s_{i-1} = s_{i-1}L_{i+\frac{1}{2}}$, for $i = 2, 3, \dots$
- (9) $L_{i+\frac{1}{2}}p_{i-1} = p_{i-1}L_{i+\frac{1}{2}}$, for $i = 2, 3, \dots$

Proof. (1) Using Proposition 3.1,

$$\begin{aligned} L_{i+1}p_{i+\frac{1}{2}} &= -s_iL_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_is_ip_{i+\frac{1}{2}} + p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}} + s_iL_is_ip_{i+\frac{1}{2}} + \sigma_{i+1}p_{i+\frac{1}{2}} \\ &= -s_iL_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_ip_{i+\frac{1}{2}} + p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}} + s_iL_ip_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} \\ &= -s_iL_ip_{i+\frac{1}{2}} - p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}} + s_iL_ip_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} \\ &= p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}}, \end{aligned}$$

as required.

(2) From Proposition 3.1, we obtain $\sigma_{i+1}p_i = \sigma_{i+1}p_ip_{i+\frac{1}{2}}p_i = s_iL_ip_{i+\frac{1}{2}}p_i$. Thus

$$\begin{aligned} L_{i+1}p_i &= -s_iL_ip_{i+\frac{1}{2}}p_i - p_{i+\frac{1}{2}}L_is_ip_i + p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}}p_i + s_iL_is_ip_i + \sigma_{i+1}p_i \\ &= (\sigma_{i+1}p_i - s_iL_ip_{i+\frac{1}{2}}p_i) + (p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}}p_i - p_{i+\frac{1}{2}}L_is_ip_i) + s_iL_ip_{i+1}s_i \\ &= (\sigma_{i+1}p_i - \sigma_{i+1}p_i) + (p_{i+\frac{1}{2}}L_is_ip_i - p_{i+\frac{1}{2}}L_is_ip_i) + s_iL_ip_{i+1}s_i, \end{aligned}$$

and the statement now follows from the fact that the right hand side of the above expression is fixed under the * anti-involution on $\mathcal{A}_{i+1}(z)$.

(3) We first show that

- (i) $p_{i+\frac{1}{2}}L_is_ip_{i-\frac{1}{2}} = p_{i+\frac{1}{2}}\sigma_is_i$,
- (ii) $s_iL_ip_{i+\frac{1}{2}}p_{i-\frac{1}{2}} = s_ip_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}}p_{i+\frac{1}{2}}$,
- (iii) $p_{i+\frac{1}{2}}L_ip_{i+1}p_{i+\frac{1}{2}}p_{i-\frac{1}{2}} = p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}}p_{i+\frac{1}{2}}$,
- (iv) $\sigma_{i+1}p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}\sigma_{i+1}$.

(i) The definition (3.1) gives

$$\begin{aligned}
p_{i+\frac{1}{2}} L_i s_i p_{i-\frac{1}{2}} &= -p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i-\frac{1}{2}} - p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} s_i p_{i-\frac{1}{2}} \\
&\quad + p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} s_i p_{i-\frac{1}{2}} + p_{i+\frac{1}{2}} s_{i-1} L_{i-1} s_{i-1} s_i p_{i-\frac{1}{2}} + p_{i+\frac{1}{2}} \sigma_i s_i p_{i-\frac{1}{2}} \\
&= -p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_{i+\frac{1}{2}} s_{i-1} s_i \\
&\quad + p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i+\frac{1}{2}} s_{i-1} s_i + p_{i+\frac{1}{2}} \sigma_i s_i p_{i-\frac{1}{2}} \\
&= -p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} - p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} s_i \\
&\quad + p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} + p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} s_i + p_{i+\frac{1}{2}} \sigma_i s_i p_{i-\frac{1}{2}} \\
&= p_{i+\frac{1}{2}} \sigma_i s_i,
\end{aligned}$$

where the last equality follows from Proposition 3.1.

(ii) The definition (3.1) gives

$$\begin{aligned}
s_i L_i p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} &= -s_i s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} - s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} \\
&\quad + s_i p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} + s_i s_{i-1} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} + s_i \sigma_i p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} \\
&= -s_i s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} - s_i p_{i-\frac{1}{2}} L_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} \\
&\quad + s_i p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} + s_i s_{i-1} L_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} + p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} \\
&= s_i p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}}
\end{aligned}$$

since $p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}$.

(iii) The definition (3.1) gives

$$\begin{aligned}
p_{i+\frac{1}{2}} L_i p_{i+1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} &= -p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} \\
&\quad + p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} s_{i-1} L_{i-1} s_{i-1} p_{i+1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} \\
&\quad + p_{i+\frac{1}{2}} \sigma_i p_{i+1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} \\
&= -p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} \\
&\quad + p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} \\
&\quad + p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} \\
&= -p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} \\
&\quad + p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} \\
&= p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}}.
\end{aligned}$$

(iv) Was demonstrated in Theorem 3.5.

Now, using

$$\begin{aligned}
s_i L_{i-1} s_{i-1} p_{i-\frac{1}{2}} &= -s_i s_{i-1} L_{i-1} p_{i-\frac{1}{2}} - p_{i-\frac{1}{2}} L_{i-1} s_{i-1} s_i p_{i-\frac{1}{2}} + s_i p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} s_i p_{i-\frac{1}{2}} \\
&\quad + s_i s_{i-1} L_{i-1} s_{i-1} s_i p_{i-\frac{1}{2}} + s_i \sigma_i s_i p_{i-\frac{1}{2}} \\
&= -s_i s_{i-1} L_{i-1} p_{i-\frac{1}{2}} - p_{i-\frac{1}{2}} L_{i-1} s_{i-1} s_i + s_i p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \\
&\quad + s_i s_{i-1} L_{i-1} s_{i-1} s_i p_{i-\frac{1}{2}} + p_{i+\frac{1}{2}} \sigma_i s_i,
\end{aligned}$$

we obtain

$$\begin{aligned}
L_{i+1}p_{i-\frac{1}{2}} &= -s_i L_i p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} - p_{i+\frac{1}{2}} L_i s_i p_{i-\frac{1}{2}} + p_{i+\frac{1}{2}} L_i p_{i+1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} + s_i L_i s_i p_{i-\frac{1}{2}} + \sigma_{i+1} p_{i-\frac{1}{2}} \\
&= -s_i p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}} \sigma_i s_i + p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} \\
&\quad + s_i L_i s_i p_{i-\frac{1}{2}} + \sigma_{i+1} p_{i-\frac{1}{2}} \\
&= p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} + s_i s_{i-1} L_{i-1} s_{i-1} s_i p_{i-\frac{1}{2}} + \sigma_{i+1} p_{i-\frac{1}{2}}.
\end{aligned}$$

Since the right hand side of the last expression is fixed under the $*$ anti-involution on $\mathcal{A}_{i+1}(z)$, the proof of (3) is complete.

(4) We show that after substituting the expression

$$L_i = -p_{i-\frac{1}{2}} L_{i-1} s_{i-1} - s_{i-1} L_{i-1} p_{i-\frac{1}{2}} + p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} + s_{i-1} L_{i-1} s_{i-1} + \sigma_i$$

into the definition (3.1), conjugation by s_{i-1} permutes the summands of L_{i+1} as follows:

- (i) $s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} s_i s_{i-1} = p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} s_i,$
- (ii) $s_{i-1} p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i s_{i-1} = s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}},$
- (iii) $s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} s_i s_{i-1} = p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+1} p_{i+\frac{1}{2}},$
- (iv) $s_{i-1} p_{i+\frac{1}{2}} s_{i-1} L_{i-1} s_{i-1} s_i s_{i-1} = s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} s_i,$
- (v) $s_{i-1} p_{i+\frac{1}{2}} \sigma_i s_i s_{i-1} = s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}},$
- (vi) $s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} s_{i-1} = p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}},$
- (vii) $s_{i-1} p_{i+\frac{1}{2}} s_{i-1} L_{i-1} s_{i-1} p_{i+1} p_{i+\frac{1}{2}} s_{i-1} = s_i p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} s_i,$
- (viii) $s_{i-1} p_{i+\frac{1}{2}} \sigma_i p_{i+1} p_{i+\frac{1}{2}} s_{i-1} = s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i,$
- (ix) $s_{i-1} s_i s_{i-1} L_{i-1} s_{i-1} s_i s_{i-1} = s_i s_{i-1} L_{i-1} s_{i-1} s_i,$
- (x) $s_{i-1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_{i-1} = p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}}.$

For item (i),

$$s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} s_i s_{i-1} = p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_i s_{i-1} s_i = p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} s_i.$$

For item (ii),

$$\begin{aligned}
s_{i-1} p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i s_{i-1} &= s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} s_i s_{i-1} \\
&= s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i s_{i-1} = s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}}.
\end{aligned}$$

For item (iii),

$$s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i p_{i-\frac{1}{2}} s_i s_{i-1} = p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_i s_i s_{i-1} p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+1} p_{i+\frac{1}{2}}.$$

For item (iv),

$$\begin{aligned}
s_{i-1} p_{i+\frac{1}{2}} s_{i-1} L_{i-1} s_{i-1} s_i s_{i-1} &= s_i p_{i-\frac{1}{2}} s_i L_{i-1} s_{i-1} s_i s_{i-1} \\
&= s_i p_{i-\frac{1}{2}} L_{i-1} s_i s_{i-1} s_i s_{i-1} = s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} s_i.
\end{aligned}$$

For the left hand side of item (v),

$$s_{i-1} p_{i+\frac{1}{2}} \sigma_i s_i s_{i-1} = s_{i-1} \sigma_i s_i p_{i-\frac{1}{2}} s_{i-1} = \sigma_{i-\frac{1}{2}} s_i p_{i-\frac{1}{2}},$$

and for the right hand side of item (v),

$$\begin{aligned}
s_i p_{i-\frac{1}{2}} L_{i-1} s_i s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} &= s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i s_{i-1} p_i p_{i-\frac{1}{2}} = s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{1}{2}} \\
&= s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} p_{i-1} s_i p_{i-\frac{1}{2}} = s_i p_{i-\frac{1}{2}} s_i \sigma_i p_i p_{i-\frac{1}{2}} p_{i-1} s_i p_{i-\frac{1}{2}} = s_i p_{i-\frac{1}{2}} s_i \sigma_i s_{i-1} p_{i-1} s_i p_{i-\frac{1}{2}} \\
&= s_i s_i \sigma_i p_{i+\frac{1}{2}} s_{i-1} s_i p_{i-1} p_{i-\frac{1}{2}} = \sigma_i s_{i-1} s_i p_{i-\frac{1}{2}} p_{i-1} p_{i-\frac{1}{2}} = \sigma_{i-\frac{1}{2}} s_i p_{i-\frac{1}{2}}
\end{aligned}$$

The item (vi) follows from the relation $s_{i-1}p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}s_{i-1} = p_{i-\frac{1}{2}}$. For item (vii),

$$s_{i-1}p_{i+\frac{1}{2}}s_{i-1}L_{i-1}s_{i-1}p_{i+1}p_{i+\frac{1}{2}}s_{i-1} = s_i p_{i-\frac{1}{2}}s_i L_{i-1}p_{i+1}s_i p_{i-\frac{1}{2}}s_i = s_i p_{i-\frac{1}{2}}L_{i-1}p_i p_{i-\frac{1}{2}}s_i.$$

For item (viii),

$$\begin{aligned} s_{i-1}p_{i+\frac{1}{2}}\sigma_i p_{i+1}p_{i+\frac{1}{2}}s_{i-1} &= s_{i-1}\sigma_i s_i p_{i-\frac{1}{2}}s_i p_{i+1}p_{i+\frac{1}{2}}s_{i-1} = \sigma_{i-\frac{1}{2}}s_i p_{i-\frac{1}{2}}p_i p_{i+\frac{1}{2}}s_{i-1} \\ &= \sigma_{i-\frac{1}{2}}s_i p_{i-\frac{1}{2}}p_i s_{i-1}s_i p_{i-\frac{1}{2}}s_i = \sigma_{i-\frac{1}{2}}s_i p_{i-\frac{1}{2}}p_{i-1}s_i p_{i-\frac{1}{2}}s_i = \sigma_{i-\frac{1}{2}}s_i p_{i-\frac{1}{2}}s_i p_{i-1}p_{i-\frac{1}{2}}s_i \\ &= \sigma_{i-\frac{1}{2}}s_i p_{i-\frac{1}{2}}s_i p_{i-1}p_{i-\frac{1}{2}}s_i = \sigma_{i-\frac{1}{2}}s_{i-1}p_{i+\frac{1}{2}}s_{i-1}p_{i-1}p_{i-\frac{1}{2}}s_i = s_{i-1}p_{i+\frac{1}{2}}s_{i-1}\sigma_{i-\frac{1}{2}}p_{i-1}p_{i-\frac{1}{2}}s_i \\ &= s_i p_{i-\frac{1}{2}}s_i \sigma_{i-\frac{1}{2}}p_{i-1}p_{i-\frac{1}{2}}s_i = s_i p_{i-\frac{1}{2}}s_i L_{i-1}p_{i-\frac{1}{2}}s_i. \end{aligned}$$

(ix) Follows from the Coxeter relations, and (x) from the relation $s_{i-1}p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}s_{i-1} = p_{i-\frac{1}{2}}$.

(5) By parts (2) and (4), L_{i+1} commutes with $\langle p_i, s_{i-1} \rangle$, and so with $p_{i-1} = s_{i-1}p_i s_{i-1}$.

(6) From item (3) of Proposition 3.1, $\sigma_{i+\frac{1}{2}}p_i = s_i \sigma_{i+1}p_i p_{i+\frac{1}{2}}p_i = L_i p_{i+\frac{1}{2}}p_i$, and

$$\begin{aligned} L_{i+\frac{1}{2}}p_i &= -p_{i+\frac{1}{2}}L_i p_i - L_i p_{i+\frac{1}{2}}p_i + p_{i+\frac{1}{2}}L_i p_i p_{i+\frac{1}{2}}p_i + s_i L_{i-\frac{1}{2}}s_i p_i + \sigma_{i+\frac{1}{2}}p_i \\ &= -p_{i+\frac{1}{2}}L_i p_i - L_i p_{i+\frac{1}{2}}p_i + p_{i+\frac{1}{2}}L_i p_i + s_i L_{i-\frac{1}{2}}p_{i+1}s_i + L_i p_{i+\frac{1}{2}}p_i \\ &= s_i L_{i-\frac{1}{2}}p_{i+1}s_i. \end{aligned}$$

Since $s_i L_{i-\frac{1}{2}}p_{i+1}s_i = s_i p_{i+1}L_{i-\frac{1}{2}}s_i$, this completes the proof of (6).

(7) We show that

$$(3.6) \quad L_{i+\frac{1}{2}}p_{i-\frac{1}{2}} = -p_{i-\frac{1}{2}}L_{i-1}p_i p_{i-\frac{1}{2}}p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}p_i p_{i-\frac{1}{2}} \\ + s_i s_{i-1}L_{i-\frac{3}{2}}p_{i+\frac{1}{2}}s_{i-1}s_i + s_i s_{i-1}\sigma_{i-\frac{1}{2}}p_{i+\frac{1}{2}}s_{i-1}s_i.$$

From the definition (3.3),

$$(3.7) \quad L_{i+\frac{1}{2}}p_{i-\frac{1}{2}} = (-L_i p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}}L_i + p_{i+\frac{1}{2}}L_i p_i p_{i+\frac{1}{2}} + s_i L_{i-\frac{1}{2}}s_i + \sigma_{i+\frac{1}{2}})p_{i-\frac{1}{2}}.$$

Using part (1) the first three summands in the right hand side of (3.7) are transformed as:

$$L_i p_{i+\frac{1}{2}}p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}L_{i-1}p_i p_{i-\frac{1}{2}}p_{i+\frac{1}{2}} \quad \text{and} \quad p_{i+\frac{1}{2}}L_i p_{i-\frac{1}{2}} = p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}p_i p_{i-\frac{1}{2}},$$

and

$$\begin{aligned} p_{i+\frac{1}{2}}L_i p_i p_{i+\frac{1}{2}}p_{i-\frac{1}{2}} &= p_{i+\frac{1}{2}}p_i L_i p_{i-\frac{1}{2}}p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}p_i p_{i-\frac{1}{2}}L_{i-1}p_i p_{i-\frac{1}{2}}p_{i+\frac{1}{2}} \\ &= p_{i+\frac{1}{2}}p_i L_{i-1}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}} = L_{i-1}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}}. \end{aligned}$$

Next,

$$\begin{aligned} s_i L_{i-\frac{1}{2}}s_i p_{i-\frac{1}{2}} &= -s_i L_{i-1}p_{i-\frac{1}{2}}s_i p_{i-\frac{1}{2}} - s_i p_{i-\frac{1}{2}}L_{i-1}s_i p_{i-\frac{1}{2}} + s_i p_{i-\frac{1}{2}}L_{i-1}p_{i-1}p_{i-\frac{1}{2}}s_i p_{i-\frac{1}{2}} \\ &\quad + s_i s_{i-1}L_{i-\frac{3}{2}}s_{i-1}s_i p_{i-\frac{1}{2}} + s_i \sigma_{i-\frac{1}{2}}s_i p_{i-\frac{1}{2}} \\ &= -s_i L_{i-1}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}} - s_i p_{i-\frac{1}{2}}L_{i-1}s_i p_{i-\frac{1}{2}} + s_i p_{i-\frac{1}{2}}L_{i-1}p_{i-1}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}} \\ &\quad + s_i s_{i-1}L_{i-\frac{3}{2}}s_{i-1}s_i p_{i-\frac{1}{2}} + s_i s_{i-1}\sigma_{i-\frac{1}{2}}s_i p_{i-\frac{1}{2}} \\ &= -L_{i-1}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}} - s_i p_{i-\frac{1}{2}}L_{i-1}s_i p_{i-\frac{1}{2}} + p_{i-\frac{1}{2}}L_{i-1}p_{i-1}p_{i-\frac{1}{2}}p_{i+\frac{1}{2}} \\ &\quad + s_i s_{i-1}L_{i-\frac{3}{2}}p_{i+\frac{1}{2}}s_{i-1}s_i + s_i s_{i-1}\sigma_{i-\frac{1}{2}}p_{i+\frac{1}{2}}s_{i-1}s_i. \end{aligned}$$

Substituting each of the above into (3.7), and using part (1) of Theorem 3.5 gives (3.6). The right hand side of (3.6) being fixed under the * anti-involution on $\mathcal{A}_{i+\frac{1}{2}}(z)$, the proof of (7) is complete.

(8) We show that, after substituting the expression

$$L_{i-\frac{1}{2}} = -p_{i-\frac{1}{2}}L_{i-1} - L_{i-1}p_{i-\frac{1}{2}} + p_{i-\frac{1}{2}}L_{i-1}p_{i-1}p_{i-\frac{1}{2}} + s_{i-1}L_{i-\frac{1}{2}}s_{i-1} + \sigma_{i-\frac{1}{2}}$$

into the definition (3.3), conjugation by s_{i-1} permutes the summands of $L_{i+\frac{1}{2}}$ as follows:

- (i) $s_{i-1}(p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}s_{i-1})s_{i-1} = p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}},$
- (ii) $s_{i-1}p_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_{i-1} = s_ip_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}},$
- (iii) $s_{i-1}p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}}s_{i-1} = p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}},$
- (iv) $s_{i-1}(p_{i+\frac{1}{2}}s_{i-1}L_{i-1}s_{i-1})s_{i-1} = s_ip_{i-\frac{1}{2}}L_{i-1}s_i,$
- (v) $s_{i-1}p_{i+\frac{1}{2}}\sigma_is_{i-1} = \sigma_ip_{i+\frac{1}{2}},$
- (vi) $s_{i-1}p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_ip_{i+\frac{1}{2}}s_{i-1} = p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_ip_{i+\frac{1}{2}},$
- (vii) $s_{i-1}p_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1} = s_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_i,$
- (viii) $s_{i-1}p_{i+\frac{1}{2}}s_{i-1}L_{i-1}s_{i-1}p_ip_{i+\frac{1}{2}}s_{i-1} = s_ip_{i-\frac{1}{2}}L_{i-1}p_{i-1}p_{i-\frac{1}{2}}s_i,$
- (ix) $s_{i-1}p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}s_{i-1} = p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_{i+\frac{1}{2}}p_ip_{i-\frac{1}{2}}.$

The left hand side of (i) is

$$s_{i-1}(p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}s_{i-1})s_{i-1} = p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}$$

and the right hand side of (i) is

$$p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}p_ip_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}.$$

The left hand side of (ii) is

$$s_{i-1}p_{2+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_{i-1} = s_ip_{i-\frac{1}{2}}s_iL_{i-1}p_{i-\frac{1}{2}} = s_ip_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}},$$

which is the same as the right hand side of (ii). The left hand side of (iii) is

$$s_{i-1}p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}}s_{i-1} = p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}p_ip_{i-\frac{1}{2}},$$

which is the same as the right hand side of (iii). The left hand side of (iv) is

$$s_{i-1}(p_{i+\frac{1}{2}}s_{i-1}L_{i-1}s_{i-1})s_{i-1} = s_ip_{i-\frac{1}{2}}s_iL_{i-1} = s_ip_{i-\frac{1}{2}}L_{i-1}s_i,$$

which is the same as the right hand side of (iv). The left hand side of (v) is

$$s_{i-1}p_{i+\frac{1}{2}}\sigma_is_{i-1} = s_{i-1}p_{i+\frac{1}{2}}\sigma_{i-\frac{1}{2}} = s_{i-1}\sigma_{i-\frac{1}{2}}p_{i+\frac{1}{2}} = \sigma_ip_{i+\frac{1}{2}},$$

which is the same as the right hand side of (v). The left hand side of (vi) is

$$\begin{aligned} s_{i-1}p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_ip_{i+\frac{1}{2}}s_{i-1} &= p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}s_{i-1}p_is_{i-1}s_ip_{i-\frac{1}{2}}s_i \\ &= p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}p_{i-1}s_ip_{i-\frac{1}{2}}s_i \\ &= p_{i+\frac{1}{2}}p_{i-\frac{1}{2}}L_{i-1}p_{i-1}p_{i-\frac{1}{2}}, \end{aligned}$$

which is the same as the right hand side of (vi). The left hand side of (vii) is

$$\begin{aligned} s_{i-1}p_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1} &= s_ip_{i-\frac{1}{2}}s_iL_{i-1}p_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1} \\ &= s_ip_{i-\frac{1}{2}}L_{i-1}s_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1} = s_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}p_{i+\frac{1}{2}}s_{i-1}s_ip_{i-\frac{1}{2}}s_i \\ &= s_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}s_ip_{i-\frac{1}{2}}s_i = s_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}s_ip_{i-\frac{1}{2}}s_i \\ &= s_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}\sigma_{i-\frac{1}{2}}p_ip_{i-\frac{1}{2}}s_i = s_ip_{i-\frac{1}{2}}p_ip_{i+\frac{1}{2}}s_{i-1}L_{i-1}p_{i-\frac{1}{2}}s_i, \end{aligned}$$

which is the same as the right hand side of (vii). The left hand side of (viii) is

$$\begin{aligned} s_{i-1}p_{i+\frac{1}{2}}s_{i-1}L_{i-1}s_{i-1}p_ip_{i+\frac{1}{2}}s_{i-1} &= s_ip_{i-\frac{1}{2}}s_iL_{i-1}s_{i-1}p_is_{i-1}s_ip_{i-\frac{1}{2}}s_i \\ &= s_ip_{i-\frac{1}{2}}L_{i-1}s_is_{i-1}p_is_{i-1}s_ip_{i-\frac{1}{2}}s_i \\ &= s_ip_{i-\frac{1}{2}}L_{i-1}p_{i-1}p_{i-\frac{1}{2}}s_i, \end{aligned}$$

which is the same as the right hand side of (viii). Since the statement (ix) is evident from the relation $s_{i-1}p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}s_{i-1} = p_{i-\frac{1}{2}}$, the proof of part (7) is complete.

(9) By parts (6) and (8), $L_{i+\frac{1}{2}}$ commutes with $\langle p_i, s_{i-1} \rangle$, and so with $p_{i-1} = s_{i-1}p_i s_{i-1}$. \square

Theorem 3.7. *If $i = 1, 2, \dots$, then*

- (1) L_{i+1} commutes with $\mathcal{A}_{i+\frac{1}{2}}(z)$, and σ_{i+1} commutes with $\mathcal{A}_{i-\frac{1}{2}}(z)$,
- (2) $L_{i+\frac{1}{2}}$ commutes with $\mathcal{A}_i(z)$, and $\sigma_{i+\frac{1}{2}}$ commutes with $\mathcal{A}_{i-1}(z)$.

Consequently, the family of elements $(L_i, L_{i+\frac{1}{2}} : i = 0, 1, \dots)$ is pairwise commutative.

Proof. (1) Observe that L_1 commutes with $\mathcal{A}_{1-\frac{1}{2}}(z)$ and L_2 commutes with $\mathcal{A}_{1+\frac{1}{2}}(z)$, while σ_2 commutes with $\mathcal{A}_{1-\frac{1}{2}}(z)$ and σ_3 commutes with $\mathcal{A}_{2-\frac{1}{2}}(z)$. Since

$$\begin{aligned} L_{i+1} \text{ commutes with } \langle s_{i-1}, p_{i-1}, p_i, p_{i-\frac{1}{2}}, p_{i+\frac{1}{2}} \rangle, & \text{ if } i \geq 2, \text{ and} \\ \sigma_{i+1} \text{ commutes with } \langle s_{i-2}, p_{i-2}, p_{i-1}, p_{i-\frac{3}{2}}, p_{i-\frac{1}{2}} \rangle, & \text{ if } i \geq 3, \end{aligned}$$

it suffices to show that

- (i) L_{i+1} commutes with $\mathcal{A}_{i-\frac{3}{2}}(z)$, if $i \geq 2$, and
- (ii) σ_{i+1} commutes with $\mathcal{A}_{i-\frac{5}{2}}(z)$, if $i \geq 3$.

If $i \geq 2$, then induction on i shows that L_{i+1} commutes with $\mathcal{A}_{i-2}(z)$, while, if $i \geq 3$, the fact that L_{i+1} commutes with $p_{i-\frac{3}{2}}$ follows from induction on i , and the fact that σ_{i+1} commutes with $p_{i-\frac{3}{2}}$. Similarly, if $i \geq 3$, then induction on i shows that σ_{i+1} commutes with $\mathcal{A}_{i-3}(z)$, and that, if $i \geq 4$, then σ_{i+1} commutes with $p_{i-\frac{5}{2}}$.

(2) Observe that $L_{0+\frac{1}{2}}$ commutes with $\mathcal{A}_0(z)$ and $L_{1+\frac{1}{2}}$ commutes with $\mathcal{A}_1(z)$, while $\sigma_{1+\frac{1}{2}}$ commutes with $\mathcal{A}_0(z)$ and $\sigma_{2+\frac{1}{2}}$ commutes with $\mathcal{A}_1(z)$. Since

$$\begin{aligned} L_{i+\frac{1}{2}} \text{ commutes with } \langle s_{i-1}, p_{i-1}, p_i, p_{i-\frac{1}{2}} \rangle, & \text{ if } i \geq 2, \text{ and} \\ \sigma_{i+\frac{1}{2}} \text{ commutes with } \langle s_{i-2}, p_{i-2}, p_{i-1}, p_{i-\frac{3}{2}} \rangle, & \text{ if } i \geq 3, \end{aligned}$$

it suffices to show that

- (i) $L_{i+\frac{1}{2}}$ commutes with $\mathcal{A}_{i-\frac{3}{2}}(z)$, if $i \geq 2$, and
- (ii) $\sigma_{i+\frac{1}{2}}$ commutes with $\mathcal{A}_{i-\frac{5}{2}}(z)$, if $i \geq 3$.

If $i \geq 2$, then induction on i shows that $L_{i+\frac{1}{2}}$ commutes with $\mathcal{A}_{i-2}(z)$, while, if $i \geq 3$, the fact that $L_{i+\frac{1}{2}}$ commutes with $p_{i-\frac{3}{2}}$ follows from induction on i , and the fact that $\sigma_{i+\frac{1}{2}}$ commutes with $p_{i-\frac{3}{2}}$. Similarly, if $i \geq 3$, then induction on i shows that $\sigma_{i+\frac{1}{2}}$ commutes with $\mathcal{A}_{i-3}(z)$, and that, if $i \geq 4$, then $\sigma_{i+\frac{1}{2}}$ commutes with $p_{i-\frac{5}{2}}$. \square

Proposition 3.8. *For $i = 1, 2, \dots$, the following statements hold:*

- (1) $(L_{i+\frac{1}{2}} + L_{i+1})p_{i+1} = p_{i+1}(L_{i+\frac{1}{2}} + L_{i+1}) = zp_{i+1}$,
- (2) $(L_i + L_{i+\frac{1}{2}})p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}(L_i + L_{i+\frac{1}{2}}) = zp_{i+\frac{1}{2}}$,
- (3) $(L_{i-\frac{1}{2}} + L_i + L_{i+\frac{1}{2}} + L_{i+1})s_i = s_i(L_{i-\frac{1}{2}} + L_i + L_{i+\frac{1}{2}} + L_{i+1})$.

Proof. (1) It suffices to observe that:

- (i) $L_i p_{i+\frac{1}{2}} p_{i+1} = \sigma_{i+1} p_{i+1}$,
- (ii) $p_{i+\frac{1}{2}} L_i p_{i+1} = p_{i+\frac{1}{2}} L_i p_{i+1} p_{i+\frac{1}{2}} p_{i+1}$,
- (iii) $p_{i+\frac{1}{2}} L_i p_i p_{i+\frac{1}{2}} p_{i+1} = p_{i+\frac{1}{2}} L_i s_i p_{i+1}$,
- (iv) $s_i L_{i-\frac{1}{2}} s_i p_{i+1} + s_i L_i s_i p_{i+1} = zp_{i+1}$,
- (v) $\sigma_{i+\frac{1}{2}} p_{i+1} = s_i L_i p_{i+\frac{1}{2}} p_{i+1}$.

With the exception of (iv), each of the statements above is evident from the defining relations or from what we have already shown. Since $(L_{\frac{1}{2}} + L_1)p_1 = zp_1$,

$$(3.8) \quad s_i L_{i-\frac{1}{2}} s_i p_{i+1} + s_i L_i s_i p_{i+1} = s_i (L_{i-\frac{1}{2}} p_i + L_i p_i) s_i = z s_i p_i s_i = z p_{i+1},$$

gives (iv) by induction.

(2) Using the expression (3.8), we have

$$\begin{aligned} z p_{i+\frac{1}{2}} &= z p_{i+\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} \\ &= p_{i+\frac{1}{2}} (s_i L_{i-\frac{1}{2}} s_i p_{i+1} + s_i L_i s_i p_{i+1}) p_{i+\frac{1}{2}} \\ &= p_{i+\frac{1}{2}} L_{i-\frac{1}{2}} s_i p_{i+1} p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} L_i s_i p_{i+1} p_{i+\frac{1}{2}} \\ &= L_{i-\frac{1}{2}} p_{i+\frac{1}{2}} p_i p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} L_i p_i p_{i+\frac{1}{2}} \\ &= L_{i-\frac{1}{2}} p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} L_i p_i p_{i+\frac{1}{2}}, \end{aligned}$$

which yields

$$\begin{aligned} (L_i + L_{i+\frac{1}{2}}) p_{i+\frac{1}{2}} &= L_i p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}} L_i p_{i+\frac{1}{2}} - L_i p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} L_i p_i p_{i+\frac{1}{2}} \\ &\quad + s_i L_{i-\frac{1}{2}} p_{i+\frac{1}{2}} + \sigma_{i+\frac{1}{2}} p_{i+\frac{1}{2}} \\ &= -p_{i+\frac{1}{2}} + z p_{i+\frac{1}{2}} - L_{i-\frac{1}{2}} p_{i+\frac{1}{2}} + L_{i-\frac{1}{2}} p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} \\ &= z p_{i+\frac{1}{2}}. \end{aligned}$$

(3) Since $\sigma_{i+1} s_i = \sigma_{i+\frac{1}{2}} = s_i \sigma_{i+1}$, we have

$$\begin{aligned} (L_{i-\frac{1}{2}} + L_i + L_{i+\frac{1}{2}} + L_{i+1}) s_i &= L_{i-\frac{1}{2}} s_i + L_i s_i - p_{i+\frac{1}{2}} L_i s_i - L_i p_{i+\frac{1}{2}} + p_{i+\frac{1}{2}} L_i p_i p_{i+\frac{1}{2}} \\ &\quad + s_i L_{i-\frac{1}{2}} + \sigma_{i+\frac{1}{2}} s_i - p_{i+\frac{1}{2}} L_i - s_i L_i p_{i+\frac{1}{2}} \\ &\quad + p_{i+\frac{1}{2}} L_i p_{i+1} p_{i+\frac{1}{2}} + s_i L_i + \sigma_{i+1} s_i \\ &= s_i (L_{i-\frac{1}{2}} + L_i + L_{i+\frac{1}{2}} + L_{i+1}), \end{aligned}$$

as required. \square

Theorem 3.9. If $k = 0, 1, \dots$, then

- (1) the element $z_{k+\frac{1}{2}} = L_{\frac{1}{2}} + L_1 + L_{1+\frac{1}{2}} + \dots + L_{k+\frac{1}{2}}$ is central in $\mathcal{A}_{k+\frac{1}{2}}(z)$,
- (2) the element $z_{k+1} = L_{\frac{1}{2}} + L_1 + L_{1+\frac{1}{2}} + \dots + L_{k+1}$ is central in $\mathcal{A}_{k+1}(z)$.

Proof. (1) We first show that $z_{k+\frac{1}{2}}$ commutes with p_1, \dots, p_k and $p_{1+\frac{1}{2}}, \dots, p_{k+\frac{1}{2}}$. If $i = 1, \dots, k$, then p_i commutes with z_{i-1} and with $L_{i+\frac{1}{2}} + L_{i+1} + \dots + L_{k+\frac{1}{2}}$. Since p_i also commutes with $L_{i-\frac{1}{2}} + L_i$, it follows that p_i commutes with $z_{k+\frac{1}{2}}$. Similarly, since $p_{i+\frac{1}{2}}$ commutes with $z_{i-\frac{1}{2}}$, $L_i + L_{i+\frac{1}{2}}$, and with $L_{i+1} + L_{i+\frac{3}{2}} + \dots + L_{k+\frac{1}{2}}$, it follows that $p_{i+\frac{1}{2}}$ commutes with $z_{k+\frac{1}{2}}$. Next, suppose that $i = 1, \dots, k-1$. Since s_i commutes with z_{i-1} , $L_{i-\frac{1}{2}} + L_i + L_{i+\frac{1}{2}} + L_{i+1}$, and with $L_{i+\frac{3}{2}} + L_{i+2} + \dots + L_{k+\frac{1}{2}}$, it is evident that s_i commutes with $z_{k+\frac{1}{2}}$. As $\mathcal{A}_{k+\frac{1}{2}}(z)$ is generated by p_1, \dots, p_k , $p_{1+\frac{1}{2}}, \dots, p_{k+\frac{1}{2}}$, and s_1, \dots, s_{k-1} , we conclude that $z_{k+\frac{1}{2}}$ is central in $\mathcal{A}_{k+\frac{1}{2}}(z)$.

(2) Given that $z_{k+\frac{1}{2}}$ is central in $\mathcal{A}_{k+\frac{1}{2}}(z)$, it suffices to observe that s_k commutes with z_{k+1} . \square

Proposition 3.10. For $i = 1, 2, \dots$, the following statements hold:

- (1) $(L_i + L_{i+\frac{1}{2}} + L_{i+1}) \sigma_{i+1} = \sigma_{i+1} (L_i + L_{i+\frac{1}{2}} + L_{i+1})$,
- (2) $(L_{i-\frac{1}{2}} + L_i + L_{i+\frac{1}{2}}) \sigma_{i+\frac{1}{2}} = \sigma_{i+\frac{1}{2}} (L_{i-\frac{1}{2}} + L_i + L_{i+\frac{1}{2}})$.

Proof. (1) Since σ_{i+1} commutes with

$$z_{i+1} = z_{i-1} + L_{i-\frac{1}{2}} + L_i + L_{i+\frac{1}{2}} + L_{i+1},$$

and σ_{i+1} also commutes with $z_{i-1} + L_{i-\frac{1}{2}} \in \mathcal{A}_{i-\frac{1}{2}}(z)$, it follows that σ_{i+1} commutes with $L_i + L_{i+\frac{1}{2}} + L_{i+1}$.

(2) Given that $\sigma_{1+\frac{1}{2}} = 1$, we may suppose that $i = 2, 3, \dots$. Since $\sigma_{i+\frac{1}{2}}$ commutes with

$$z_{i+\frac{1}{2}} = z_{i-\frac{3}{2}} + L_{i-1} + L_{i-\frac{1}{2}} + L_i + L_{i+\frac{1}{2}},$$

and $\sigma_{i+\frac{1}{2}}$ also commutes with $z_{i-\frac{3}{2}} + L_{i-1} \in \mathcal{A}_{i-1}(z)$, it follows that $\sigma_{i+\frac{1}{2}}$ commutes with $L_{i-\frac{1}{2}} + L_i + L_{i+\frac{1}{2}}$. \square

4. A PRESENTATION FOR THE PARTITION ALGEBRAS

The goal here is to rewrite the presentation for $\mathcal{A}_k(z)$ given by [HR] in Theorem 2.1 in terms of the elements $\sigma_i, \sigma_{i+\frac{1}{2}}, p_i, p_{i+\frac{1}{2}}$.

Theorem 4.1. *If $k = 1, 2, \dots$, then $\mathcal{A}_k(z)$ is the unital associative algebra presented by the generators*

$$p_1, p_2, \dots, p_k, p_{1+\frac{1}{2}}, \dots, p_{k-\frac{1}{2}}, \sigma_2, \sigma_3, \dots, \sigma_k, \sigma_{2+\frac{1}{2}}, \sigma_{3+\frac{1}{2}}, \dots, \sigma_{k-\frac{1}{2}},$$

and the relations:

- (1) (Involutions)
 - (a) $\sigma_{i+\frac{1}{2}}^2 = 1$, for $i = 2, \dots, k-1$.
 - (b) $\sigma_{i+1}^2 = 1$, for $i = 1, \dots, k-1$.
- (2) (Braid-like relations)
 - (a) $\sigma_{i+1}\sigma_{j+\frac{1}{2}} = \sigma_{j+\frac{1}{2}}\sigma_{i+1}$, if $j \neq i+1$.
 - (b) $\sigma_i\sigma_j = \sigma_j\sigma_i$, if $j \neq i+1$.
 - (c) $\sigma_{i+\frac{1}{2}}\sigma_{j+\frac{1}{2}} = \sigma_{j+\frac{1}{2}}\sigma_{i+\frac{1}{2}}$, if $j \neq i+1$.
 - (d) $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$, for $i = 1, \dots, k-2$, if $s_{j-1} = \sigma_{j-\frac{1}{2}}\sigma_j$, for $j = 2, \dots, k$.
- (3) (Idempotent relations)
 - (a) $p_i^2 = zp_i$, for $i = 1, \dots, k$.
 - (b) $p_{i+\frac{1}{2}}^2 = p_{i+\frac{1}{2}}$, for $i = 1, \dots, k-1$.
 - (c) $\sigma_{i+1}p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}\sigma_{i+1} = p_{i+\frac{1}{2}}$, for $i = 1, \dots, k-1$.
 - (d) $\sigma_{i+\frac{1}{2}}p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}\sigma_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}$, for $i = 1, \dots, k-1$.
 - (e) $\sigma_{i+\frac{1}{2}}p_i p_{i+1} = \sigma_{i+1}p_i p_{i+1}$, for $i = 1, \dots, k-1$.
 - (f) $p_i p_{i+1} \sigma_{i+\frac{1}{2}} = p_i p_{i+1} \sigma_{i+1}$, for $i = 1, \dots, k-1$.
- (4) (Commutation relations)
 - (a) $p_i p_j = p_j p_i$, for $i, j = 1, \dots, k$.
 - (b) $p_{i+\frac{1}{2}} p_{j+\frac{1}{2}} = p_{j+\frac{1}{2}} p_{i+\frac{1}{2}}$, for $i, j = 1, \dots, k-1$.
 - (c) $p_{i+\frac{1}{2}} p_j = p_j p_{i+\frac{1}{2}}$, for $j \neq i, i+1$.
 - (d) $\sigma_i p_j = p_j \sigma_i$ if $j \neq i-1, i$.
 - (e) $\sigma_i p_{j+\frac{1}{2}} = p_{j+\frac{1}{2}} \sigma_i$, if $j \neq i$.
 - (f) $\sigma_{i+\frac{1}{2}} p_j = p_j \sigma_{i+\frac{1}{2}}$, if $j \neq i, i+1$.
 - (g) $\sigma_{i+\frac{1}{2}} p_{j+\frac{1}{2}} = p_{j+\frac{1}{2}} \sigma_{i+\frac{1}{2}}$, if $j \neq i-1$.
 - (h) $\sigma_{i+\frac{1}{2}} p_i \sigma_{i+\frac{1}{2}} = \sigma_{i+1} p_{i+1} \sigma_{i+1}$, for $i = 1, \dots, k-1$.
 - (i) $\sigma_{i+\frac{1}{2}} p_{i-\frac{1}{2}} \sigma_{i+\frac{1}{2}} = \sigma_i p_{i+\frac{1}{2}} \sigma_i$, for $i = 2, \dots, k-1$.
- (5) (Contraction relations)
 - (a) $p_{i+\frac{1}{2}} p_j p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}$, for $j = i, i+1$.
 - (b) $p_i p_{j-\frac{1}{2}} p_i = p_i$, for $j = i, i+1$.

Proof. We first show that the relations given in the statement above are a consequence the presentation given by [HR] in Theorem 2.1.

(1a), (1b) Proposition 4.2 shows that $\sigma_{i+\frac{1}{2}}$ and σ_{i+1} are involutions in $\mathcal{A}_k(z)$.

(2a) To see that σ_{i+1} and $\sigma_{i+\frac{1}{2}}$ commute, we use Proposition 3.3 and Proposition 4.2 to obtain

$$(4.1) \quad \sigma_{i+1} = \sigma_{i+\frac{1}{2}} s_i = s_i \sigma_{i+\frac{1}{2}} \quad \text{and} \quad \sigma_{i+\frac{1}{2}} \sigma_{i+1} = \sigma_{i+1} \sigma_{i+\frac{1}{2}} = s_i \quad (\text{for } i = 1, \dots, k-1)$$

where s_i is a Coxeter generator for $\mathfrak{S}_k \subset \mathcal{A}_k(z)$. Theorem 3.7 shows that σ_j commutes with $\mathcal{A}_{j-\frac{3}{2}}(z)$, and hence that σ_j commutes with $\sigma_{1+\frac{1}{2}}, \dots, \sigma_{j-\frac{3}{2}}$. From Theorem 3.7, $\sigma_{i+\frac{1}{2}}$ commutes with $\mathcal{A}_{i-1}(z)$, and hence with $\sigma_2, \dots, \sigma_{i-1}$. In general however, σ_i and $\sigma_{i+\frac{1}{2}}$ do not commute.

(2b), (2c) Theorem 3.7 shows that σ_{i+1} commutes with $\sigma_2, \dots, \sigma_{i-1}$ and that $\sigma_{i+\frac{1}{2}}$ commutes with $\sigma_{1+\frac{1}{2}}, \dots, \sigma_{i-\frac{3}{2}}$.

(2d) Follows from (4.1), whereby for $j = 1, \dots, k-1$, each product $\sigma_{j+\frac{1}{2}} \sigma_{j+1} = s_j$ is a Coxeter generator for $\mathfrak{S}_k \subset \mathcal{A}_k(z)$.

(3a), (3b) Are included in the set of relations given by [HR].

(3c), (3d) That $\sigma_{i+1} p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}} \sigma_{i+1} = p_{i+\frac{1}{2}}$ is given in Proposition 3.1. Proposition 3.3 shows that $\sigma_{i+\frac{1}{2}} p_{i+\frac{1}{2}} = s_i \sigma_{i+1} p_{i+\frac{1}{2}} = s_i p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}$.

(3e), (3f) Proposition 3.3 and Proposition 4.2 show that

$$p_i p_{i+1} = s_i p_i p_{i+1} = \sigma_{i+\frac{1}{2}} \sigma_{i+1} p_i p_{i+1} \quad \text{and} \quad \sigma_{i+\frac{1}{2}} p_i p_{i+1} = \sigma_{i+1} p_i p_{i+1}.$$

(4a)–(4c) Are included in the set of relations given by [HR].

(4d), (4e) By Theorem 3.7, σ_i commutes with $\mathcal{A}_{i-\frac{3}{2}}(z)$, and hence with p_1, \dots, p_{i-2} and with $p_{1+\frac{1}{2}}, \dots, p_{i-\frac{3}{2}}$. Proposition 3.1 shows that σ_i commutes with $p_{i-\frac{1}{2}}$.

(4f), (4g) By Theorem 3.7, $\sigma_{i+\frac{1}{2}}$ commutes with $\mathcal{A}_{i-1}(z)$, and hence with p_1, \dots, p_{i-1} and with $p_{1+\frac{1}{2}}, \dots, p_{i-\frac{3}{2}}$. From (3d), it follows that $\sigma_{i+\frac{1}{2}}$ commutes with $p_{i+\frac{1}{2}}$.

(4h) Proposition 3.3 and Proposition 4.2 show that

$$p_i = s_i p_{i+1} s_i = \sigma_{i+\frac{1}{2}} \sigma_{i+1} p_{i+1} \sigma_{i+1} \sigma_{i+\frac{1}{2}} \quad \text{and} \quad \sigma_{i+\frac{1}{2}} p_i \sigma_{i+\frac{1}{2}} = \sigma_{i+1} p_{i+1} \sigma_{i+1}.$$

(4i) Proposition 3.1 shows that $p_{i-\frac{1}{2}} s_i \sigma_i = s_i \sigma_i p_{i+\frac{1}{2}}$. Proposition 3.3 and Proposition 4.2, together with the fact that σ_{i+1} commutes with $p_{i-\frac{1}{2}}$ give

$$p_{i-\frac{1}{2}} \sigma_{i+1} \sigma_{i+\frac{1}{2}} \sigma_i = \sigma_{i+1} \sigma_{i+\frac{1}{2}} \sigma_i p_{i+\frac{1}{2}} \quad \text{and} \quad p_{i-\frac{1}{2}} \sigma_{i+\frac{1}{2}} \sigma_i = \sigma_{i+\frac{1}{2}} \sigma_i p_{i+\frac{1}{2}}.$$

Multiplying both sides of the last expression by $\sigma_i \sigma_{i+\frac{1}{2}}$ on the left, and using Proposition 4.2 once more shows that

$$\sigma_i \sigma_{i+\frac{1}{2}} p_{i-\frac{1}{2}} \sigma_{i+\frac{1}{2}} \sigma_i = \sigma_i \sigma_{i+\frac{1}{2}} \sigma_{i+\frac{1}{2}} \sigma_i p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}} \quad \text{and} \quad \sigma_{i+\frac{1}{2}} p_{i-\frac{1}{2}} \sigma_{i+\frac{1}{2}} = \sigma_i p_{i+\frac{1}{2}} \sigma_i,$$

as required.

(5a), (5b) Are included in the set of relations given by [HR].

Next, we derive the relations given by [HR] in Theorem 2.1 from the relations (1a)–(5b) above.

(1i) By the relations (1a), (1b) and (2a),

$$\sigma_{i+1} \sigma_{i+\frac{1}{2}} = \sigma_{i+\frac{1}{2}} \sigma_{i+\frac{1}{2}} \quad \text{and} \quad (\sigma_{i+\frac{1}{2}} \sigma_{i+1})^2 = 1, \quad \text{for } i = 1, \dots, k-1.$$

Thus, writing $s_i = \sigma_{i+\frac{1}{2}} \sigma_{i+1}$, for $i = 1, \dots, k-1$, we recover (1i).

(1ii) If $j \neq i+1$, then, by (2b) and (2c),

$$s_i s_j = \sigma_{i+\frac{1}{2}} \sigma_{i+1} \sigma_{j+\frac{1}{2}} \sigma_{j+1} = \sigma_{j+\frac{1}{2}} \sigma_{j+1} \sigma_{i+\frac{1}{2}} \sigma_{i+1} = s_j s_i,$$

as required.

(1iii) Is equivalent to (2d) with $s_i = \sigma_{i+\frac{1}{2}} \sigma_{i+1}$, for $i = 1, \dots, k-1$.

(2i), (2ii) Are identical to the relations (3a) and (3b).

(2iii) With $s_i = \sigma_{i+1}\sigma_{i+\frac{1}{2}}$, the relations (3c) and (3d) give

$$s_i p_{i+\frac{1}{2}} = \sigma_{i+\frac{1}{2}} \sigma_{i+1} p_{i+\frac{1}{2}} = \sigma_{i+\frac{1}{2}} p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}}$$

and

$$p_{i+\frac{1}{2}} s_i = p_{i+\frac{1}{2}} \sigma_{i+1} \sigma_{i+\frac{1}{2}} = p_{i+\frac{1}{2}} \sigma_{i+\frac{1}{2}} = p_{i+\frac{1}{2}},$$

as required.

(2iv) With $s_i = \sigma_{i+1}\sigma_{i+\frac{1}{2}}$, the relations (2a), (3e) and (3f) give

$$s_i p_i p_{i+1} = \sigma_{i+1} \sigma_{i+\frac{1}{2}} p_i p_{i+1} = p_i p_{i+1} \quad \text{and} \quad p_i p_{i+1} s_i = p_i p_{i+1} \sigma_{i+1} \sigma_{i+\frac{1}{2}} = p_i p_{i+1},$$

as required.

(3i)-(3iii) Are identical to the relations (4a)-(4c).

(3iv) If $j \neq i, i+1$, then the relations (4d) and (4f) give

$$s_i p_j = \sigma_{i+\frac{1}{2}} \sigma_{i+1} p_j = \sigma_{i+\frac{1}{2}} p_j \sigma_{i+1} = p_j \sigma_{i+\frac{1}{2}} \sigma_{i+1} = p_j s_i,$$

as required.

(3v) If $j \neq i-1, i+1$, then the relations (4e) and (4f) give

$$s_i p_{j+\frac{1}{2}} = \sigma_{i+\frac{1}{2}} \sigma_{i+1} p_{j+\frac{1}{2}} = \sigma_{i+\frac{1}{2}} p_{j+\frac{1}{2}} \sigma_{i+1} = p_{j+\frac{1}{2}} \sigma_{i+\frac{1}{2}} \sigma_{i+1} = p_{j+\frac{1}{2}} s_i,$$

as required.

(3vi) From the relations (1b) and (4h),

$$s_i p_i s_i = \sigma_{i+1} \sigma_{i+\frac{1}{2}} p_i \sigma_{i+\frac{1}{2}} \sigma_{i+1} = \sigma_{i+1}^2 p_{i+1} \sigma_{i+1}^2 = p_{i+1},$$

as required.

(3vii) From the relations (4e), (4g) and (1a), (1b),

$$s_i p_{i-\frac{1}{2}} s_i = \sigma_{i+\frac{1}{2}} \sigma_{i+1} p_{i-\frac{1}{2}} \sigma_{i+1} \sigma_{i+\frac{1}{2}} = \sigma_i \sigma_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} \sigma_i = s_{i-1} p_{i+\frac{1}{2}} s_{i-1},$$

as required.

(4i), (4ii) Are identical to the relations (5a) and (5b). \square

Proposition 4.2. *If $i = 0, 1, \dots$, then $(\sigma_{i+\frac{1}{2}})^2 = 1$ and $(\sigma_{i+1})^2 = 1$.*

Proof. Given that $\sigma_{i+\frac{1}{2}} s_i = s_i \sigma_{i+\frac{1}{2}} = \sigma_{i+1}$, we obtain

$$\sigma_{i+\frac{1}{2}}^2 = (s_i \sigma_{i+1})^2 = \sigma_{i+1}^2.$$

It therefore suffices to show that $\sigma_{i+1}^2 = 1$. By definition $\sigma_1 = 1$, so we assume that $\sigma_i^2 = 1$ proceed by induction. After taking the square of the right hand side of the definition (3.2), the statement that $\sigma_{i+1}^2 = 1$ is equivalent to the following set of relations:

$$(4.2) \quad (p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i)^2 = (p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i)(p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}});$$

$$(4.3) \quad (p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i)(s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}}) \\ = (p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i)(s_{i-1} s_i \sigma_i s_i s_{i-1});$$

$$(4.4) \quad (p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i)(s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i) \\ = (s_{i-1} s_i \sigma_i s_i s_{i-1})(s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i);$$

$$(4.5) \quad (s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}}) (p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i) \\ = (s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}}) (s_{i-1} s_i \sigma_i s_i s_{i-1});$$

$$(4.6) \quad (s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}})^2 = (p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}}) (s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}})$$

$$(4.7) \quad (s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}}) (p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}}) = (s_{i-1} s_i \sigma_i s_i s_{i-1}) (p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}});$$

$$(4.8) \quad (p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}}) (p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i) = (p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}}) (s_{i-1} s_i \sigma_i s_i s_{i-1});$$

$$(4.9) \quad (s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}}) (s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i) \\ = (p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}}) (s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i);$$

$$(4.10) \quad (p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}})^2 = (s_{i-1} s_i \sigma_i s_i s_{i-1}) (s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}});$$

$$(4.11) \quad (s_{i-1} s_i \sigma_i s_i s_{i-1}) (p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i) = (s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i)^2;$$

$$(4.12) \quad (s_{i-1} s_i \sigma_i s_i s_{i-1})^2 = 1;$$

$$(4.13) \quad (s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i) (p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i) \\ = (s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i) (p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}});$$

$$(4.14) \quad (s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i) (s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}}) \\ = (s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i) (s_{i-1} s_i \sigma_i s_i s_{i-1}).$$

Considering the left hand side of (4.2), using the fact that $p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}$, together with $p_{i+\frac{1}{2}} s_{i-1} p_{i+\frac{1}{2}} = p_{i-\frac{1}{2}} p_{i+\frac{1}{2}}$,

$$\begin{aligned} (p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i)^2 &= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i \\ &= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} p_{i+\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} \\ &= p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} \\ &= p_{i-\frac{1}{2}} p_{i+\frac{1}{2}}. \end{aligned}$$

Similarly, considering the right hand side of (4.2),

$$\begin{aligned} (p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i) (p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}}) &= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} \\ &= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} \\ &= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} p_{i+\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} \\ &= p_{i-\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} \\ &= p_{i-\frac{1}{2}} p_{i+\frac{1}{2}}, \end{aligned}$$

Which demonstrates (4.2). Now consider the left hand side of (4.3)

$$\begin{aligned} (p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i) (s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}}) \\ = p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} \\ = p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i-1} \sigma_i p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}}. \end{aligned}$$

Considering the right hand side of (4.3),

$$\begin{aligned}
(p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i) (s_{i-1} s_i \sigma_i s_i s_{i-1}) &= (p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} s_i) (s_{i-1} s_i \sigma_i s_i s_{i-1}) \\
&= p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i s_i s_{i-1} p_{i+\frac{1}{2}} \sigma_i s_i s_{i-1} \\
&= p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} p_{i+1} p_{i+\frac{1}{2}} \sigma_i s_i s_{i-1} \\
&= p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i+1} p_{i+\frac{1}{2}} \sigma_i s_i s_{i-1} \\
&= p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_i s_i p_{i+\frac{1}{2}} \sigma_i s_i s_{i-1} \\
&= p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} \sigma_i s_i s_{i-1} \\
&= p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} s_{i-1} s_i s_{i-1} \\
&= p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} p_i \sigma_{i-\frac{1}{2}} p_{i+\frac{1}{2}} s_{i-1} s_i s_{i-1} \\
&= p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} p_i \sigma_{i-\frac{1}{2}} s_{i-1} s_i p_{i-\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} p_i \sigma_i s_i p_{i-\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} \\
&= (p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}}) (p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}}).
\end{aligned}$$

Now,

$$\begin{aligned}
(4.15) \quad (p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}}) (p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}}) &= p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} p_i \sigma_i s_i p_{i-\frac{1}{2}} s_i \sigma_i p_i p_{i-\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} p_i \sigma_i s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \sigma_i p_i p_{i-\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} p_i \sigma_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i p_{i-\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} p_i (\sigma_{i-\frac{1}{2}})^2 p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} \\
&= p_{i-\frac{1}{2}},
\end{aligned}$$

which completes the proof of (4.3). For the left hand side of (4.4),

$$\begin{aligned}
(p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i) (s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i) &= p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_i \sigma_i s_i p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i \sigma_i s_i p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i \sigma_i s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \\
&= p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i \sigma_{i-\frac{1}{2}} p_{i+\frac{1}{2}} s_{i-1} \\
&= p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} s_{i-1} \\
&= p_{i-\frac{1}{2}} L_{i-1} s_{i-1} \sigma_{i-\frac{1}{2}} p_{i+\frac{1}{2}} s_{i-1} \\
&= p_{i-\frac{1}{2}} p_i (\sigma_i)^2 p_{i+\frac{1}{2}} s_{i-1} \\
&= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1}.
\end{aligned}$$

Considering the right hand side of (4.4),

$$\begin{aligned}
(s_{i-1}s_i\sigma_i s_i s_{i-1})(s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i) &= s_{i-1}s_i\sigma_i p_{i+\frac{1}{2}} s_{i-1}s_i L_{i-1} s_i p_{i-\frac{1}{2}} s_i \\
&= s_{i-1}s_i\sigma_i p_{i+\frac{1}{2}} s_{i-1}s_i L_{i-1} s_i p_{i-\frac{1}{2}} s_i \\
&= s_{i-1}s_i s_{i-1}\sigma_{i-\frac{1}{2}} p_{i+\frac{1}{2}} s_{i-1}s_i L_{i-1} s_i p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} s_i s_{i-1}\sigma_{i-\frac{1}{2}} s_{i-1}s_i L_{i-1} s_i p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} s_i \sigma_{i-\frac{1}{2}} (s_{i-1})^2 (s_i)^2 L_{i-1} p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} s_i \sigma_{i-\frac{1}{2}} \sigma_i p_i p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} s_i s_{i-1} p_i p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} p_{i-1} s_i p_{i-\frac{1}{2}} s_i \\
&= p_{i-\frac{1}{2}} p_{i-1} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \\
&= p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1},
\end{aligned}$$

which demonstrates (4.4). Considering the left hand side of (4.5),

$$\begin{aligned}
(s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}})(q_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i) &= s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i \\
&= s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} s_i \\
&= s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i \\
&= s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \sigma_i p_i p_{i-\frac{1}{2}} p_i \sigma_i s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \\
&= s_{i-1} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i \sigma_{i-\frac{1}{2}} p_{i+\frac{1}{2}} s_{i-1} \\
&= s_{i-1} (\sigma_{i-\frac{1}{2}})^2 p_{i+\frac{1}{2}} s_{i-1} \\
&= s_{i-1} p_{i+\frac{1}{2}} s_{i-1},
\end{aligned}$$

while, for the right hand side of (4.5),

$$\begin{aligned}
(s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}})(s_{i-1}s_i\sigma_i s_i s_{i-1}) &= s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} s_i \sigma_i s_i s_{i-1} \\
&= s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i s_i s_{i-1} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} s_i s_{i-1} \\
&= s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i + 1 s_{i-1} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} s_i s_{i-1} \\
&= s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} p_{i+1} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} s_i s_{i-1} \\
&= s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i+1} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} s_i s_{i-1} \\
&= s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} p_i s_i p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} s_i s_{i-1} \\
&= s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \sigma_i p_i p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} s_i s_{i-1} \\
&= s_{i-1} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} s_i s_{i-1} \\
&= s_{i-1} \sigma_{i-\frac{1}{2}} p_{i+\frac{1}{2}} p_i p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} s_i s_{i-1} \\
&= s_{i-1} p_{i+\frac{1}{2}} (\sigma_{i-\frac{1}{2}})^2 s_{i-1} \\
&= s_{i-1} p_{i+\frac{1}{2}} s_{i-1},
\end{aligned}$$

which demonstrates (4.5). The statement (4.6) is equivalent to (4.2) which has already been verified. The right hand side of (4.7) comes to,

$$\begin{aligned}
(s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}})(p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}}) &= s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} p_{i-1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} \\
&= s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} \\
&= s_{i-1} p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} \\
&= s_{i-1} p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i-1} s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} \\
&= s_{i-1} p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i-1} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} \\
&= s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \sigma_i p_i p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} \sigma_i p_i p_{i-\frac{1}{2}} \\
&= s_{i-1} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i p_{i-\frac{1}{2}} \\
&= s_{i-1} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i p_{i-\frac{1}{2}} \\
&= s_{i-1} p_{i+\frac{1}{2}} (\sigma_{i-\frac{1}{2}})^2 p_i p_{i-\frac{1}{2}} \\
&= s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}},
\end{aligned}$$

while

$$\begin{aligned}
(s_{i-1} s_i \sigma_i s_i s_{i-1})(p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}}) &= s_{i-1} s_i \sigma_i s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} \\
&= s_{i-1} s_i \sigma_i s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} \\
&= s_{i-1} s_i \sigma_i s_i s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \sigma_i p_i p_{i-\frac{1}{2}} \\
&= s_{i-1} s_i \sigma_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i p_{i-\frac{1}{2}} \\
&= s_{i-1} s_i p_{i+\frac{1}{2}} (\sigma_{i-\frac{1}{2}})^2 p_i p_{i-\frac{1}{2}} \\
&= s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}},
\end{aligned}$$

as required. Since the statement (4.8) is equivalent to (4.7), we consider (4.9). Using the relation $p_{i-\frac{1}{2}} s_i p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}} p_{i+\frac{1}{2}}$,

$$\begin{aligned}
(s_i p_{i-\frac{1}{2}} L_{i-1} p_{i+\frac{1}{2}} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}})(s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i) &= s_i p_{i-\frac{1}{2}} L_{i-1} p_{i+\frac{1}{2}} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i \\
&= s_i p_{i-\frac{1}{2}} L_{i-1} p_{i+\frac{1}{2}} s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} \\
&= s_i p_{i-\frac{1}{2}} L_{i-1} p_{i+\frac{1}{2}} s_{i-1} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} \\
&= p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}} \\
&= p_{i+\frac{1}{2}} p_{i-\frac{1}{2}}.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
(p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}})(s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i) &= p_{i-\frac{1}{2}} L_{i-1} s_i p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i \\
&= p_{i+\frac{1}{2}} (p_{i-\frac{1}{2}} L_{i-1} p_{i-\frac{1}{2}})^2 \\
&= p_{i+\frac{1}{2}} p_{i-\frac{1}{2}}.
\end{aligned}$$

The left hand side of (4.10) is given by (4.15), and the right hand side by

$$\begin{aligned}
(s_{i-1}s_i\sigma_i s_i s_{i-1})(s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}}) &= s_{i-1}s_i\sigma_i s_i s_{i-1} s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} p_{i-1} p_{i-\frac{1}{2}} \\
&= s_{i-1}s_i\sigma_i s_i s_{i-1} s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{1}{2}} \\
&= s_{i-1}s_i\sigma_i s_i s_{i-1} s_i p_{i-\frac{1}{2}} s_i L_{i-1} p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{1}{2}} \\
&= s_{i-1}s_i\sigma_i s_i (s_{i-1})^2 p_{i+\frac{1}{2}} s_{i-1} \sigma_i p_i p_{i-\frac{1}{2}} s_i p_{i-1} p_{i-\frac{1}{2}} \\
&= s_{i-1}s_i s_{i-1} \sigma_{i-\frac{1}{2}} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i-1} s_i p_{i-\frac{1}{2}} \\
&= s_{i-1} p_{i-\frac{1}{2}} s_i s_{i-1} (\sigma_{i-\frac{1}{2}})^2 p_i p_{i-\frac{1}{2}} p_{i-1} s_i p_{i-\frac{1}{2}} \\
&= p_{i-\frac{1}{2}} s_i s_{i-1} p_i p_{i-\frac{1}{2}} p_{i-1} s_i p_{i-\frac{1}{2}} \\
&= p_{i-\frac{1}{2}},
\end{aligned}$$

as required. Considering the left hand side of (4.11),

$$\begin{aligned}
(s_{i-1}s_i\sigma_i s_i s_{i-1})(p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i) &= s_{i-1}s_i\sigma_i s_i p_{i-\frac{1}{2}} s_i p_{i+1} p_{i+\frac{1}{2}} s_{i-1} \sigma_i p_i p_{i-\frac{1}{2}} s_i \\
&= s_{i-1}s_i\sigma_i s_{i-1} p_{i+\frac{1}{2}} s_{i-1} p_{i+1} p_{i+\frac{1}{2}} s_{i-1} \sigma_i p_i p_{i-\frac{1}{2}} s_i \\
&= s_{i-1}s_i\sigma_{i-\frac{1}{2}} p_{i+\frac{1}{2}} s_{i-1} p_{i+1} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i p_{i-\frac{1}{2}} s_i \\
&= s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \sigma_{i-\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i p_{i-\frac{1}{2}} s_i \\
&= s_i p_{i-\frac{1}{2}} s_i p_{i+1} p_{i+\frac{1}{2}} (\sigma_{i-\frac{1}{2}})^2 p_i p_{i-\frac{1}{2}} s_i \\
&= s_i p_{i-\frac{1}{2}} s_i p_{i+1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} s_i \\
&= s_i p_{i-\frac{1}{2}} s_i,
\end{aligned}$$

which, by (4.15), is equal to the right hand side of (4.11). Since (4.9) is equivalent to (4.13), while (4.14) is equivalent to (4.4), the proof of the proposition is complete. \square

We record for later reference further consequences of the presentation given in Theorem 2.1.

Proposition 4.3. *For $i = 1, 2, \dots$, the following statements hold:*

- (1) $p_{i+1}\sigma_{i+1}p_{i+1} = L_i p_{i+1}$,
- (2) $p_{i+1}\sigma_{i+\frac{1}{2}}p_{i+1} = (z - L_{i-\frac{1}{2}})p_{i+1}$,
- (3) $p_{i+\frac{3}{2}}\sigma_{i+1}p_{i+\frac{3}{2}} = p_{i+\frac{1}{2}}p_{i+\frac{3}{2}}$.

Proof. (1) From Proposition 3.1, we obtain $p_{i+\frac{1}{2}}p_{i+1}\sigma_{i+1} = p_{i+\frac{1}{2}}L_i$. Thus

$$p_{i+1}\sigma_{i+1} = p_{i+1}p_{i+\frac{1}{2}}p_{i+1}\sigma_{i+1} = p_{i+1}p_{i+\frac{1}{2}}L_i \quad \text{and} \quad p_{i+1}\sigma_{i+1}p_{i+1} = p_{i+1}p_{i+\frac{1}{2}}L_i p_{i+1} = L_i p_{i+1},$$

as required.

(2) We first compute

$$\begin{aligned}
p_{i+1}s_i p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+1} &= s_i p_i p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+1} \\
&= s_i p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+1} \\
&= s_i p_i p_{i+\frac{1}{2}} \sigma_i p_{i-1} p_{i-\frac{1}{2}} p_i s_i \\
&= s_i p_i p_{i+\frac{1}{2}} \sigma_i s_{i-1} p_i s_i \\
&= s_i p_i p_{i+\frac{1}{2}} \sigma_{i-\frac{1}{2}} p_i s_i \\
&= \sigma_{i-\frac{1}{2}} p_{i+1}.
\end{aligned}$$

Now observe that $p_2\sigma_{1+\frac{1}{2}}p_2 = (z - L_{\frac{1}{2}})p_2$ and hence, by induction,

$$\begin{aligned}
p_{i+1}\sigma_{i+\frac{1}{2}}p_{i+1} &= s_i s_{i-1} p_i \sigma_{i-\frac{1}{2}} p_i s_{i-1} s_i + p_{i+1} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i+1} \\
&\quad + p_{i+1} s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_{i+1} - p_{i+1} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i+1} \\
&\quad - p_{i+1} s_i p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+1} \\
&= s_{i-1} s_i p_i \sigma_{i-\frac{1}{2}} p_i s_i s_{i-1} + p_{i-\frac{1}{2}} L_{i-1} p_{i+1} + L_{i-1} p_{i-\frac{1}{2}} p_{i+1} \\
&\quad - p_{i-\frac{1}{2}} L_{i-1} p_{i-1} p_{i-\frac{1}{2}} p_{i+1} - \sigma_{i-\frac{1}{2}} p_{i+1} \\
&= (z - L_{i-\frac{1}{2}})p_{i+1}.
\end{aligned}$$

(3) Observe that $p_{2+\frac{1}{2}}\sigma_2 p_{2+\frac{1}{2}} = p_{2+\frac{1}{2}}s_1 p_{2+\frac{1}{2}} = p_{2+\frac{1}{2}}p_{1+\frac{1}{2}}$ and, by induction,

$$\begin{aligned}
p_{i+\frac{3}{2}} s_{i-1} s_i \sigma_i s_i s_{i-1} p_{i+\frac{3}{2}} &= s_{i-1} p_{i+\frac{3}{2}} s_i \sigma_i s_i p_{i+\frac{3}{2}} s_{i-1} \\
&= s_{i-1} s_i s_{i+1} p_{i+\frac{1}{2}} s_{i+1} \sigma_i s_{i+1} p_{i+\frac{1}{2}} s_i s_{i+1} s_{i-1} \\
&= s_{i-1} s_i s_{i+1} p_{i+\frac{1}{2}} \sigma_i p_{i+\frac{1}{2}} s_i s_{i+1} s_{i-1} \\
&= s_{i-1} s_i s_{i+1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} s_i s_{i+1} s_{i-1} \\
&= p_{i+\frac{1}{2}} p_{i+\frac{3}{2}}.
\end{aligned}$$

Therefore, using the definition (3.2) and the fact that $p_{i-\frac{1}{2}}L_{i-1}p_{i-\frac{1}{2}} = p_{i-\frac{1}{2}}$,

$$\begin{aligned}
p_{i+\frac{3}{2}}\sigma_{i+1}p_{i+\frac{3}{2}} &= p_{i+\frac{3}{2}}p_{i+\frac{1}{2}} + p_{i+\frac{3}{2}}s_i p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i p_{i+\frac{3}{2}} \\
&\quad + p_{i+\frac{3}{2}}p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} p_{i+\frac{3}{2}} - p_{i+\frac{3}{2}}s_i p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} p_{i+\frac{3}{2}} \\
&\quad - p_{i+\frac{3}{2}}p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} s_i p_{i+\frac{3}{2}} \\
&= p_{i+\frac{3}{2}}p_{i+\frac{1}{2}} + p_{i+\frac{3}{2}}s_i p_{i+\frac{3}{2}} p_{i-\frac{1}{2}} L_{i-1} s_i p_{i-\frac{1}{2}} s_i \\
&\quad + p_{i-\frac{1}{2}} L_{i-1} p_{i+\frac{3}{2}} s_i p_{i+\frac{3}{2}} p_{i-\frac{1}{2}} - p_{i+\frac{3}{2}}s_i p_{i+\frac{3}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} \\
&\quad - p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{3}{2}} s_i p_{i+\frac{3}{2}} \\
&= p_{i+\frac{3}{2}}p_{i+\frac{1}{2}} + p_{i+\frac{3}{2}}p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} s_{i-1} \\
&\quad + p_{i-\frac{1}{2}} L_{i-1} p_{i+\frac{3}{2}} p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} - p_{i+\frac{3}{2}}p_{i+\frac{1}{2}} p_{i-\frac{1}{2}} L_{i-1} s_{i-1} p_{i+\frac{1}{2}} p_i p_{i-\frac{1}{2}} \\
&\quad - p_{i-\frac{1}{2}} p_i p_{i+\frac{1}{2}} s_{i-1} L_{i-1} p_{i-\frac{1}{2}} p_{i+\frac{1}{2}} p_{i+\frac{3}{2}} \\
&= p_{i+\frac{3}{2}}p_{i+\frac{1}{2}},
\end{aligned}$$

as required. \square

The next statement establishes an alternative formula for the family $(L_{i+\frac{1}{2}} : i = 0, 1, \dots)$, for use in §5 and [En].

Theorem 4.4. *If $i = 1, 2, \dots$, then*

$$L_{i+\frac{1}{2}} = -L_i p_{i+\frac{1}{2}} - p_{i+\frac{1}{2}} L_i + (z - L_{i-\frac{1}{2}})p_{i+\frac{1}{2}} + s_i L_{i-\frac{1}{2}} s_i + \sigma_{i+\frac{1}{2}}.$$

Proof. By Proposition 3.1 and Proposition 4.3,

$$p_{i+\frac{1}{2}} L_i p_i p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}} p_{i+1} \sigma_{i+1} p_i p_{i+\frac{1}{2}} = p_{i+\frac{1}{2}} p_{i+1} \sigma_{i+\frac{1}{2}} p_{i+1} p_{i+\frac{1}{2}} = (z - L_{i-\frac{1}{2}})p_{i+\frac{1}{2}}.$$

Substituting the above expression into the definition of $L_{i+\frac{1}{2}}$ given in (3.3) provides the required statement. \square

5. SCHUR–WEYL DUALITY

This section uses Schur–Weyl duality to establish that the family $(L_{i+\frac{1}{2}}, L_{i+1} : i = 1, 2, \dots)$ defined above and the Jucys–Murphy elements given by [HR] are in fact equal.

Let $n = 1, 2, \dots$, and V be a vector space over \mathbb{C} with basis v_1, \dots, v_n . If $r = 1, 2, \dots$, the tensor product

$$V^{\otimes r} = \underbrace{V \otimes V \otimes \cdots \otimes V}_{r \text{ factors}} \quad \text{has basis} \quad \{v_{i_1} \otimes v_{i_2} \otimes \cdots \otimes v_{i_r} \mid 1 \leq i_1, \dots, i_r \leq n\},$$

and is equipped, via the inclusion $\mathfrak{S}_n \subset GL_n(\mathbb{C})$, with the diagonal \mathfrak{S}_n -action

$$w(v_{i_1} \otimes v_{i_2} \otimes \cdots \otimes v_{i_r})w = v_{(i_1)w^{-1}} \otimes v_{(i_2)w^{-1}} \otimes \cdots \otimes v_{(i_r)w^{-1}}, \quad \text{for } w \in \mathfrak{S}_n.$$

Let $A_r(n) = \mathcal{A}_r(z) \otimes_{\mathbb{Z}[z]} \mathbb{C}$, where z acts on \mathbb{C} as multiplication by n . The action of $A_r(n)$ on $V^{\otimes r}$ is given (§3 of [HR]) by

$$u(v_{i_1} \otimes v_{i_2} \otimes \cdots \otimes v_{i_r}) = v_{i_{(1)u^{-1}}} \otimes v_{i_{(2)u^{-1}}} \otimes \cdots \otimes v_{i_{(r)u^{-1}}}, \quad \text{for } u \in \mathfrak{S}_r,$$

and for $k = 1, \dots, r-1$,

$$p_{k+\frac{1}{2}}(v_{i_1} \otimes v_{i_2} \otimes \cdots \otimes v_{i_r}) = \begin{cases} v_{i_1} \otimes v_{i_2} \otimes \cdots \otimes v_{i_r} & \text{if } i_k = i_{k+1}, \\ 0 & \text{otherwise,} \end{cases}$$

and for $k = 1, \dots, r$,

$$p_k(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}} \otimes v_{i_k} \otimes v_{i_{k+1}} \otimes \cdots \otimes v_{i_r}) = \sum_{j=1}^n v_{i_1} \otimes \cdots \otimes v_{i_{k-1}} \otimes v_j \otimes v_{i_{k+1}} \otimes \cdots \otimes v_{i_r}.$$

The $A_{r+\frac{1}{2}}(n)$ -action on $V^{\otimes r}$ is obtained in §3 of [HR] from the action of $A_{r+1}(n)$ on $V^{\otimes r+1}$ by restricting to the subspace $V^{\otimes r} \otimes v_n$ and identifying $V^{\otimes r}$ with $V^{\otimes r} \otimes v_n$. The next statement asserts that \mathfrak{S}_n and $A_r(n)$ act as commuting operators on $V^{\otimes r}$.

Theorem 5.1 (Theorem 3.22 of [HR]). *Let $n, r \in \mathbb{Z}_{\geq 0}$. Let S_n^λ denote the irreducible \mathfrak{S}_n -module indexed by λ .*

(1) *As $(\mathbb{C}\mathfrak{S}_n, A_r(n))$ -bimodules*

$$V^{\otimes r} \cong \bigoplus_{\lambda \in \hat{A}_r(n)} S_n^\lambda \otimes A_r^\lambda(n),$$

where $\lambda \in \hat{A}_r(n)$ is an indexing set for the irreducible $A_r^\lambda(n)$ -modules, and the vector spaces $A_r^\lambda(n)$, for $\lambda \in \hat{A}_r(n)$, are irreducible $A_r^\lambda(n)$ -modules.

(2) *As $(\mathbb{C}\mathfrak{S}_{n-1}, A_{r+\frac{1}{2}}(n))$ -bimodules*

$$V^{\otimes r} \cong \bigoplus_{\lambda \in \hat{A}_{r+\frac{1}{2}}(n)} S_{n-1}^\lambda \otimes A_{r+\frac{1}{2}}^\lambda(n),$$

where $\lambda \in \hat{A}_{r+\frac{1}{2}}(n)$ is an indexing set for the irreducible $A_{r+\frac{1}{2}}^\lambda(n)$ -modules, and the vector spaces $A_{r+\frac{1}{2}}^\lambda(n)$, for $\lambda \in \hat{A}_{r+\frac{1}{2}}(n)$, are irreducible $A_{r+\frac{1}{2}}^\lambda(n)$ -modules.

By Theorem 3.6 of [HR], the homomorphism $A_r(n) \rightarrow \text{End}_{\mathfrak{S}_n}(V^{\otimes r})$ in Theorem 5.1 is an isomorphism whenever $n \geq 2r$.

If $1 \leq i, j \leq n$, let $s_{i,j} \in \mathfrak{S}_n$ denote the transposition which interchanges i and j . The next statement gives the action of the group $\langle \sigma_{i+\frac{1}{2}}, \sigma_{i+1} \mid i = 1, \dots, r-1 \rangle$ on $V^{\otimes r}$.

Proposition 5.2. If $k = 1, 2, \dots$, and $v_{i_1} \otimes \cdots \otimes v_{i_{k+1}} \in V^{\otimes k+1}$, then

$$(5.1) \quad \sigma_{k+\frac{1}{2}}(v_{i_1} \otimes \cdots \otimes v_{i_{k+1}}) = \underbrace{s_{i_k, i_{k+1}} \otimes \cdots \otimes s_{i_k, i_{k+1}}}_{k-1 \text{ factors}}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_{i_k} \otimes v_{i_{k+1}}$$

and

$$(5.2) \quad \sigma_{k+1}(v_{i_1} \otimes \cdots \otimes v_{i_{k+1}}) = \underbrace{s_{i_k, i_{k+1}} \otimes \cdots \otimes s_{i_k, i_{k+1}}}_{k+1 \text{ factors}}(v_{i_1} \otimes \cdots \otimes v_{i_k} \otimes v_{i_{k+1}}).$$

Proof. The proposition is true when $k = 1$. If $k = 2, 3, \dots$, observe that the linear endomorphisms defined, for $v_{i_1} \otimes \cdots \otimes v_{i_{k+1}} \in V^{\otimes k+1}$, by

$$\theta_{k+\frac{1}{2}} : v_{i_1} \otimes \cdots \otimes v_{i_{k+1}} \mapsto \underbrace{s_{i_k, i_{k+1}} \otimes \cdots \otimes s_{i_k, i_{k+1}}}_{k-1 \text{ factors}}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_{i_k} \otimes v_{i_{k+1}}$$

and

$$\theta_{k+1} : v_{i_1} \otimes \cdots \otimes v_{i_{k+1}} \mapsto \underbrace{s_{i_k, i_{k+1}} \otimes \cdots \otimes s_{i_k, i_{k+1}}}_{k+1 \text{ factors}}(v_{i_1} \otimes \cdots \otimes v_{i_k} \otimes v_{i_{k+1}})$$

commute with the diagonal action of \mathfrak{S}_n on $V^{\otimes k+1}$. Thus, by Theorem 5.1, $\theta_{k+\frac{1}{2}}$ and θ_{k+1} , lie in the image of the map $A_{k+1}(n) \mapsto \text{End}_{\mathfrak{S}_n}(V^{\otimes k+1})$. Observe also that the action of $\theta_{k+\frac{1}{2}}$ on $V^{\otimes k+1}$ commutes with the action of $A_{k-1}(n)$ on $V^{\otimes k+1}$, and the action of θ_{k+1} on $V^{\otimes k+1}$ commutes with the action of $A_{k-\frac{1}{2}}(n)$ on $V^{\otimes k+1}$. Since $A_{k+1}(n)$ is generated by $A_{k-\frac{1}{2}}(n)$ together with $\langle \sigma_{k+\frac{1}{2}}, \sigma_{k+1}, p_k, p_{k+\frac{1}{2}} \rangle$, to show that the map $A_{k+1}(n) \rightarrow \text{End}_{\mathfrak{S}_n}(V^{\otimes k+1})$ sends

$$\sigma_{k+\frac{1}{2}} \mapsto \theta_{k+\frac{1}{2}} \quad \text{and} \quad \sigma_{k+1} \mapsto \theta_{k+1},$$

it now suffices to verify that, as operators on $V^{\otimes k+1}$,

- (1) $\theta_{k+\frac{1}{2}}^2 = \theta_{k+1}^2 = 1$,
- (2) $\theta_{k+\frac{1}{2}} \theta_{k+1} = \theta_{k+1} \theta_{k+\frac{1}{2}} = s_k$,
- (3) $\theta_{k+\frac{1}{2}} p_{k+\frac{1}{2}} = p_{k+\frac{1}{2}} \theta_{k+\frac{1}{2}} = p_{k+\frac{1}{2}}$,
- (4) $\theta_{k+\frac{1}{2}} p_k p_{k+1} = \theta_{k+1} p_k p_{k+1}$,
- (5) $p_k p_{k+1} \theta_{k+\frac{1}{2}} = p_k p_{k+1} \theta_{k+1}$,
- (6) $\theta_{k+\frac{1}{2}} p_k \theta_{k+\frac{1}{2}} = \theta_{k+1} p_{k+1} \theta_{k+1}$,
- (7) $\theta_{k+\frac{1}{2}} p_{k-\frac{1}{2}} \theta_{k+\frac{1}{2}} = \sigma_k p_{k+\frac{1}{2}} \sigma_k$.

Since each of the above relations can be verified by inspection, the proof of the proposition is complete. \square

Proposition 5.3. If $k = 1, 2, \dots$, and $v_{i_1} \otimes \cdots \otimes v_{i_k} \in V^{\otimes k}$ then

$$(5.3) \quad L_{k-\frac{1}{2}}(v_{i_1} \otimes \cdots \otimes v_{i_k}) = nv_{i_1} \otimes \cdots \otimes v_{i_k} - \sum_{j=1}^n s_{i_k, j} \otimes \cdots \otimes s_{i_k, j}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_{i_k},$$

and

$$(5.4) \quad L_k(v_{i_1} \otimes \cdots \otimes v_{i_k}) = \sum_{j=1}^n s_{i_k, j} \otimes \cdots \otimes s_{i_k, j}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_j.$$

Proof. Identify $V^{\otimes k}$ as the subspace $V^{\otimes k} \otimes \sum_{j=1}^n v_j \subseteq V^{\otimes k+1}$. Then

$$\begin{aligned} p_{k+1}\sigma_{k+\frac{1}{2}}p_{i+1}(v_{i_1} \otimes \cdots \otimes v_{i_k} \otimes v_n) &= p_{k+1} \sum_{j=1}^n \sigma_{k+\frac{1}{2}}(v_{i_1} \otimes \cdots \otimes v_{i_k} \otimes v_j) \\ &= p_{k+1} \sum_{j=1}^n s_{i_k,j} \otimes \cdots \otimes s_{i_k,j}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_{i_k} \otimes v_j \\ &= \sum_{j,\ell=1}^n s_{i_k,j} \otimes \cdots \otimes s_{i_k,j}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_{i_k} \otimes v_\ell. \end{aligned}$$

By Proposition 4.3, as operators on $V^{\otimes k}$,

$$\begin{aligned} p_{k+1}\sigma_{k+\frac{1}{2}}p_{i+1}(v_{i_1} \otimes \cdots \otimes v_{i_k} \otimes v_n) &= (n - L_{i-\frac{1}{2}})p_{i+1}(v_{i_1} \otimes \cdots \otimes v_{i_k} \otimes v_n) \\ &= \sum_{\ell=1}^n (n - L_{i-\frac{1}{2}})(v_{i_1} \otimes \cdots \otimes v_{i_k} \otimes v_\ell), \end{aligned}$$

and the statement (5.3) follows. Next,

$$\begin{aligned} p_{k+1}\sigma_{k+1}p_{k+1}(v_{i_1} \otimes \cdots \otimes v_{i_k} \otimes v_n) &= p_{k+1} \sum_{j=1}^n \sigma_{k+1}(v_{i_1} \otimes \cdots \otimes v_{i_k} \otimes v_j) \\ &= p_{k+1} \sum_{j=1}^n \sigma_{k+\frac{1}{2}}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}} \otimes v_j \otimes v_{i_k}) \\ &= p_{k+1} \sum_{j=1}^n s_{i_k,i_{k+1}} \otimes \cdots \otimes s_{i_k,i_{k+1}}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}} \otimes v_j) \otimes v_{i_k} \\ &= \sum_{j,\ell=1}^n s_{i_k,i_{k+1}} \otimes \cdots \otimes s_{i_k,i_{k+1}}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}} \otimes v_j) \otimes v_\ell. \end{aligned}$$

By Proposition 4.3, as operators on $V^{\otimes k}$,

$$\begin{aligned} p_{k+1}\sigma_{k+1}p_{k+1}(v_{i_1} \otimes \cdots \otimes v_{i_k} \otimes v_n) &= L_k p_{k+1}(v_{i_1} \otimes \cdots \otimes v_{i_k} \otimes v_n) \\ &= \sum_{\ell=1}^n L_k(v_{i_1} \otimes \cdots \otimes v_{i_k} \otimes v_\ell), \end{aligned}$$

which yields (5.4). \square

As in §3 of [HR], let κ_n be the central element which is the class sum corresponding to the conjugacy class of transpositions in $\mathbb{C}\mathfrak{S}_n$,

$$\kappa_n = \sum_{1 \leq i < j \leq n} s_{i,j}.$$

For $\ell = 1, \dots, n$, we also define

$$\kappa_{n,\ell} = \sum_{\substack{1 \leq i < j \leq n \\ i,j \neq \ell}} s_{i,j},$$

so that $\kappa_{n,n} = \kappa_{n-1}$.

Proposition 5.4. Let $n = \dim(V)$ and $r \in \mathbb{Z}_{>0}$. If $z_{r-\frac{1}{2}} \in A_{r-\frac{1}{2}}(n)$ and $z_r \in A_r(n)$ are the central elements defined by Theorem 3.9, and $v_{i_1} \otimes \cdots \otimes v_{i_r} \in V^{\otimes r}$, then

$$(5.5) \quad z_r(v_{i_1} \otimes \cdots \otimes v_{i_r}) = \kappa_n(v_{i_1} \otimes \cdots \otimes v_{i_r}) - \left(\binom{n}{2} - rn\right)(v_{i_1} \otimes \cdots \otimes v_{i_r}),$$

and, if $r = 2, 3, \dots$, then,

$$(5.6) \quad z_{r-\frac{1}{2}}(v_{i_1} \otimes \cdots \otimes v_{i_r}) = \kappa_{n,i_r}(v_{i_1} \otimes \cdots \otimes v_{i_r}) - \left(\binom{n}{2} - rn + 1\right)(v_{i_1} \otimes \cdots \otimes v_{i_r}).$$

Proof. The proof is by induction on r . Let $i = 1, \dots, n$. Since $z_1 = p_1$,

$$\begin{aligned} \kappa_n v_i &= \sum_{i>j} s_{j,i} v_i + \sum_{j>i} s_{i,j} v_j + \sum_{\substack{j,\ell \\ j,\ell \neq i}} s_{j,\ell} v_i \\ &= \sum_{j=1}^n v_j + \left(\binom{n-1}{2} - 1\right) v_i \\ &= z_1 v_i + \left(\binom{n}{2} - n\right) v_i, \end{aligned}$$

which verifies (5.5) when $r = 1$. Now observe that if $v_{i_1} \otimes \cdots \otimes v_{i_k} \in V^{\otimes k}$, then the diagonal action of κ_n and κ_{n,i_r} on $V^{\otimes k}$ allows us to write

$$\kappa_n(v_{i_1} \otimes \cdots \otimes v_{i_k}) = \kappa_{n,i_k}(v_{i_1} \otimes \cdots \otimes v_{i_k}) + \sum_{\substack{j=1 \\ j \neq i_k}}^n s_{i_k,j} \otimes \cdots \otimes s_{i_k,j}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_j,$$

and

$$\begin{aligned} \kappa_{n,i_k}(v_{i_1} \otimes \cdots \otimes v_{i_k}) &= \kappa_n(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_{i_k} \\ &\quad - \sum_{\substack{j=1 \\ j \neq i_k}}^n s_{i_k,j} \otimes \cdots \otimes s_{i_k,j}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_{i_k}. \end{aligned}$$

Assuming that (5.5) holds for $r = 1, \dots, k-1$, we obtain

$$\begin{aligned} z_{k-\frac{1}{2}}(v_{i_1} \otimes \cdots \otimes v_{i_k}) &= z_{k-1}(v_{i_1} \otimes \cdots \otimes v_{i_k}) + L_{k-\frac{1}{2}}(v_{i_1} \otimes \cdots \otimes v_{i_k}) \\ &= \kappa_n(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_{i_k} - \left(\binom{n}{2} - (k-1)n\right)(v_{i_1} \otimes \cdots \otimes v_{i_k}) \\ &\quad + nv_{i_1} \otimes \cdots \otimes v_{i_k} - \sum_{j=1}^n s_{i_k,j} \otimes \cdots \otimes s_{i_k,j}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_{i_k} \\ &= \kappa_n(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_{i_k} - \left(\binom{n}{2} - kn + 1\right)(v_{i_1} \otimes \cdots \otimes v_{i_k}) \\ &\quad - \sum_{\substack{j=1 \\ j \neq i_k}}^n s_{i_k,j} \otimes \cdots \otimes s_{i_k,j}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_{i_k} \\ &= \kappa_{n,i_k}(v_{i_1} \otimes \cdots \otimes v_{i_k}) - \left(\binom{n}{2} - kn + 1\right)(v_{i_1} \otimes \cdots \otimes v_{i_k}), \end{aligned}$$

which verifies (5.6) for $r = k$. Next, since (5.6) holds for $r = 2, 3, \dots, k$, we obtain

$$\begin{aligned}
z_k(v_{i_1} \otimes \cdots \otimes v_{i_k}) &= z_{k-\frac{1}{2}}(v_{i_1} \otimes \cdots \otimes v_{i_k}) + L_k(v_{i_1} \otimes \cdots \otimes v_{i_k}) \\
&= \kappa_{n,i_k}(v_{i_1} \otimes \cdots \otimes v_{i_k}) - \binom{n}{2}(kn+1)(v_{i_1} \otimes \cdots \otimes v_{i_k}) \\
&\quad + \sum_{j=1}^n s_{i_k,j} \otimes \cdots \otimes s_{i_k,j}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_j \\
&= \kappa_{n,i_k}(v_{i_1} \otimes \cdots \otimes v_{i_k}) - \binom{n}{2}(kn)(v_{i_1} \otimes \cdots \otimes v_{i_k}) \\
&\quad + \sum_{\substack{j=1 \\ j \neq i_k}}^n s_{i_k,j} \otimes \cdots \otimes s_{i_k,j}(v_{i_1} \otimes \cdots \otimes v_{i_{k-1}}) \otimes v_j \\
&= \kappa_n(v_{i_1} \otimes \cdots \otimes v_{i_k}) - \binom{n}{2}(kn)(v_{i_1} \otimes \cdots \otimes v_{i_k}),
\end{aligned}$$

which verifies (5.5) for $r = k$. \square

Let $Z_k \in A_k(n)$ and $Z_{k+\frac{1}{2}} \in A_{k+\frac{1}{2}}(n)$ denote the central element defined by Halverson and Ram in §3 of [HR]. Then the Jucys–Murphy elements of [HR] are given by

$$M_{k-\frac{1}{2}} = Z_{k-\frac{1}{2}} - Z_{k-1} \quad \text{and} \quad M_k = Z_k - Z_{k-\frac{1}{2}} \quad \text{for } k = 1, 2, \dots$$

Theorem 5.5. if $k = 1, 2, \dots$, then $M_{k+\frac{1}{2}} = L_{k+\frac{1}{2}}$ and $M_{k+1} = L_{k+1}$ as elements of $A_{k+1}(n)$.

Proof. By Theorem 3.6 of [HR], the homomorphism $A_k(n) \rightarrow \text{End}_{\mathfrak{S}_n}(V^{\otimes k})$ is an isomorphism whenever $n \geq 2k$. Since the coefficients in the expansions of $Z_k, Z_{k+\frac{1}{2}}$ and $z_k, z_{k+\frac{1}{2}}$, in terms of the basis of diagrams for $A_{k+1}(n)$, are polynomials in n , and the map $A_k(n) \rightarrow \text{End}_{\mathfrak{S}_n}(V^{\otimes k})$ is an isomorphism for infinitely many values of n , to prove the theorem, it suffices to compare the action of z_k and Z_k (*resp.* $z_{k+\frac{1}{2}}$ and $Z_{k+\frac{1}{2}}$) on $V^{\otimes k}$ for an arbitrary choice of n . Identifying $V^{\otimes k}$ with the subspace $V^{\otimes k} \otimes v_n \subseteq V^{\otimes k+1}$, and κ_{n-1} with $\kappa_{n,n}$, then Theorem 3.35 of [HR] states that, as operators on $V^{\otimes k}$,

$$(5.7) \quad Z_k = \kappa_n - \binom{n}{2}(kn) \quad \text{and} \quad Z_{k+\frac{1}{2}} = \kappa_{n-1} - \binom{n}{2}((k+1)n+1).$$

Comparing (5.7) with the action of z_k and $z_{k+\frac{1}{2}}$ on $V^{\otimes k}$ in (5.5) and (5.6) completes the proof. \square

Remark 5.6. Whereas $L_{\frac{1}{2}} = 0$ and $L_1 = p_1$, in [HR], the first three Jucys–Murphy elements are $M_0 = M_{\frac{1}{2}} = 1$, and $M_1 = p_1 - 1$. Thus, although $z_1 = Z_1$ as elements of $A_1(n)$, Theorem 5.5 cannot be extended to the case $k = 0$.

REFERENCES

- [En] J. Enyang, *A seminormal form for partition algebras*, in preparation.
- [GG] F.M. Goodman and J. Gruber, *Cellularity and the Jones basic construction*, Adv. in App. Math. (to appear).
- [GG1] F.M. Goodman and J. Gruber, *On cellular algebras with Jucys–Murphy elements*, preprint, 2009.
- [HR] T. Halverson and A. Ram, *Partition algebras*, European J. Combin. **26** (2005), 869–921.
- [Jo] V.F.R Jones, *The Potts model and the symmetric group*, in: Subfactors: Proceedings of the Taniguchi Symposium on Operator Algebras (Kuzeso, 1993), River Edge, NJ, World Sci. Publishing, 1994.
- [Ma] P. Martin, *Potts models and related problems in statistical mechanics*, Series on Advances in Statistical Mechanics, vol. 5, World Scientific Publishing Co. Inc., Teaneck NJ, 1991.
- [MS] A. Mathas and M. Soriano, *Seminormal forms and Gram determinants for cellular algebras*, J. Reine Angew. Math., **619** (2008), 141–173. With an appendix by M. Soriano.
- [Na] M. Nazarov, *Young’s orthogonal form for Brauer’s centralizer algebra*, J. Algebra **182** (1996), 664–693.
- [Ow] W. Owens, *The Partition Algebra*, Honors Thesis, Macalester College, 2002 (May).
- [Xi] C.C. Xi, *Partition algebras are cellular*, Comp. Math. **119** (1999), 99–109.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, PARKVILLE, VIC 3010 AUSTRALIA

E-mail address: jenyang@unimelb.edu.au