

A Note on Intuitionistic Fuzzy Hypervector Spaces

Sanjay Roy, T. K. Samanta

Department of Mathematics, South Bantra Ramkrishna Institution , West Bengal, India.

e-mail: sanjaypuremath@gmail.com

Department of Mathematics, Uluberia College, West Bengal, India.

e-mail: mumpu_tapas5@yahoo.co.in

Abstract

The notion of Intuitionistic fuzzy hypervector space has been generalized and a few basic properties on this concept are studied. It has been shown that the intersection and union of an arbitrary family of Intuitionistic fuzzy hypervector spaces are also Intuitionistic fuzzy hypervector space. Lastly, the notion of a linear transformation on a hypervector space is introduced and established an important theorem relative to Intuitionistic fuzzy hypervector spaces.

Keywords: *Intuitionistic fuzzy hyperfield, Intuitionistic fuzzy hypervector spaces, Linear transformation.*

2010 Mathematics Subject Classification: 03F55, 08A72

1 Introduction

The notion of hyperstructure was introduced by F. Marty in 1934. Then he established the definition of hypergroup [4] in 1935. Since then many researchers have studied and developed (for example see [5] , [6]) the concept of different types of hyperstructures in different views. In 1990 M. S. Tallini

[10] introduced the notion of hypervector spaces. Then in 2005 R. Ameri [1] also studied this spaces extensively. In our previous papers ([7], [8]), we also introduced the notion of a hypervector spaces in more general form than the previous concept of hypervector space and thereafter established a few useful theorems in this space.

The concept of intuitionistic Fuzzy set, as a generalization of a fuzzy set was first introduced by Atanassov [3]. Then many researchers ([2], [9], [12]) applied this notion to norm, Continuity and Uniform Convergence etc. At the present time many researchers (for example [11]) are trying to apply this concept on the hyperstructure theory.

In this paper, the concept of Intuitionistic fuzzy hypervector space is introduced and a few basic properties are developed. Further it has been shown that the intersection and union of a arbitrary family of Intuitionistic fuzzy hypervector spaces are also Intuitionistic fuzzy hypervector space. Lastly we have introduced the notion of a linear transformation on a hypervector space and established an important theorem relative to Intuitionistic fuzzy hypervector space.

2 Preliminaries

This section contain some basic definition and preliminary results which will be needed.

Definition 2.1 [6] A **hyperoperation** over a non empty set X is a mapping of $X \times X$ into the set of all non empty subsets of X .

Definition 2.2 [6] A non empty set R with exactly one hyperoperation ' $\#$ ' is a **hypergroupoid**.

Let $(X, \#)$ be a hypergroupoid. For every point $x \in X$ and every non empty

subset A of X , we defined $x \# A = \bigcup_{a \in A} \{x \# a\}$.

Definition 2.3 [6] A hypergroupoid $(X, \#)$ is called a **hypergroup** if

- (i) $x \# (y \# z) = (x \# y) \# z$
- (ii) $\exists 0 \in X$ such that for every $a \in X$, there is unique element $b \in X$ for which $0 \in a \# b$ and $0 \in b \# a$. Here b is denoted by $-a$.
- (iii) For all $a, b, c \in X$ if $a \in b \# c$, then $b \in a \# (-c)$.

Proposition 2.4 [6] (i) In a hypergroup $(X, \#)$, $-(-a) = a$, $\forall a \in X$.

(ii) $0 \# a = \{a\}$, $\forall a \in X$, if $(X, \#)$ is a commutative hypergroup.

(iii) In a commutative hypergroup $(X, \#)$, 0 is unique.

Definition 2.5 [7] A **hyperring** is a non empty set equipped with a hyperaddition ' $\#$ ' and a multiplication ' \cdot ' such that $(X, \#)$ is a commutative hypergroup and (X, \cdot) is a semigroup and the multiplication is distributive across the hyperaddition both from the left and from the right and $a.0 = 0.a = 0$, $\forall a \in X$, where 0 is the zero element of the hyperring.

Definition 2.6 [7] A **hyperfield** is a non empty set X equipped with a hyperaddition ' $\#$ ' and a multiplication ' \cdot ' such that

- (i) $(X, \#, \cdot)$ is a hyperring.
- (ii) \exists an element $1 \in X$, called the identity element such that $a.1 = a$, $\forall a \in X$
- (iii) For each non zero element a in X , \exists an element a^{-1} such that $a.a^{-1}=1$
- (iv) $a.b = b.a$, $\forall a, b \in X$.

Definition 2.7 [8] Let (F, \oplus, \cdot) be a hyperfield and $(V, \#)$ be an additive commutative hypergroup. Then V is said to be a **hypervector space** over the hyperfield F if there exist a hyperoperation $* : F \times V \rightarrow P^*(V)$ such that

- (i) $a * (\alpha \# \beta) \subseteq a * \alpha \# a * \beta$, $\forall a \in F$ and $\forall \alpha, \beta \in V$

- (ii) $(a \oplus b) * \alpha \subseteq a * \alpha \# b * \alpha , \quad \forall a, b \in F \text{ and } \forall \alpha \in V$
- (iii) $(a . b) * \alpha = a * (b * \alpha) , \quad \forall a, b \in F \text{ and } \forall \alpha \in V.$
- (iv) $(-a) * \alpha = a * (-\alpha), \quad \forall \alpha \in V \text{ and } \forall a \in F.$
- (v) $\alpha \in 1_F * \alpha, \theta \in 0 * \alpha \text{ and } 0 * \theta = \theta, \quad \forall \alpha \in V.$

where 1_F is the identity element of F , 0 is the zero element of F and θ is zero vector of V and $P^*(V)$ is the set of all non empty subset of V .

Definition 2.8 [1] Let $f : X \rightarrow Y$ be a mapping and $\nu \in FS(Y)$. Then we define $f^{-1}(\nu) \in FS(X)$ as follows:

$$f^{-1}(\nu)(x) = \nu(f(x)), \forall x \in X.$$

Definition 2.9 [12] Let E be a any set. An Intuitionistic fuzzy set (IFS) A of E is an object of the form $A = \{ (x, \mu_A(x), \nu_A(x)) : x \in E \}$, where the functions $\mu_A : E \rightarrow [0, 1]$ and $\nu_A : E \rightarrow [0, 1]$ denotes the degree of membership and the non-membership of the element $x \in E$ respectively and for every $x \in E$, $0 \leq \mu_A(x) + \nu_A(x) \leq 1$.

3 Intuitionistic Fuzzy hypervector Space

In this section we established the definition of intuitionistic fuzzy hypervector Spaces and deduce some important theorems.

Definition 3.1 Let $(F, \oplus, .)$ be a hyperfield. An **intuitionistic fuzzy hypervector field** on F is an object of the form $A = \{ (a, \mu_F(a), \nu_F(a)) : a \in F \}$ satisfies the following conditions :

- (i) $\wedge_{x \in a \oplus b} \mu_F(x) \geq \mu_F(a) \wedge \mu_F(b), \forall a, b \in F$
- (ii) $\mu_F(-a) \geq \mu_F(a), \forall a \in F$
- (iii) $\mu_F(a.b) \geq \mu_F(a) \wedge \mu_F(b) \forall a, b \in F$
- (iv) $\mu_F(a^{-1}) \geq \mu_F(a), \forall a (\neq 0) \in F$
- (v) $\vee_{x \in a \oplus b} \nu_F(x) \leq \nu_F(a) \vee \nu_F(b), \forall a, b \in F$

- (vi) $\nu_F(-a) \leq \nu_F(a), \forall a \in F$
- (vii) $\nu_F(a.b) \leq \nu_F(a) \vee \nu_F(b), \forall a, b \in F$
- (viii) $\nu_F(a^{-1}) \leq \nu_F(a), \forall a(\neq 0) \in F.$

Result 3.2 If A is a intuitionistic fuzzy hyperfield of F , then

- (i) $\mu_F(0) \geq \mu_F(a), \forall a \in F$
- (ii) $\mu_F(1) \geq \mu_F(a), \forall a \in F \setminus \{0\}$
- (iii) $\mu_F(0) \geq \mu_F(1)$
- (iv) $\nu_F(0) \leq \nu_F(a), \forall a \in F$
- (v) $\nu_F(1) \leq \nu_F(a), \forall a \in F \setminus \{0\}$
- (vi) $\nu_F(0) \leq \nu_F(1)$

Proof : Obvious.

Definition 3.3 Let $(V, \#, *)$ be a hypervector space over a hyperfield $(F, \oplus, .)$ and A be a intuitionistic fuzzy hyperfield in F . A intuitionistic fuzzy subset $B = \{(x, \mu_V(x), \nu_V(x)) : x \in V\}$ of V is said to be a **intuitionistic fuzzy hypervector space** of V over a intuitionistic fuzzy hyperfield A , if the following conditions are satisfied:

- (i) $\wedge_{\alpha \in x \# y} \mu_V(\alpha) \geq \mu_V(x) \wedge \mu_V(y), \forall x, y \in V$
- (ii) $\mu_V(-x) \geq \mu_V(x), \forall x \in V$
- (iii) $\wedge_{y \in a * x} \mu_V(y) \geq \mu_V(x) \wedge \mu_F(a), \forall a \in F \text{ and } \forall x \in V$
- (iv) $\mu_F(1) \geq \mu_V(\theta), \text{ where } \theta \text{ be the null vector of } V.$
- (v) $\vee_{\alpha \in x \# y} \nu_V(\alpha) \leq \nu_V(x) \vee \nu_V(y), \forall x, y \in V$
- (vi) $\nu_V(-x) \leq \nu_V(x), \forall x \in V$
- (vii) $\vee_{y \in a * x} \nu_V(y) \leq \nu_V(x) \vee \nu_F(a), \forall a \in F \text{ and } \forall x \in V$
- (viii) $\nu_F(1) \leq \nu_V(\theta), \text{ where } \theta \text{ be the null vector of } V.$

Here we say that B is a intuitionistic fuzzy hypervector space over a intuitionistic fuzzy hyperfield A .

Result 3.4 *If B is a intuitionistic fuzzy hypervector space over a intuitionistic fuzzy hyperfield A , then*

- (i) $\mu_F(0) \geq \mu_V(\theta)$,
- (ii) $\mu_V(\theta) \geq \mu_V(x), \forall x \in V$
- (iii) $\mu_F(0) \geq \mu_V(x), \forall x \in V$
- (iv) $\nu_F(0) \leq \nu_V(\theta)$,
- (v) $\nu_V(\theta) \leq \nu_V(x), \forall x \in V$
- (vi) $\nu_F(0) \leq \nu_V(x), \forall x \in V$

Proof : Obvious.

Theorem 3.5 *Let V be a hypervector space over a hyperfield F and A be a intuitionistic fuzzy hyperfield. Let $B \in IFS(V)$. Then B is a intuitionistic fuzzy hypervector space over A iff*

- (i) $\Lambda_{z \in a*x \# b*y} \mu_V(z) \geq (\mu_F(a) \wedge \mu_V(x)) \wedge (\mu_F(b) \wedge \mu_V(y)), \forall x, y \in V$ and
 $\forall a, b \in F$
- (ii) $\mu_F(1) \geq \mu_V(\theta)$ where θ be the null vector of V .
- (iii) $\vee_{z \in a*x \# b*y} \nu_V(z) \leq (\nu_F(a) \vee \nu_V(x)) \vee (\nu_F(b) \vee \nu_V(y)), \forall x, y \in V$ and
 $\forall a, b \in F$
- (iv) $\nu_F(1) \leq \nu_V(\theta)$ where θ be the null vector of V .

Proof: First we suppose that B is a intuitionistic fuzzy hypervector space over the intuitionistic fuzzy hyperfield A . Then for $a, b \in F$ and $x, y \in V$, we have

$$\begin{aligned}
\Lambda_{z \in a*x \# b*y} \mu_V(z) &= \Lambda_{z \in \alpha \# \beta, \alpha \in a*x, \beta \in b*y} \mu_V(z) \\
&= \Lambda_{\alpha \in a*x, \beta \in b*y} (\Lambda_{z \in \alpha \# \beta} \mu_V(z)) \\
&\geq \Lambda_{\alpha \in a*x, \beta \in b*y} (\mu_V(\alpha) \wedge \mu_V(\beta)) \\
&= (\Lambda_{\alpha \in a*x} \mu_V(\alpha)) \wedge (\Lambda_{\beta \in b*y} \mu_V(\beta)) \\
&\geq (\mu_F(a) \wedge \mu_V(x)) \wedge (\mu_F(b) \wedge \mu_V(y))
\end{aligned}$$

$$\begin{aligned}
\vee_{z \in a*x \# b*y} \nu_V(z) &= \vee_{z \in \alpha \# \beta, \alpha \in a*x, \beta \in b*y} \nu_V(z) \\
&= \vee_{\alpha \in a*x, \beta \in b*y} (\vee_{z \in \alpha \# \beta} \nu_V(z)) \\
&\leq \vee_{\alpha \in a*x, \beta \in b*y} (\nu_V(\alpha) \vee \nu_V(\beta)) \\
&= (\vee_{\alpha \in a*x} \nu_V(\alpha)) \vee (\vee_{\beta \in b*y} \nu_V(\beta)) \\
&\leq (\nu_F(a) \vee \nu_V(x)) \vee (\nu_F(b) \vee \nu_V(y))
\end{aligned}$$

The second and fourth inequalities are directly follow.

conversely suppose that the inequalities of the theorem hold for all $x, y \in V$ and $\forall a, b \in F$.

$$\begin{aligned}
\text{Then } \wedge_{z \in 1*x \# 1*y} \mu_V(z) &\geq (\mu_F(1) \wedge \mu_V(x)) \wedge (\mu_F(1) \wedge \mu_V(y)) \\
&\geq (\mu_V(\theta) \wedge \mu_V(x)) \wedge (\mu_V(\theta) \wedge \mu_V(y)) \\
&= \mu_V(x) \wedge \mu_V(y)
\end{aligned}$$

i.e $\wedge_{z \in x \# y} \mu_V(z) \geq \mu_V(x) \wedge \mu_V(y)$, as $x \in 1*x$ and $y \in 1*y$

$$\begin{aligned}
\mu_V(-x) &= \wedge_{z \in -1*x} \mu_V(z), \text{ as } -x \in -1*x \\
&\geq \wedge_{z \in -1*x \# 0*x} \mu_V(z) \geq (\mu_F(-1) \wedge \mu_V(x)) \wedge (\mu_F(0) \wedge \mu_V(x)) \\
&\geq (\mu_F(1) \wedge \mu_V(x)) \wedge \mu_V(x) \\
&\geq (\mu_V(\theta) \wedge \mu_V(x)) \wedge \mu_V(x) \\
&= \mu_V(x) \wedge \mu_V(x) \\
&= \mu_V(x)
\end{aligned}$$

i.e $\mu_V(-x) \geq \mu_V(x)$

$$\begin{aligned}
\wedge_{y \in a*x} \mu_V(y) &\geq \wedge_{y \in a*x \# 0*x} \mu_V(y) \geq (\mu_F(a) \wedge \mu_V(x)) \wedge (\mu_F(0) \wedge \mu_V(x)) \\
&= (\mu_V(x) \wedge \mu_F(a)) \wedge \mu_V(x), \text{ as } \mu_F(0) \geq \mu_V(x) \\
&= \mu_V(x) \wedge \mu_F(a)
\end{aligned}$$

The fourth inequality of definition 3.3 is directly follows.

$$\begin{aligned}
\text{Next } \vee_{z \in 1*x \# 1*y} \nu_V(z) &\leq (\nu_F(1) \vee \nu_V(x)) \vee (\nu_F(1) \vee \nu_V(y)) \\
&= (\nu_V(\theta) \vee \nu_V(x)) \vee (\nu_V(\theta) \vee \nu_V(y)) \\
&= \nu_V(x) \vee \nu_V(y)
\end{aligned}$$

i.e $\vee_{z \in x \# y} \nu_V(z) \leq \nu_V(x) \vee \nu_V(y)$, as $x \in 1*x$ and $y \in 1*y$

$$\begin{aligned}
\nu_V(-x) &\leq \bigvee_{z \in -1*x} \nu_V(z), \text{ as } -x \in -1*x \\
&\leq \bigvee_{z \in -1*x \# 0*x} \nu_V(z) \\
&\leq (\nu_F(-1) \vee \nu_V(x)) \vee (\nu_F(0) \vee \nu_V(x)) \\
&\leq (\nu_F(1) \vee \nu_V(x)) \vee \nu_V(x) \\
&\leq (\nu_V(\theta) \vee \nu_V(x)) \vee \nu_V(x) \\
&= \nu_V(x) \vee \nu_V(x) \\
&= \nu_V(x)
\end{aligned}$$

i.e $\nu_V(-x) \leq \nu_V(x)$

$$\begin{aligned}
\bigvee_{y \in a*x} \nu_V(y) &\leq \bigvee_{y \in a*x \# 0*x} \nu_V(y) \\
&\leq (\nu_F(a) \vee \nu_V(x)) \vee (\nu_F(0) \vee \nu_V(x)) \\
&= (\nu_V(x) \vee \nu_F(a)) \vee \nu_V(x), \text{ as } \nu_F(0) \leq \nu_V(x) \\
&= \nu_V(x) \vee \nu_F(a)
\end{aligned}$$

The eighth inequality of definition 3.3 is obvious.

Therefore B is a intuitionistic fuzzy hypervector space over A.

This completes the proof.

Definition 3.6 Let $B_{(\alpha \in \Lambda)}^\alpha = \{x, \mu_V^\alpha(x), \nu_V^\alpha(x) : x \in V\}$ be a family of intuitionistic fuzzy hypervector spaces of a hypervector space V over the same intuitionistic fuzzy hyperfield $A = \{(x, \mu_F(x), \nu_F(x)) : x \in F\}$. Then the **intersection** of those intuitionistic fuzzy hypervector spaces is defined as $(\bigcap_{\alpha \in \Lambda} B^\alpha)(x) = \{x, \bigwedge_{\alpha \in \Lambda} \mu_V^\alpha(x), \bigwedge_{\alpha \in \Lambda} \nu_V^\alpha(x) : \forall x \in V\}$

and the **union** of those intuitionistic fuzzy hypervector spaces is defined as

$$(\bigcup_{\alpha \in \Lambda} B^\alpha)(x) = \{x, \bigvee_{\alpha \in \Lambda} \mu_V^\alpha(x), \bigvee_{\alpha \in \Lambda} \nu_V^\alpha(x) : \forall x \in V\}$$

Theorem 3.7 The intersection of any family of intuitionistic fuzzy hypervector spaces of a hypervector space V is a intuitionistic fuzzy hypervector space.

Proof: Let $\{B^\alpha : \alpha \in \Lambda\}$ be a family of intuitionistic fuzzy hypervector spaces of V over the same intuitionistic fuzzy hyperfield $A = \{(x, \mu_F(x), \nu_F(x)) : x \in F\}$.

$$\text{Let } B = (\cap_{\alpha \in \Lambda} B^\alpha)(x) = \{(x, \mu_V(x), \nu_V(x)) : x \in V\}$$

$$\text{where } \mu_V(x) = \bigwedge_{\alpha \in \Lambda} \mu_V^\alpha(x) \text{ and } \nu_V(x) = \bigwedge_{\alpha \in \Lambda} \nu_V^\alpha(x)$$

Let $x, y \in V$ and $a, b \in F$

$$\begin{aligned} \bigwedge_{z \in a*x \# b*y} \mu_V(z) &= \bigwedge_{z \in a*x \# b*y} (\bigwedge_{\alpha \in \Lambda} \mu_V^\alpha(z)) \\ &= \bigwedge_{\alpha \in \Lambda} (\bigwedge_{z \in a*x \# b*y} \mu_V^\alpha(z)) \\ &\geq \bigwedge_{\alpha \in \Lambda} \{(\mu_F(a) \wedge \mu_V^\alpha(x)) \wedge (\mu_F(b) \wedge \mu_V^\alpha(y))\} \\ &= \{(\mu_F(a) \wedge (\bigwedge_{\alpha \in \Lambda} \mu_V^\alpha(x)))\} \wedge \{(\mu_F(b) \wedge (\bigwedge_{\alpha \in \Lambda} \mu_V^\alpha(y)))\} \\ &= (\mu_F(a) \wedge \mu_V(x)) \wedge (\mu_F(b) \wedge \mu_V(y)) \end{aligned}$$

$$\text{Therefore } \bigwedge_{z \in a*x \# b*y} \mu_V(z) \geq (\mu_F(a) \wedge \mu_V(x)) \wedge (\mu_F(b) \wedge \mu_V(y))$$

$$\text{Again } \mu_F(1) \geq \mu_V^\alpha(\theta), \forall \alpha \in \Lambda$$

$$\text{Therefore } \mu_F(1) \geq \bigwedge_{\alpha \in \Lambda} \mu_V^\alpha(\theta)$$

$$\text{i.e. } \mu_F(1) \geq \mu_V(\theta)$$

$$\begin{aligned} \text{Next } \bigvee_{z \in a*x \# b*y} \nu_V(z) &= \bigvee_{z \in a*x \# b*y} (\bigwedge_{\alpha \in \Lambda} \nu_V^\alpha(z)) \\ &\leq \bigwedge_{\alpha \in \Lambda} (\bigvee_{z \in a*x \# b*y} \nu_V^\alpha(z)) \\ &\leq \bigwedge_{\alpha \in \Lambda} \{(\nu_F(a) \vee \nu_V^\alpha(x)) \vee (\nu_F(b) \vee \nu_V^\alpha(y))\} \\ &= \{(\nu_F(a) \vee (\bigwedge_{\alpha \in \Lambda} \nu_V^\alpha(x)))\} \vee \{(\nu_F(b) \vee (\bigwedge_{\alpha \in \Lambda} \nu_V^\alpha(y)))\} \\ &= (\nu_F(a) \vee \nu_V(x)) \vee (\nu_F(b) \vee \nu_V(y)) \end{aligned}$$

$$\text{Therefore } \bigvee_{z \in a*x \# b*y} \nu_V(z) \leq (\nu_F(a) \vee \nu_V(x)) \vee (\nu_F(b) \vee \nu_V(y))$$

$$\text{Again } \nu_F(1) \leq \nu_V^\alpha(\theta), \forall \alpha \in \Lambda$$

$$\text{Therefore } \nu_F(1) \leq \bigwedge_{\alpha \in \Lambda} \nu_V^\alpha(\theta)$$

$$\text{i.e. } \nu_F(1) \leq \nu_V(\theta)$$

Therefore B is also a intuitionistic fuzzy hypervector space over A .

This completes the proof.

Theorem 3.8 *The union of any family of intuitionistic fuzzy hypervector spaces of a hypervector space V is a intuitionistic fuzzy hypervector space.*

Proof: Let $\{B^\alpha : \alpha \in \Lambda\}$ be a family of intuitionistic fuzzy hypervector spaces of V over the same intuitionistic fuzzy hyperfield $A = \{(x, \mu_F(x), \nu_F(x)) : x \in F\}$.

$$\text{Let } B = \bigvee_{\alpha \in \Lambda} \mu_V^\alpha(x) = \{(x, \mu_V(x), \nu_V(x)) : x \in V\}$$

$$\text{where } \mu_V(x) = \bigvee_{\alpha \in \Lambda} \mu_V^\alpha(x) \text{ and } \nu_V(x) = \bigvee_{\alpha \in \Lambda} \nu_V^\alpha(x)$$

Let $x, y \in V$ and $a, b \in F$

$$\begin{aligned} \wedge_{z \in a*x \# b*y} \mu_V(z) &= \wedge_{z \in a*x \# b*y} (\bigvee_{\alpha \in \Lambda} \mu_V^\alpha(z)) \\ &\geq \bigvee_{\alpha \in \Lambda} (\wedge_{z \in a*x \# b*y} \mu_V^\alpha(z)) \\ &\geq \bigvee_{\alpha \in \Lambda} \{(\mu_F(a) \wedge \mu_V^\alpha(x)) \wedge (\mu_F(b) \wedge \mu_V^\alpha(y))\} \\ &= \{(\mu_F(a) \wedge (\bigvee_{\alpha \in \Lambda} \mu_V^\alpha(x)))\} \wedge \{(\mu_F(b) \wedge (\bigvee_{\alpha \in \Lambda} \mu_V^\alpha(y)))\} \\ &= (\mu_F(a) \wedge \mu_V(x)) \wedge (\mu_F(b) \wedge \mu_V(y)) \end{aligned}$$

$$\text{Therefore } \wedge_{z \in a*x \# b*y} \mu_V(z) \geq (\mu_F(a) \wedge \mu_V(x)) \wedge (\mu_F(b) \wedge \mu_V(y))$$

$$\text{Again } \mu_F(1) \geq \mu_V^\alpha(\theta), \forall \alpha \in \Lambda$$

$$\text{Therefore } \mu_F(1) \geq \bigvee_{\alpha \in \Lambda} \mu_V^\alpha(\theta)$$

$$\text{i.e. } \mu_F(1) \geq \mu_V(\theta)$$

$$\begin{aligned} \text{Next } \bigvee_{z \in a*x \# b*y} \nu_V(z) &= \bigvee_{z \in a*x \# b*y} (\bigvee_{\alpha \in \Lambda} \nu_V^\alpha(z)) \\ &= \bigvee_{\alpha \in \Lambda} (\bigvee_{z \in a*x \# b*y} \nu_V^\alpha(z)) \\ &\leq \bigvee_{\alpha \in \Lambda} \{(\nu_F(a) \vee \nu_V^\alpha(x)) \vee (\nu_F(b) \vee \nu_V^\alpha(y))\} \\ &= \{(\nu_F(a) \vee (\bigvee_{\alpha \in \Lambda} \nu_V^\alpha(x)))\} \vee \{(\nu_F(b) \vee (\bigvee_{\alpha \in \Lambda} \nu_V^\alpha(y)))\} \\ &= (\nu_F(a) \vee \nu_V(x)) \vee (\nu_F(b) \vee \nu_V(y)) \end{aligned}$$

$$\text{Therefore } \bigvee_{z \in a*x \# b*y} \nu_V(z) \leq (\nu_F(a) \vee \nu_V(x)) \vee (\nu_F(b) \vee \nu_V(y))$$

$$\text{Again } \nu_F(1) \leq \nu_V^\alpha(\theta), \forall \alpha \in \Lambda$$

$$\text{Therefore } \nu_F(1) \leq \bigvee_{\alpha \in \Lambda} \nu_V^\alpha(\theta)$$

$$\text{i.e. } \nu_F(1) \leq \nu_V(\theta)$$

Therefore B is also a intuitionistic fuzzy hypervector space over A.

This completes the proof.

4 Linear Transformation

In this section we established the definition of a Linear transformation on hypervector Spaces and deduce a important theorem relative to a intuitionistic fuzzy concept.

Definition 4.1 Let $(V, \#, *)$ and $(W, \#', *')$ be two hypervector space over the same hyperfield $(F, \oplus, .)$. A mapping $T : V \rightarrow W$ is called Linear transformation iff

- (i) $T(x \# y) \subseteq T(x) \#' T(y)$,
- (ii) $T(a * x) \subseteq a *' T(x)$, $\forall x, y \in V$ and $a \in F$
- (iii) $T(\theta) = \theta'$

Theorem 4.2 Let $(V, \#, *)$ and $(W, \#', *')$ be two hypervector space over the same hyperfield $(F, \oplus, .)$ and $T : V \rightarrow W$ be a linear transformation. Let $B = \{(y, \mu_W(y), \nu_W(y)) : y \in W\}$ be a intuitionistic fuzzy hypervector space over $A = \{(a, \mu_F(a), \nu_F(a)) : a \in F\}$. Then $T^{-1}(B) = \{x, T^{-1}(\mu_W)(x), T^{-1}(\nu_W)(x) : x \in V\}$ is a intuitionistic fuzzy hypervector space of V over A .

Proof: Let $a, b \in F$ and $\alpha, \beta \in V$.

$$\begin{aligned}
 \text{Then } \bigwedge_{x \in a * \alpha \# b * \beta} T^{-1}(\mu_W)(x) &= \bigwedge_{x \in a * \alpha \# b * \beta} \mu_W(T(x)) \\
 &= \bigwedge_{T(x) \in T(a * \alpha \# b * \beta)} \mu_W(T(x)) \\
 &\geq \bigwedge_{T(x) \in a *' T(\alpha) \#' b *' T(\beta)} \mu_W(T(x)) \\
 &\geq (\mu_F(a) \wedge \mu_W(T(\alpha)) \wedge (\mu_F(b) \wedge \mu_W(T(\beta)))) \\
 &= (\mu_F(a) \wedge T^{-1}(\mu_W)(\alpha)) \wedge (\mu_F(b) \wedge T^{-1}(\mu_W)(\beta))
 \end{aligned}$$

Again $T^{-1}(\mu_W)(\theta) = \mu_W(T(\theta)) = \mu_W(\theta')$, where θ' be a null vector of W.

$$\leq \mu_F(1)$$

i.e $\mu_F(1) \geq T^{-1}(\mu_W)(\theta)$

$$\begin{aligned} \text{And } \bigvee_{x \in a * \alpha \# b * \beta} T^{-1}(\nu_W)(x) &= \bigvee_{x \in a * \alpha \# b * \beta} \nu_W(T(x)) \\ &= \bigvee_{T(x) \in T(a * \alpha \# b * \beta)} \nu_W(T(x)) \\ &\leq \bigvee_{T(x) \in a' * T(\alpha) \# b' * T(\beta)} \nu_W(T(x)) \\ &\leq (\nu_F(a) \vee \nu_W(T(\alpha))) \vee (\nu_F(b) \vee \nu_W(T(\beta))) \\ &= (\nu_F(a) \vee T^{-1}(\nu_W)(\alpha)) \vee (\nu_F(b) \vee T^{-1}(\nu_W)(\beta)) \end{aligned}$$

Again $T^{-1}(\nu_W)(\theta) = \nu_W(T(\theta)) = \nu_W(\theta')$, where θ' be a null vector of W.

$$\geq \nu_F(1)$$

i.e $\nu_F(1) \leq T^{-1}(\nu_W)(\theta)$

Therefore $T^{-1}(B)$ is a intuitionistic fuzzy hypervector space of a hypervector space V over A.

References

- [1] AMERI R. *Fuzzy Hypervector Spaces Over Valued Field*, Iranian J. of Fuzzy Systems Vol. 2, No. 1, (2005)pp. 37-47.
- [2] Dinda B, Samanta T. K. *Intuitionistic Fuzzy Continuity and Uniform Convergence*, Int. J. Open Problems Compt.Math., Vol 3, No.1 (2010) 8 - 26.
- [3] K. Atanassov *Intuitionistic fuzzy sets*, Fuzzy Sets and Systems 20 (1986) 87 - 96.
- [4] MARTY F. *Role de la notion de hypergroupe dans l'etude de groupes non abeliens*, comptes Renclus Acad. Sci. Paris, 201, 636-638,(1935).

- [5] MARTY F. *Sur les groupes et les hypergroupes*, attaches a une fraction rationnelle, Ann. Sci. Ecole Norm. Sup. (3) 53, 82-123, (1936).
- [6] Nakassis A. *Expository and survey article recent results in hyperring and hyperfield theory*, internet. J. Math. and Math. Sci.,11(2)(1988)209 - 220.
- [7] Roy S, Samanta T. K. *A note on Hypervector Spaces* ,(communicated)
- [8] Roy S, Samanta T. K. *Hyper innerproduct Spaces* ,(communicated)
- [9] Samanta T. K, Jebril IH. *Finite dimentional intuitionistic fuzzy normed linear space*, Int. J. Open Problems Compt. Math., Vol 2, No. 4 (2009) 574-591.
- [10] Tallini M S. *Hypervector Spaces*, Proceedings of fourth Int. Congress in Algebraic Hyperstructures and Applications, Xanthi,Greece, World Scientific(1990)167 - 174.
- [11] Torkzadeh, L. - Abbasi, M. - Zahedi, M.M. *Some results of intuitionistic fuzzy weak dual hyper K-ideals*, Iranian J. of Fuzzy Systems, Vol. 5, No. 1., (2008), 65-78.
- [12] Vijayabalaji S, Thillaigovindan N, Jun YB. *Intuitionistic Fuzzy n-normed linear space* , Bull. Korean Math. Soc. 44 (2007) 291 - 308.