

SPECTRAL PROPERTIES OF GRAIN BOUNDARIES AT SMALL ANGLES OF ROTATION

RAINER HEMPEL, MARTIN KOHLMANN

ABSTRACT. We study some spectral properties of a simple two-dimensional model for small angle defects in crystals and alloys. Starting from a periodic potential $V: \mathbb{R}^2 \rightarrow \mathbb{R}$, we let $V_\vartheta(x, y) = V(x, y)$ in the right half-plane $\{x \geq 0\}$ and $V_\vartheta = V \circ M_{-\vartheta}$ in the left half-plane $\{x < 0\}$, where $M_\vartheta \in \mathbb{R}^{2 \times 2}$ is the usual matrix describing rotation of the coordinates in \mathbb{R}^2 by an angle ϑ . As a main result, it is shown that spectral gaps of the periodic Schrödinger operator $H_0 = -\Delta + V$ fill with spectrum of $R_\vartheta = -\Delta + V_\vartheta$ as $0 \neq \vartheta \rightarrow 0$. Moreover, we obtain upper and lower bounds for a quantity pertaining to an integrated density of states measure for the surface states.

CONTENTS

Introduction	1
1. The dislocation problem on a strip and for the plane	5
2. The rotation problem for small angles	6
3. Integrated density of states bounds	8
4. Muffin tin potentials	12
5. Some extensions and remarks	15
6. Appendix	16
References	19

INTRODUCTION

In the quantum theory of solids one first studies periodic structures which can often be modelled by Schrödinger operators with periodic potentials. Other models deal with situations where periodicity holds only in subsets of the sample; more precisely, the sample is the disjoint union of subsets such that, in each subset, the potential is obtained by restricting different periodic potentials to the corresponding subsets. Such zones or “grains” occur frequently in crystals and in alloys; some typical examples are shown in Figure 1. It is an important issue to understand how the interface between two grains will influence the energy spectrum of the sample. Typically, the grain boundaries appear to be (piecewise) linear, and one is led to

2000 *Mathematics Subject Classification.* Primary 35J10, 35P20, 81Q10.

Key words and phrases. Schrödinger operators, eigenvalues, spectral gaps.

study problems on \mathbb{R}^2 with a potential $W = W(x, y)$ defined by

$$W(x, y) := \begin{cases} V_r(x, y), & x \geq 0, \\ V_\ell(x, y), & x < 0, \end{cases} \quad (0.1)$$

where $V_r, V_\ell: \mathbb{R}^2 \rightarrow \mathbb{R}$ are periodic. In many situations, V_ℓ is obtained from V_r by a translation or a rotation about the origin.




Figure 1: Edge dislocation and small angle grain boundary. The pictures on the right were taken with a TEM; references:

http://www.fys.uio.no/bate/?page_\,id=7

<http://pruffle.mit.edu/~ccarter/NANOAM/images/>

In this paper, we will use some results on a translational problem to obtain spectral information about rotational problems in the limit of small angles. Our main theorem deals with the following situation. Let $V: \mathbb{R}^2 \rightarrow \mathbb{R}$ be a Lipschitz-continuous function which is periodic w.r.t. the lattice \mathbb{Z}^2 . For $\vartheta \in (0, \pi/2)$, let

$$M_\vartheta := \begin{pmatrix} \cos \vartheta & -\sin \vartheta \\ \sin \vartheta & \cos \vartheta \end{pmatrix} \in \mathbb{R}^{2 \times 2}, \quad (0.2)$$

and

$$V_{\vartheta}(x, y) := \begin{cases} V(x, y), & x \geq 0, \\ V(M_{-\vartheta}(x, y)) & x < 0. \end{cases} \quad (0.3)$$

We then let H_0 denote the (unique) self-adjoint extension of $-\Delta \upharpoonright C_c^\infty(\mathbb{R}^2)$, acting in the Hilbert space $L_2(\mathbb{R}^2)$, and

$$R_{\vartheta} := H_0 + V_{\vartheta}, \quad D(R_{\vartheta}) = D(H_0). \quad (0.4)$$

Then R_{ϑ} is essentially self-adjoint on $C_c^\infty(\mathbb{R}^2)$ and semi-bounded from below. Our main assumption is that the periodic Hamiltonian $H := H_0 + V = R_0$ has a gap (a, b) in the essential spectrum $\sigma_{\text{ess}}(H)$, i.e., we assume that there exist $a < b \in \mathbb{R}$ that satisfy $\inf \sigma_{\text{ess}}(H) < a$ and $(a, b) \cap \sigma(H) = \emptyset$; we do not need to assume that a, b are the actual gap edges. It is easy to see (using, e.g., [RS-I; Thm. VIII.25]) that the operators R_{ϑ} converge to R_{ϑ_0} in the strong resolvent sense as $\vartheta \rightarrow \vartheta_0 \in [0, \pi/2)$; in particular, R_{ϑ} converges to H in the strong resolvent sense as $\vartheta \rightarrow 0$. Recall that strong resolvent convergence implies upper semi-continuity of the spectrum while the spectrum may contract considerably when the limit is reached. In the present paper, we are dealing with a situation where the spectrum in fact behaves discontinuously at $\vartheta = 0$ since, counter to first intuition, the spectrum of R_{ϑ} “fills” the gap (a, b) as $\vartheta \rightarrow 0$ with $\vartheta > 0$. This implies, in particular, that R_{ϑ} cannot converge to H in the norm resolvent sense, as $\vartheta \rightarrow 0$.

0.1. Theorem. *Let H, R_{ϑ} and (a, b) as above. Then, for any $\varepsilon > 0$ there exists $0 < \vartheta_\varepsilon < \pi/2$ such that for any $E \in (a, b)$ we have*

$$\sigma(R_{\vartheta}) \cap (E - \varepsilon, E + \varepsilon) \neq \emptyset, \quad \forall 0 < \vartheta < \vartheta_\varepsilon. \quad (0.5)$$

0.2. Remarks. (i) Roughly speaking, the moment we start rotating the potential on the left-hand side by a tiny angle the gap (a, b) is suddenly full of spectrum of R_{ϑ} in the sense that, for $0 < \vartheta < \vartheta_\varepsilon$, no gap of R_{ϑ} in the interval (a, b) can have length larger than 2ε . It is conceivable that for most ϑ the spectrum of R_{ϑ} covers the interval (a, b) , but there are examples (cf. Section 4) where R_{ϑ} has gaps in (a, b) for some ϑ .

(ii) It seems to be quite hard to determine the nature of the spectrum of R_{ϑ} for general $\vartheta \in (0, \pi/2)$; however, there are some special angles for which a result from [DS] excludes singular continuous spectrum (cf. Section 5).

(iii) In addition to what is stated in Theorem 0.1 we obtain lower and upper bounds for the spectral densities in the intervals $(E - \varepsilon, E + \varepsilon)$ on a scale that is appropriate to surface states (without knowing that an integrated surface density of states as in [EKSchrS, KS] exists for R_{ϑ}); cf. Theorems 3.1 and 3.2.

There is a simple, intuitive connection between the rotational problem and the related translational problem, given as follows: Starting from the same periodic potential V as above, we now look at

$$W_t(x, y) := \begin{cases} V(x, y), & x \geq 0, \\ V(x + t, y), & x < 0, \end{cases} \quad 0 \leq 1 \leq t, \quad (0.6)$$

and define $D_t := -\Delta + W_t$, acting in $L_2(\mathbb{R}^2)$. In the 1-dimensional case, this problem has been studied in great detail by Korotyaev [Kor1, Kor2]. A different approach was most recently implemented in [HK] where it is shown that some of the results of [Kor1, Kor2] can be recovered with a rather crude variational technique. This method can be easily generalized to the dislocation problem on a strip $\mathbb{R} \times (0, 1)$ and then to the plane \mathbb{R}^2 ; one finds that spectrum of D_t crosses the gap as t varies between 0 and 1. Now our key observation consists in the following: for any given $\varepsilon > 0$ and $n \in \mathbb{N}$, we can find points $(0, \eta)$ on the y -axis such that

$$|V_\vartheta(x, y) - W_t(x, y)| < \varepsilon, \quad (x, y) \in Q_n(0, \eta), \quad (0.7)$$

with $Q_n(0, \eta) = (-n, n) \times (\eta - n, \eta + n)$, provided $\vartheta > 0$ is small enough and satisfies a condition which ensures an appropriate alignment of the period cells on the y -axis. This basic observation is somewhat reminiscent of a key idea in [HHK] where rotationally symmetric Schrödinger operators of the type $-\Delta + U(|x|)$ in $L_2(\mathbb{R}^n)$ with $U: \mathbb{R} \rightarrow \mathbb{R}$ periodic and Lipschitz-continuous are considered: here, far away from the origin, the potential $U(|x|)$ looks very much like a potential depending only on the x_1 -coordinate if we restrict our attention to balls of fixed size with centers far out on the x_1 -axis.

The paper is organized as follows. In Section 1, we briefly summarize some results of [HK] on translational lattice dislocations for the strip and for the plane. What we will use in the sequel is the simple fact that, for any $E \in (a, b)$, there is some $t = t_E \in (0, 1)$ with the following property: for any $\varepsilon > 0$, there is a compactly supported approximate eigenfunction u in the domain of D_t that satisfies $\|(D_t - E)u\| < \varepsilon$ and $\|u\| = 1$.

In Section 2, we first employ the Birkhoff Ergodic Theorem to obtain a set $\Theta \subset (0, \pi/2)$ with countable complement so that (0.7) can be established for small $\vartheta \in \Theta$ and suitable $\eta \in \mathbb{R}$. Then the above approximate eigenfunctions u will also be approximate eigenfunctions of R_ϑ for small $\vartheta \in \Theta$, after an appropriate (ϑ -dependent) translation along the y -axis. This then gives Theorem 0.1.

Suitable points $(0, \eta)$ for the construction of Section 2 occur with a certain density and we expect a lower bound for the integrated surface density of states. Since we do not know whether the i.d.s. measure or the integrated surface density of states measure exist, we only provide lower and upper bounds for the number of Dirichlet eigenvalues in subintervals of the gap (a, b) for our operators R_ϑ , restricted to large squares $Q_n = (-n, n)^2$. Theorem 3.1 in Section 3 provides a lower bound of the form $c_1 n$ for n large with a positive constant c_1 , while Theorem 3.2 gives an upper bound by $c_2 n \log n$, for n large. Note that Theorem 3.2 deals with a much more general situation: in fact, we allow here for two different potentials V_ℓ and V_r on the left and right which are not required to be periodic; all we need is a common gap.

In Section 4 we discuss examples of “muffin tin”-type which come in three versions: in the simplest case, the muffin tins are circular wells, arranged on a periodic grid, with infinitely high walls (so that the Schrödinger operator is just the direct sum of a countable number of Dirichlet Laplacians on circles). We then approximate

by muffin tins of finite height, and, finally, by muffin tins with Lipschitz potentials. We obtain spectral results for the rotation problem for all three versions.

In Section 5, finally, we first explain a simplified model for small angle grain boundaries where we assume rotation by an angle $\vartheta/2$ in both half-planes in opposite direction; this problem is somewhat easier to deal with. We furthermore discuss special rotation angles ϑ for which the operators R_ϑ are periodic in the y -direction and thus have no singular continuous spectrum [DS]. All these questions hinge on an approximate or exact matching up between the given periodic potential in the right half-plane and its rotated version in the left half-plane. This is somewhat connected with the question of coincidence between a lattice and some of its rotated versions (CSL-lattices).

For basic results on the spectral theory of self-adjoint operators or, more specifically, periodic Schrödinger operators, we refer to [K], [RS-IV], [E] and [Ku]. We will use results from these references mostly without specific reference.

Acknowledgements. The authors would like to thank Andreas Ruschhaupt, Univ. Hannover, for several fruitful discussions.

1. THE DISLOCATION PROBLEM ON A STRIP AND FOR THE PLANE

Let $V: \mathbb{R}^2 \rightarrow \mathbb{R}$ be \mathbb{Z}^2 -periodic and bounded and let $\Sigma := \mathbb{R} \times (0, 1)$ denote the infinite strip of width 1. As above, we write $H := -\Delta + V$ for the (self-adjoint) Schrödinger operator with potential V acting in $L_2(\mathbb{R}^2)$. Then $\sigma(H)$, the spectrum of H , has band structure, i.e., it is the (locally finite) union of compact intervals [RS-IV]. The intervals of spectrum, the *bands*, may be separated by (open) intervals, the *gaps*. Moreover, $\sigma(H)$ is purely absolutely continuous. For $0 \leq t \leq 1$, we introduce the families

$$\begin{aligned} S_t &:= -\Delta + W_t, & \text{acting in } L_2(\Sigma), \\ D_t &:= -\Delta + W_t, & \text{acting in } L_2(\mathbb{R}^2), \end{aligned} \tag{1.1}$$

of self-adjoint operators, where S_t has periodic boundary conditions in the y -variable and W_t is as in (0.6). The operator $-\Delta + W_t$ in $L_2(\Sigma)$ with ϑ -periodic boundary conditions is denoted by $S_t(\vartheta)$. In [HK], it is shown that the essential spectrum of S_t does not depend on the dislocation parameter $t \in [0, 1]$. However, discrete eigenvalues of S_t cross the spectral gaps of H as t increases from 0 to 1. The existence of these eigenvalues is obtained by counting the eigenvalues below E for the operators $S_t^{(n)} := -\Delta + W_t$ on $(-n - t, n) \times (0, 1)$ with periodic boundary conditions in x and y . As t increases from 0 to 1, the number of eigenvalues below E grows by 1. The eigenfunctions of $S_t^{(n)}$ associated with an eigenvalue E in the gap (a, b) are localized near the interface $\{x = 0\}$ and we may take the limit $n \rightarrow \infty$ to obtain eigenfunctions of S_t .

Periodic continuation of the eigenfunctions on the strip and appropriate cut-offs yield compactly supported approximate eigenfunctions of D_t . This leads to the following result (cf. Thm. 4.3 and Remark 4.4 of [HK]).

1.1. Proposition. *Let (a, b) denote a spectral gap of H and let $E \in (a, b)$. Then there exists $t = t_E \in (0, 1)$ such that $E \in \sigma(D_t)$.*

Moreover, for any $n \in \mathbb{N}$ there are functions $v_n = v_n(x, y)$ in the domain of D_t satisfying $\|v_n\| = 1$, $\text{supp } v_n \subset [-n, n]^2$ and $(D_t - E)v_n \rightarrow 0$ as $n \rightarrow \infty$.

For a detailed discussion and further results, we refer to [HK].

2. THE ROTATION PROBLEM FOR SMALL ANGLES

In this section, we study the spectrum of the operators $(R_\vartheta; 0 < \vartheta < \pi/2)$ where the R_ϑ , defined in (0.4), are self-adjoint operators in the Hilbert space $L_2(\mathbb{R}^2)$.

In view of a proof of Theorem 0.1, consider a fixed $E \in (a, b)$. Then, by Proposition 1.1, there is some $t \in (0, 1)$ such that E is in the spectrum of the dislocation operator D_t on the plane. We wish to find angles ϑ with the property that the potential V_ϑ is approximately equal to W_t on a sufficiently large square $Q_n(0, \eta)$ of side-length $2n$, centered at some point $(0, \eta)$ on the y -axis. This leads to the following requirements: If we imagine the grid $\Gamma = \{(x, y) \in \mathbb{R}^2; x \in \mathbb{Z} \text{ or } y \in \mathbb{Z}\}$ of lines describing the period cells, we have to make sure that, inside $Q_n(0, \eta)$, the alignment between the horizontal lines of Γ in the right half-plane with the rotated horizontal lines of $M_\vartheta\Gamma$ in the left half-plane is nearly perfect on the y -axis and that the rotated vertical lines of $M_\vartheta\Gamma$ in the left half-plane have, roughly, distance t (modulo \mathbb{Z}) from the y -axis. More precisely, we wish to find $m \in \mathbb{N}$ such that $m/\cos\vartheta$ is integer, up to a small error, and $m \tan\vartheta = t \pmod{\mathbb{Z}}$, again up to a small error, inside $Q_n(0, \eta)$.

We first prepare a lemma which deals with ergodicity on the flat torus $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$, as in [RS-I], [CFS]. We consider transformations $T_\vartheta: \mathbb{T}^2 \rightarrow \mathbb{T}^2$ defined by

$$T_\vartheta(x, y) := (x + \tan\vartheta, y + 1/\cos\vartheta). \quad (2.1)$$

2.1. Lemma. *There is a set $\Theta \subset (0, \pi/2)$ with countable complement such that the transformation T_ϑ in (2.1) is ergodic for all $\vartheta \in \Theta$.*

Proof. T_ϑ is ergodic if and only if the numbers 1, $\tan\vartheta$, and $1/\cos\vartheta$ are independent over the rationals, i.e., $(n_1, n_2, n_3) \in \mathbb{Z}^3$ and

$$n_1 + n_2 \tan\vartheta + \frac{n_3}{\cos\vartheta} = 0 \quad (2.2)$$

implies $n_1 = n_2 = n_3 = 0$. Write $\mathcal{Z}_3 := \mathbb{Z}^3 \setminus \{(0, 0, 0)\}$. For any triple $(n_1, n_2, n_3) \in \mathcal{Z}_3$ the set of points $(x, y) \in \mathbb{R}^2$ that satisfy $n_1 + n_2x + n_3y = 0$ is a line $\ell = \ell_{(n_1, n_2, n_3)} \subset \mathbb{R}^2$. Consider the (countable) set

$$\Lambda := \{\ell_{(n_1, n_2, n_3)}; (n_1, n_2, n_3) \in \mathcal{Z}_3\}. \quad (2.3)$$

In (2.2), the variables x and y are of the special form $x = \tan\vartheta$, $y = 1/\cos\vartheta$ and so $y = \sqrt{1+x^2}$. Since $F(x) := \sqrt{1+x^2}$ is convex, each $\ell \in \Lambda$ has at most two intersection points with the graph Γ of F . Then

$$S := \cup_{\ell \in \Lambda} \{\ell \cap \Gamma\} \quad (2.4)$$

is countable and so $\Gamma \setminus S$ has full measure. Let $\Gamma_+ := \Gamma \cap \{x > 0\}$. We map $\Gamma_+ \setminus S$ to $(0, \pi/2)$ by $h: (x, F(x)) \mapsto \arctan x$. Since $h: \Gamma_+ \rightarrow (0, \pi/2)$ is diffeomorphic, $\Theta := h(\Gamma_+ \setminus S)$ is as desired. \square

Let us write x_\sim for the fractional part of $x > 0$, i.e., $x_\sim = x - [x]$ if $x > 0$. In the proof of our main theorem, we will need natural numbers m such that, for $t \in (0, 1)$ given, $(m \tan \vartheta)_\sim$ is approximately equal to t and $(m/\cos \vartheta)_\sim$ almost equals 0. The existence of such numbers m follows from Lemma 2.1 and Birkhoff's Ergodic Theorem. Let $\vartheta \in \Theta$, $\varepsilon > 0$, and let us denote by χ_Q the characteristic function of the set $Q := (t - \varepsilon, t + \varepsilon) \times (-\varepsilon, \varepsilon) \subset \mathbb{T}^2$. Then, for all $x \in \mathbb{T}^2$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=0}^{n-1} \chi_Q(T_\vartheta^m x) = \int_Q dx = 4\varepsilon^2 > 0. \quad (2.5)$$

By setting $x = (0, 0)$ we are done. We are now ready for a first main result which establishes the existence of surface states in the gaps of H and shows that, in fact, any gap (a, b) of H is filling up with spectrum of R_ϑ as $\vartheta \rightarrow 0$.

2.2. Proposition. *Let (a, b) be a spectral gap of H and let $[\alpha, \beta] \subset (a, b)$, $\alpha < \beta$. Then there is a $\vartheta_0 = \vartheta_0(\alpha, \beta) > 0$ such that*

$$\sigma(R_\vartheta) \cap (\alpha, \beta) \neq \emptyset, \quad \forall \vartheta \in (0, \vartheta_0). \quad (2.6)$$

Proof. (1) We first restrict our attention to $\vartheta \in \Theta$ with Θ as in Lemma 2.1. Let $E \in (\alpha, \beta)$ and $\varepsilon := \min\{E - \alpha, \beta - E\}/2$. By Proposition 1.1, we can find $n = n_\varepsilon \in \mathbb{N}$ and a function u_n of norm 1 in the domain of D_t with $\text{supp } u_n \subset [-n, n]^2$ such that $\|(D_t - E)u_n\| < \varepsilon$. Obviously $u_{n,k}(x, y) := u_n(x, y - k)$ satisfies the same estimate for any $k \in \mathbb{N}$. If we can show that, for appropriate $k \in \mathbb{N}$,

$$|V_\vartheta(x, y) - W_t(x, y)| < \varepsilon, \quad (x, y) \in Q_n(0, k), \quad (2.7)$$

we may conclude that

$$\|(R_\vartheta - E)u_{n,k}\| < 2\varepsilon; \quad (2.8)$$

but then the Spectral Theorem implies that R_ϑ has spectrum inside the interval $(E - 2\varepsilon, E + 2\varepsilon) \subset (\alpha, \beta)$.

For a proof of (2.7), we first observe that by the properties of V and the definitions of V_ϑ and W_t , we have the following estimate:

$$|V_\vartheta(x, y) - W_t(x, y)|^2 \leq \min_{j_1, j_2 \in \mathbb{Z}} L^2((X - j_1)^2 + (Y - j_2)^2), \quad \forall (x, y) \in \mathbb{R}^2, \quad (2.9)$$

with

$$X := x(\cos \vartheta - 1) - t + y \sin \vartheta, \quad Y := -x \sin \vartheta + y(\cos \vartheta - 1) \quad (2.10)$$

and L the Lipschitz constant of V . Now for $\vartheta \in \Theta$ given, there is some $m = m_\vartheta \in \mathbb{N}$ such that

$$\left(\frac{m}{\cos \vartheta}\right)_\sim < \varepsilon/4, \quad |(m \tan \vartheta)_\sim - t| < \varepsilon/4; \quad (2.11)$$

in particular, there is some $N \in \mathbb{N}$ s.th. $|m/\cos \vartheta - N| < \varepsilon/4$.

We may now apply the estimate (2.9) to the points $(x, y) \in Q_n(0, N)$ to find

$$|V_\vartheta(x, y) - W_t(x, y)|^2 \leq L^2 ((X - \lfloor m \tan \vartheta \rfloor)^2 + (Y + N - m)^2), \quad (2.12)$$

for all $(x, y) \in Q_n(0, N)$. Here

$$|X - \lfloor m \tan \vartheta \rfloor| \leq n(1 - \cos \vartheta) + n\vartheta + |m \tan \vartheta - \lfloor m \tan \vartheta \rfloor - t| \leq 2n_\varepsilon \vartheta + |(m \tan \vartheta) \sim - t| \quad (2.13)$$

and

$$|Y + N - m| \leq 2n_\varepsilon \vartheta + |N - m / \cos \vartheta|. \quad (2.14)$$

We choose $\vartheta_0 > 0$ small enough to have $2n_\varepsilon \vartheta_0 < \varepsilon/4$ and (2.7) follows if we pick $k := N$. We have thus shown that R_ϑ has spectrum in (α, β) for all $\vartheta \in \Theta \cap (0, \vartheta_0)$.

(2) In order to remove the restriction $\vartheta \in \Theta$ we note that with each $\vartheta \in \Theta$ there comes a positive number $\eta_\vartheta > 0$ such that

$$\|(R_\sigma - E)u_{n,k}\| < 3\varepsilon, \quad \forall \sigma \in (\vartheta - \eta_\vartheta, \vartheta + \eta_\vartheta), \quad (2.15)$$

since

$$\|(V_\sigma - V_\vartheta) \upharpoonright \text{supp } u_{n,k}\|_\infty \rightarrow 0, \quad \sigma \rightarrow \vartheta. \quad (2.16)$$

As the intervals $(\vartheta - \eta_\vartheta, \vartheta + \eta_\vartheta)$ with ϑ ranging between 0 and ϑ_0 cover the interval $(0, \vartheta_0)$, the desired result follows. \square

Now it is easy to obtain Theorem 0.1 in the Introduction from Proposition 2.2:

Proof of Theorem 0.1. For $\varepsilon > 0$ given, we consider points $a = \gamma_0 < \gamma_1 < \gamma_2 < \dots < \gamma_N = b$ such that $\gamma_j - \gamma_{j-1} < \varepsilon/2$, for $j = 1, \dots, N$. For each of the intervals $I_j := (\gamma_{j-1}, \gamma_j)$, $2 \leq j \leq N-1$, Proposition 2.2 yields a constant $\vartheta_j > 0$ with the property that R_ϑ has spectrum in the interval I_j for all $0 < \vartheta < \vartheta_j$. Then $\vartheta_0 := \min_{2 \leq j \leq N-1} \vartheta_j$ has the required properties. \square

3. INTEGRATED DENSITY OF STATES BOUNDS

It is clear that ergodicity gives us not just a single m as in (2.11), for $\vartheta \in \Theta$; in fact, eqn. (2.5) guarantees that suitable m will appear with a certain frequency. We will use this observation to obtain lower bounds for a quantity which, in the limit, would translate into a (positive) lower bound for the surface i.d.s. measure if we knew that the required limit exists. This will be complemented by a similar upper bound which is of the expected order, up to a logarithmic factor. A detailed and rather complete account of the i.d.s. for (random) Schrödinger operators can be found in [V] which also contains a wealth of references. [EKSchrS] and [KS] specifically discuss the existence of a surface i.d.s. as a distribution or a measure. Some results on the surface i.d.s. measure for the translational dislocation problem can be found in [HK].

Let $R_\vartheta^{(n)}$ denote the operator $-\Delta + V_\vartheta$, acting in $L_2(Q_n)$ with Dirichlet boundary conditions, where $Q_n := (-n, n)^2 \subset \mathbb{R}^2$. For any interval $I \subset \mathbb{R}$, we denote by $N_I(R_\vartheta^{(n)})$ the number of eigenvalues of $R_\vartheta^{(n)}$ in I , each eigenvalue being counted according to its multiplicity. The existence of a surface i.d.s. measure in the gap

(a, b) would correspond to the existence of a finite limit $\lim_{n \rightarrow \infty} \frac{1}{n} N_I(R_\vartheta^{(n)})$, for any interval I with $\bar{I} \subset (a, b)$. Theorem 3.1 provides lower bounds of the form

$$\liminf_{n \rightarrow \infty} \frac{1}{n} N_I(R_\vartheta^{(n)}) > 0, \quad (3.1)$$

for (non-degenerate) subintervals I and small $\vartheta \in \Theta$, while Theorem 3.3 yields an upper bound

$$\limsup_{n \rightarrow \infty} \frac{1}{n \log n} N_I(R_\vartheta^{(n)}) < \infty. \quad (3.2)$$

We begin with a lower bound.

3.1. Theorem. *Let H, R_ϑ as above and suppose that (a, b) is a spectral gap of H . Let Θ as in Lemma 2.1.*

Then, for any $\varepsilon > 0$ there exists a $\vartheta_\varepsilon > 0$ such that (3.1) holds for all $\vartheta \in \Theta \cap (0, \vartheta_\varepsilon)$ and for any interval $I \subset (a, b)$ of length greater than ε .

Proof. (1) Let $[\alpha, \beta] \subset (a, b)$, fix $E \in (\alpha, \beta)$, and let $0 < \varepsilon < \min\{E - \alpha, \beta - E\}$. Let u_0 in the domain of D_t with compact support satisfy $\|u_0\| = 1$ and $\|(D_t - E)u_0\| < \varepsilon$, as in Prop. 1.1. Let $\nu \in \mathbb{N}$ be such that $\text{supp } u_0 \subset Q_\nu = (-\nu, \nu)^2$; note that, in this proof, ν corresponds to the parameter n of Section 2.

Let $\vartheta \in \Theta \cap (0, \pi/4]$ so that, in particular, $1/\sqrt{2} \leq \cos \vartheta \leq 1$. By ergodicity, there exists a constant $c_0 = c_0(\vartheta) > 0$ with the following properties: for $n \in \mathbb{N}$ large, there are at least $J_n := \lfloor c_0 n \rfloor$ natural numbers $m_1, \dots, m_{J_n} \in (0, n/4)$ such that (2.11) holds for $m = m_s$, $s = 1, \dots, J_n$, and such that

$$|m_s - m_r| \geq 2\nu, \quad s \neq r, \quad 1 \leq s, r \leq J_n; \quad (3.3)$$

here J_n and m_1, \dots, m_{J_n} depend on n and ϑ . It follows that for each $j = 1, \dots, J_n$ there is some $N_j \in \mathbb{N}$ such that $|m_s / \cos \vartheta - N_j| < \varepsilon/4$ and $|m_s \tan \vartheta - t|_\sim < \varepsilon/4$. We then see that the functions φ_j , defined by $\varphi_j(x, y) := u_0(x, y - N_j)$, are of norm 1 and have mutually disjoint supports contained in $(-n, n)^2$. Furthermore, for ϑ small enough, $0 < \vartheta < \vartheta_\varepsilon$, say, we can show (as in the proof of Proposition 2.2) that an estimate (2.7) holds on each square $(-\nu, \nu) \times (N_j - \nu, N_j + \nu)$. Thus (2.7) holds on the support of each φ_j and it follows that

$$\left\| (R_\vartheta^{(n)} - E)\varphi_j \right\| < \varepsilon, \quad 0 < \vartheta < \vartheta_\varepsilon, \quad j = 1, \dots, J_n. \quad (3.4)$$

Then $\mathcal{M} := \text{span}\{\varphi_j; j = 1, \dots, J_n\}$ has dimension J_n . Let \mathcal{N} denote the range of the spectral projection $P_{(\alpha, \beta)}(R_\vartheta^{(n)})$ of $R_\vartheta^{(n)}$ associated with the interval (α, β) and assume for a contradiction that $\dim \mathcal{N} < J_n$. Then we can find a function $v \in \mathcal{M} \cap \mathcal{N}^\perp$ of norm 1. By the Spectral Theorem, $\left\| (R_\vartheta^{(n)} - E)v \right\| \geq \varepsilon$. On the other hand, (3.4) and $v = \sum_{i=1}^{J_n} a_i \varphi_i$ implies $\left\| (R_\vartheta^{(n)} - E)v \right\| < \varepsilon$ because the φ_j have mutually disjoint supports.

We have therefore shown that for any interval $I = [\alpha, \beta]$ there exists some $\vartheta_0 > 0$ such that (3.1) holds for all $\vartheta \in \Theta \cap (0, \vartheta_0)$.

(2) Now let $\varepsilon > 0$. As in the proof of Theorem 0.1, given at the end of Section 2, we may cover the interval (a, b) by a finite number of subintervals of length ε ; applying the result of part (1) we then obtain the desired statement. \square

Remarks. (a) It appears that the argument used at the end of the proof of Proposition 2.2 to remove the restriction $\vartheta \in \Theta$ does not work in the context of Theorem 3.1.

(b) It follows from the proof of Thm. 3.1 that $\sigma_{\text{ess}}(R_\vartheta) \cap I \neq \emptyset$ for all $\vartheta \in \Theta \cap (0, \vartheta_\varepsilon)$ and for any interval $I \subset (a, b)$ of length greater than ε .

We now complement the lower estimate established in Theorem 3.2 by an upper bound which is of the expected order, up to a logarithmic factor. Note that we treat a situation which is far more general than the rotation or dislocation problems studied so far. In fact, we will allow for different potentials V_1 on the left and V_2 on the right which are only linked by the assumption that there is a common spectral gap; neither V_1 nor V_2 are required to be periodic. The proof uses technology which is fairly standard and based on exponential decay estimates for resolvents.

3.2. Theorem. *Let $V_1, V_2 \in L_\infty(\mathbb{R}^2, \mathbb{R})$ and suppose that the interval $(a, b) \subset \mathbb{R}$ does not intersect the spectra of the self-adjoint operators $H_k := -\Delta + V_k$, $k = 1, 2$, both acting in the Hilbert space $L_2(\mathbb{R}^2)$. Let*

$$W := \chi_{\{x < 0\}} \cdot V_1 + \chi_{\{x \geq 0\}} \cdot V_2 \quad (3.5)$$

and define $H := -\Delta + W$, a self-adjoint operator in $L_2(\mathbb{R}^2)$. Finally, we let $H^{(n)}$ denote the self-adjoint operator $-\Delta + W$ acting in $L_2(Q_n)$ with Dirichlet boundary conditions. Then, for any interval $[a', b'] \subset (a, b)$, we have

$$\limsup_{n \rightarrow \infty} \frac{1}{n \log n} N_{[a', b']}(H^{(n)}) < \infty. \quad (3.6)$$

Proof. (1) We write $N(n) := N_{[a', b']}(H^{(n)})$ and note that there is a constant $c_0 \geq 0$ such that

$$N(n) \leq c_0 n^2, \quad n \in \mathbb{N}; \quad (3.7)$$

this follows by routine min-max arguments as in [RS-IV; Section XIII.15].

(2) Let us consider the (normalized) eigenfunctions $u_{i,n}$ of $H^{(n)}$ associated with the eigenvalues $E_{i,n} \in [a', b']$, for $i = 1, \dots, N(n)$. The main idea of the proof is to show that the $u_{i,n}$ are concentrated near the boundary of Q_n or near the y -axis. To obtain the corresponding estimates, we introduce the sets

$$\Omega_j(n) := \Omega_j^-(n) \cup \Omega_j^+(n), \quad j \in \{1, 2, 3, 4\}, \quad (3.8)$$

where $\Omega_j^-(n) := \left(-\frac{n}{2} + \frac{2j}{\alpha} \log n, -\frac{2j}{\alpha} \log n\right) \times \left(-\frac{n}{2} + \frac{2j}{\alpha} \log n, \frac{n}{2} - \frac{2j}{\alpha} \log n\right)$, and $\Omega_j^+(n) := -\Omega_j^-(n)$ is the mirror-image of $\Omega_j^-(n)$ with respect to the y -axis; the parameter $\alpha > 0$ will be chosen as in eqn. (3.9) below. Note that, for $\alpha > 0$ fixed, the sets $\Omega_1(n), \dots, \Omega_4(n)$ are non-empty for n large. We have the trivial inclusions $\Omega_{j+1}(n) \subset \Omega_j(n)$ for $j = 1, 2, 3$.

We will use the following exponential decay estimate for the resolvent of the operators H_k : There are constants $C \geq 0$, $\alpha > 0$ such that for any $E \in [a', b']$ and (measurable) sets $K_1, K_2 \subset \mathbb{R}^2$ we have (cf., e.g., [AADH; Prop. 2.4])

$$\|\chi_{K_1} \partial_j^p (H_k - E)^{-1} \chi_{K_2}\| \leq C e^{-\alpha \text{dist}(K_1, K_2)}, \quad j, p \in \{0, 1\}, \quad k \in \{1, 2\}; \quad (3.9)$$

here $\partial_1 = \partial_x$, $\partial_2 = \partial_y$. We also choose cut-off functions $\varphi_n, \psi_n \in C_c^\infty(\mathbb{R}^2; \mathbb{R})$ satisfying

$$\text{supp } \varphi_n \subset \Omega_1(n), \quad \varphi_n \upharpoonright \Omega_2(n) = 1, \quad \text{supp } \psi_n \subset \Omega_3(n), \quad \psi_n \upharpoonright \Omega_4(n) = 1, \quad (3.10)$$

and $|\nabla \varphi_n|, |\nabla \psi_n|, |\partial_{ij} \varphi_n|, |\partial_{ij} \psi_n| \leq c(\log n)^{-1}$ with some constant $c \geq 0$; here $\varphi_n = \varphi_{n,\ell} + \varphi_{n,r}$ with $\varphi_{n,\ell}$ and $\varphi_{n,r}$ being supported in $\Omega_1^-(n)$ and $\Omega_1^+(n)$, respectively. By a well-known argument we can now derive the desired localization property: by the Leibniz rule, we have for $i = 1, \dots, N(n)$

$$(H_1 - E_i)(\varphi_{n,\ell} u_{i,n}) = (H^{(n)} - E_i)(\varphi_{n,\ell} u_{i,n}) = -2\nabla \varphi_{n,\ell} \cdot \nabla u_{i,n} - \Delta \varphi_{n,\ell} u_{i,n} \quad (3.11)$$

so that

$$\chi_{\Omega_3^-(n)} u_{i,n} = -\chi_{\Omega_3^-(n)} (H_1 - E_i)^{-1} \chi_{\text{supp } \nabla \varphi_{n,\ell}} [2\nabla \varphi_{n,\ell} \cdot \nabla u_{i,n} + \Delta \varphi_{n,\ell} u_{i,n}]. \quad (3.12)$$

Using that $\text{dist}(\Omega_3(n), \text{supp } \nabla \varphi_n) \geq 2\alpha^{-1} \log n$ and $|\nabla \varphi_n|, |\Delta \varphi_n| \leq c(\log n)^{-1}$, the estimate (3.9) implies that

$$\|\chi_{\Omega_3(n)} u_{i,n}\|, \|\chi_{\Omega_3(n)} \nabla u_{i,n}\| \leq C(n^2 \log n)^{-1}, \quad i = 1, \dots, N(n). \quad (3.13)$$

We now define $v_{i,n} := (1 - \psi_n)u_{i,n}$ and let $\mathcal{M}_n := \text{span}\{v_{i,n}; i = 1, \dots, N(n)\}$. We claim that

$$\dim \mathcal{M}_n = N(n), \quad n \geq n_0, \quad (3.14)$$

for some $n_0 \in \mathbb{N}$. Let $H_{Q_n \setminus \Omega_4(n)}$ be the operator $-\Delta + W$ on $Q_n \setminus \Omega_4(n)$ with Dirichlet boundary conditions. The functions $v_{i,n} := (1 - \psi_n)u_{i,n}$ are approximate eigenfunctions of $H_{Q_n \setminus \Omega_4(n)}$: in fact, using (3.13), one easily checks that

$$\|(H_{Q_n \setminus \Omega_4(n)} - E_{i,n})v_{i,n}\| \leq C(n^2 \log^2 n)^{-1} \quad (3.15)$$

and

$$\|v_{i,n} - u_{i,n}\| \leq C(n^2 \log n)^{-1}, \quad (3.16)$$

for $i = 1, \dots, N(n)$. Now (3.7) and (3.16) imply $\sum_{i=1}^{N(n)} \|u_{i,n} - v_{i,n}\|^2 < 1$ for n large and we obtain (3.14).

(3) We next show that there is $n_1 \geq n_0 \in \mathbb{N}$ such that

$$\langle H_{Q_n \setminus \Omega_4(n)} w, w \rangle < b \|w\|^2, \quad w \in \mathcal{M}_n, \quad n \geq n_1. \quad (3.17)$$

For a proof, consider an arbitrary $w = \sum_{i=1}^{N(n)} \gamma_i v_{i,n} \in \mathcal{M}_n$ with $\|w\| = 1$. Here we first observe that the coefficients γ_i satisfy a bound $|\gamma_i| \leq 2$, for n large, since (writing $\gamma^2 := \sum_i |\gamma_i|^2$ and $\eta_n^2 := \sum_i \|v_{i,n} - u_{i,n}\|^2$)

$$1 = \|w\| \geq \left\| \sum_{i=1}^{N(n)} \gamma_i u_{i,n} \right\| - \sum_{i=1}^{N(n)} |\gamma_i| \cdot \|v_{i,n} - u_{i,n}\| \geq \gamma(1 - \eta_n), \quad (3.18)$$

where $\eta_n \rightarrow 0$ as $n \rightarrow \infty$ by (3.16). Using (3.16) and the fact that $\nabla\psi_n$ and $\Delta\psi_n$ have support in $\Omega_3(n) \setminus \Omega_4(n)$, it follows that for n large

$$\|w\|^2 = \sum_{i=1}^{N(n)} |\gamma_i|^2 + r, \quad \langle H_{Q_n \setminus \Omega_4(n)} w, w \rangle = \sum_{i=1}^{N(n)} E_i |\gamma_i|^2 + r', \quad (3.19)$$

where $r, r' \leq C(\log n)^{-2}$, so that

$$\langle H_{Q_n \setminus \Omega_4(n)} w, w \rangle \leq b' \|w\|^2 + r'', \quad (3.20)$$

with $r'' \leq C(\log n)^{-2}$, for n large, and we obtain (3.17).

(4) We conclude from (3.17) that $\mathcal{M}_n \subset P_{(-\infty, b)}(H_{Q_n \setminus \Omega_4(n)})$ and then (3.14) implies that $\dim P_{(-\infty, b)}(H_{Q_n \setminus \Omega_4(n)}) \geq \dim \mathcal{M}_n = N(n)$. On the other hand, min-max arguments yield an upper bound for $\dim P_{(-\infty, b)}(H_{Q_n \setminus \Omega_4(n)})$ of the form $cn \log n$, and we are done. \square

It seems to be possible, using more powerful methods, to remove the logarithmic factor in (3.2) and (3.6) (H. Cornean, private communication, 2010).

4. MUFFIN TIN POTENTIALS

In this section, we present a class of examples where one can arrive at rather precise statements that illustrate some of the phenomena described before. Note that the results given below are derived directly, without recourse to Section 2. We will look at three types of muffin tin potentials and discuss the effect of the “filling up” of the gaps at small angles of rotation. We begin with muffin tins with walls of infinite height, then approximate by muffin tin potentials of height n , for $n \in \mathbb{N}$ large. By another approximation step, one may obtain examples with Lipschitz-continuous potentials. These examples show, among other things, that Schrödinger operators of the form R_ϑ may in fact have spectral gaps for some $\vartheta > 0$.

We consider the lattice $\mathbb{Z}^2 \subset \mathbb{R}^2$ where we first introduce the Laplacian of a periodic muffin tin with infinitely high walls separating the wells: for $0 < r < 1/2$, we let $D_r := B_r(\frac{1}{2}, \frac{1}{2})$ denote the disc of radius r centered at the point $(\frac{1}{2}, \frac{1}{2}) \in \mathbb{R}^2$, and generate from D_r the periodic sets

$$\Omega_r := \cup_{(i,j) \in \mathbb{Z}^2} (D_r + (i, j)), \quad 0 < r < 1/2. \quad (4.1)$$

The Dirichlet Laplacian H_r of Ω_r is the direct sum of a countable number of copies of the Dirichlet Laplacian on D_r ; therefore, the spectrum of H_r consists in a sequence of positive eigenvalues $(\mu_k(r))_{k \in \mathbb{N}}$ with $\mu_k(r) \rightarrow \infty$ as $k \rightarrow \infty$; we may assume that $\mu_k(r) < \mu_{k+1}(r)$ for all $k \in \mathbb{N}$. The eigenvalues $\mu_k = \mu_k(r)$ of H_r have infinite multiplicity. The μ_k correspond to the bands of a periodic problem: in fact, defining $V_r: \mathbb{R}^2 \rightarrow \mathbb{R}$ by

$$V_r(x, y) := \begin{cases} 0, & (x, y) \in \Omega_r, \\ 1, & (x, y) \notin \Omega_r, \end{cases} \quad (4.2)$$

the periodic Schrödinger operators $H_{r,n} := H_0 + nV_r$ have purely a.c. spectrum with a band/gap structure. Furthermore, norm resolvent convergence $H_{r,n} \rightarrow H_r$, obtained as in [HH], implies that the bands of $H_{r,n}$ converge to the eigenvalues μ_k

of H_r . In the sequel, denote by (a, b) one of the gaps (μ_k, μ_{k+1}) . We next look at the rotation problem where we define

$$\Omega_{r,\vartheta} := \Omega_r \cap \{x \geq 0\} \cup (M_\vartheta \Omega_r) \cap \{x < 0\}; \quad (4.3)$$

we also let $H_{r,\vartheta}$ denote the Dirichlet Laplacian on $\Omega_{r,\vartheta}$, for $0 < r < 1/2$ and $0 \leq \vartheta \leq \pi/4$.

The set $(M_\vartheta \Omega_r) \cap \{x < 0\}$ comes with two types of connected components: most (or, in some cases, all) components are discs, but typically there are also discs in $M_\vartheta \Omega_r$ with center at a distance less than r from the y -axis; those appear in $(M_\vartheta \Omega_r) \cap \{x < 0\}$ in a truncated form. It is then clear that $H_{r,\vartheta}$ has pure point spectrum.

Let us comment on some special cases before we proceed: for $\tan \vartheta$ rational, these truncated discs form a periodic pattern; furthermore, we will find a half-disc in $(M_\vartheta \Omega_r) \cap \{x < 0\}$ if and only if there is a disc in $M_\vartheta \Omega_r$ with center on the y -axis which happens if and only if $\tan \vartheta = 1/(2k+1)$ for some $k \in \mathbb{N}$. It follows that for any $\tan \vartheta \in \mathbb{Q}$ with $\tan \vartheta \notin \{1/(2k+1) \mid k \in \mathbb{N}\}$ there is some $r_0 > 0$ such that *no* component of $M_\vartheta \Omega_r$ meets the y -axis, for $0 < r < r_0$; in other words, in this case all components of $\Omega_{r,\vartheta}$ are discs.




Figure 2: The domain $\Omega_{1/4, \pi/8}$ (shaded).

We now return to the general situation. Moving up the y -axis from the origin, we denote the discs in $M_\vartheta \Omega_r$ that intersect the y -axis by $D_{j;r,\vartheta}$, $j \in J_\vartheta$, with centers $(\xi_{j;\vartheta}, \eta_{j;\vartheta})$; here $J_\vartheta = \emptyset$ or $J_\vartheta = \mathbb{N}$ according to the cases discussed above. Without restriction, we may assume that the $\eta_{j;\vartheta}$ are monotonically increasing. Let

$$C_{j;r,\vartheta} := D_{j;r,\vartheta} \cap \{x < 0\}, \quad j \in J_\vartheta. \quad (4.4)$$

Clearly, the Dirichlet eigenvalues of the components $C_{j;r,\vartheta}$ are the surface states in this model. Since the eigenvalues of the $C_{j;r,\vartheta}$ only depend on r and $\xi_{j;\vartheta}$ (but not on $\eta_{j;\vartheta}$), it is enough to study the eigenvalues of the following sample domains: For $r > 0$ and $-r < \xi \leq r$, we denote

$$C_{r,\xi} := \{(x, y) \in D_r \mid x < 1/2 + \xi\}. \quad (4.5)$$

The Dirichlet Laplacian of $C_{r,\xi}$ has eigenvalues $\lambda_k(r,\xi)$, $k \in \mathbb{N}$, which depend continuously on ξ , for $k \in \mathbb{N}$ and $r > 0$ fixed. Furthermore, $\lambda_k(r,\xi) \rightarrow \infty$ as $\xi \rightarrow -r$ and $\lambda_k(r,\xi) \rightarrow \mu_k(r)$, as $\xi \rightarrow r$. Therefore, for each $k \in \mathbb{N}$, the eigenvalues $\lambda_k(r,\xi)$ range over the interval $[\mu_k(r), \infty)$.

We now combine the above properties of $\lambda_k(r,\xi)$ with information on the distribution of the coordinates $\xi_{j;\vartheta}$. It is easy to see that, as $0 \neq \vartheta \rightarrow 0$, the $\xi_{j;\vartheta}$ partition the interval $(-r, r)$ into subintervals of smaller and smaller length. Therefore, for $\varepsilon > 0$ given, any interval $(\alpha, \beta) \subset (a, b)$ of length $\geq \varepsilon$ will contain an eigenvalue of $H_{r,\vartheta}$ for all sufficiently small $\vartheta > 0$.

We expect stronger properties for angles ϑ for which the set $\{\xi_{j;\vartheta} \mid j \in \mathbb{N}\}$ is dense in $(-r, r)$. As in Section 2, ergodic theory gives us a set $\Theta \subset (0, \pi/2)$ of full measure such that for each $\vartheta \in \Theta$ the set $\{(m \tan \vartheta)_\sim \mid m \in \mathbb{N}\}$ is dense in $(0, 1)$, which implies the desired property for the $\xi_{j;\vartheta}$. It follows that, for any $\vartheta \in \Theta$, the eigenvalues of $H_{r,\vartheta}$ will be dense in $[\mu_1(r), \infty)$.

Finally, for $\tan \vartheta$ rational the $\xi_{j;\vartheta}$ form a periodic set and then $H_{r,\vartheta}$ will only have a finite number of eigenvalues (each of infinite multiplicity) in the gap (a, b) . We thus have the following result:

4.1. Proposition. *Let $0 < r < 1/2$ be fixed.*

(a) *Each $\mu_k(r)$, $k = 1, 2, \dots$, is an eigenvalue of infinite multiplicity of $H_{r,\vartheta}$, for all $0 \leq \vartheta \leq \pi/4$. The spectrum of $H_{r,\vartheta}$ is pure point, for all $0 \leq \vartheta \leq \pi/4$.*

(b) *For any $\varepsilon > 0$ there is a $\vartheta_\varepsilon = \vartheta_\varepsilon(r) > 0$ such that any interval $(\alpha, \beta) \subset (a, b)$ with $\beta - \alpha \geq \varepsilon$ contains an eigenvalue of $H_{r,\vartheta}$ for any $0 < \vartheta < \vartheta_\varepsilon$.*

(c) *There exists a set $\Theta \subset (0, \pi/2)$ of full measure such that $\sigma(H_{r,\vartheta}) = [\mu_1(r), \infty)$. The eigenvalues different from the $\mu_k(r)$ are of finite multiplicity.*

4.2. Remark. If $\tan \vartheta$ is rational, $H_{r,\vartheta}$ has at most a finite number of eigenvalues in (a, b) , each of them of infinite multiplicity. Hence we see a drastic change in the spectrum for $\tan \vartheta \in \mathbb{Q}$ as compared with $\vartheta \in \Theta$. Furthermore, if $\tan \vartheta$ is rational with $\tan \vartheta \notin \{1/(2k+1) \mid k \in \mathbb{N}\}$, then there is some $r_\vartheta > 0$ such that $\sigma(H_{r,\vartheta}) = \sigma(H_r)$ for all $0 < r < r_\vartheta$.

We next turn to muffin tin potentials of finite height. Here we define the potential $V_{r,\vartheta}$ to be zero on $\Omega_{r,\vartheta}$ and $V_{r,\vartheta} = 1$ on the complement of $\Omega_{r,\vartheta}$, where $0 < r < 1/2$ and $0 \leq \vartheta \leq \pi/4$; we also let $H_{r,n,\vartheta} := H_0 + nV_{r,\vartheta}$. The periodic operators $H_{r,n,0}$ have purely absolutely continuous spectrum.

We first show that, for r, ϑ fixed, the operators $H_{r,n,\vartheta}$ converge to $H_{r,\vartheta}$ in the sense of norm resolvent convergence. This can be seen as follows: In view of Theorem A.1, we introduce an additional Dirichlet boundary condition on a (closed) set $S = S_\vartheta \subset \mathbb{R}^2 \setminus \Omega_{r,\vartheta}$, which we now define: Let $\Gamma \subset \mathbb{R}^2$ denote the grid $\{(x, y) \in \mathbb{R}^2 \mid x \in \mathbb{Z} \text{ or } y \in \mathbb{Z}\}$, and let $\Gamma_\vartheta := M_\vartheta \Gamma$ denote the rotated grid. For $\rho := (1/2 - r)/2$, we let S consist of $\Gamma \cap \{(x, y) \mid x > \rho\}$, $\Gamma_\vartheta \cap \{x < \rho\}$ plus the vertical line $\{x = \rho\}$. Note that S has distance $\rho > 0$ from $\Omega_{r,\vartheta}$. Let $H_{r,n,\vartheta;S}$ denote the operator $-\Delta + nV_{r,\vartheta}$ with Dirichlet boundary condition on S . We now

have

$$\begin{aligned} \|(H_{r,\vartheta} + 1)^{-1} \oplus 0 - (H_{r,n,\vartheta} + 1)^{-1}\| &\leq \|(H_{r,\vartheta} + 1)^{-1} \oplus 0 - (H_{r,n,\vartheta;S} + 1)^{-1}\| \\ &\quad + \|(H_{r,n,\vartheta;S} + 1)^{-1} - (H_{r,n,\vartheta} + 1)^{-1}\| \end{aligned} \quad (4.6)$$

where 0 denotes the zero operator on $L_2(\mathbb{R}^2 \setminus \Omega_{r,\vartheta})$. By Theorem A.1, we can find $n_0 \in \mathbb{N}$ such that the second term on the RHS is smaller than any given $\varepsilon > 0$, for $n \geq n_0$. The first term on the RHS is a direct sum of operators living on the components of $\mathbb{R}^2 \setminus S_\vartheta$. By routine arguments, we have norm resolvent convergence on each of the components, as $n \rightarrow \infty$, and all we have to do is to convince ourselves that this convergence holds uniformly on all components. The components of $\mathbb{R}^2 \setminus S_\vartheta$ fall into 4 classes: there are unrotated and rotated squares, there are rectangles in the right half-plane of the form $(\rho, 1) \times (\ell, \ell + 1)$ for $\ell \in \mathbb{Z}$, and there are polygons (triangles, quadrangles, and pentagons) in the set $\{x < \rho\}$ that are bounded to the right by the line $x = \rho$. We have no problem with uniform convergence for the first three classes and Lemma A.2 takes care of the fourth class. It is easy to see that the norm resolvent convergence $H_{r,n,\vartheta} \rightarrow H_{r,\vartheta}$ as $n \rightarrow \infty$ is uniform in $\vartheta \in [0, \pi/4]$. We then obtain from Proposition 4.1 (b) and Remark 4.2 the following results:

4.3. Proposition.

(a) For $\tan \vartheta \in \mathbb{Q}$ the spectrum of $H_{r,n,\vartheta}$ has gaps inside the interval (a, b) for n large. More precisely, if $H_{r,\vartheta}$ has a gap $(a', b') \subset (a, b)$, then, for $\varepsilon > 0$ given, the interval $(a' + \varepsilon, b' - \varepsilon)$ will be free of spectrum of $H_{r,n,\vartheta}$ for n large.

(b) For any $\varepsilon > 0$ there are $\vartheta_0 > 0$ and $n_0 > 0$ such that any interval $(c - \varepsilon, c + \varepsilon) \subset (a, b)$ contains spectrum of $H_{r,n,\vartheta}$ for all $0 < \vartheta < \vartheta_0$ and $n \geq n_0$.

By similar arguments, we can approximate $V_{r,\vartheta}$ by Lipschitz-continuous muffin tin potentials that converge monotonically (from below) to $V_{r,\vartheta}$ in such a way that norm resolvent convergence holds for the associated Schrödinger operators (again uniformly in $\vartheta \in [0, \pi/4]$). The spectral properties obtained are analogous to the ones stated in Proposition 4.3. Note, however, that the statement corresponding to part (b) in Proposition 4.3 is weaker than the result of our main Theorem 0.1.

A brief study of translational dislocation problems for muffin tin potentials can be found in [HK].

5. SOME EXTENSIONS AND REMARKS

(1) A simple variant of the rotation problem consists in rotations in the left and the right half planes through angles $\vartheta/2$ and $-\vartheta/2$, respectively, i.e., we study

$$\tilde{V}_\vartheta(x, y) = \begin{cases} (V \circ M_{-\vartheta/2})(x, y), & x \geq 0, \\ (V \circ M_{\vartheta/2})(x, y), & x < 0; \end{cases} \quad (5.1)$$

this potential might be rather close to the physical situation shown in Figure 1. Here we consider the accompanying translational dislocation potentials

$$\tilde{W}_t(x, y) = \begin{cases} V(x - t/2, y), & x \geq 0, \\ V(x + t/2, y), & x < 0. \end{cases} \quad (5.2)$$

We may then obtain results as in Theorem 0.1 without the use of Birkhoff's theorem: here we only need to take care of the second condition in eqn. (2.11) since the horizontal alignment between the left- and right-hand part of V_ϑ on the y -axis is guaranteed by the definition of \tilde{V}_ϑ .

(2) We have shown that the spectral gaps of H fill with spectrum of R_ϑ as $\vartheta \rightarrow 0$ in the sense that any interval of length $\varepsilon > 0$ inside a gap of H will contain spectrum of R_ϑ for sufficiently small angles. In general, we do not know whether the spectrum of R_ϑ in the gaps of H_0 is pure point, absolutely continuous or singular continuous. However, there are some special angles where we can exclude singular continuous spectrum: if we assume that $\cos \vartheta$ is a rational number, $\cos \vartheta = q/p$ with $p, q \in \mathbb{N}$, and p and q belong to a Pythagorean triple ($p^2 - q^2 = r^2$ for some $r \in \mathbb{N}$), then V_ϑ has period p in y -direction. In this case, a result in [DS] implies that $\sigma(H_\vartheta)$ has no singular continuous part.

(3) It is natural to ask about higher dimensions. Suppose we are given a potential $V: \mathbb{R}^3 \rightarrow \mathbb{R}$, periodic with respect to the lattice \mathbb{Z}^3 . We may then simply consider rotations of the (x, y) -plane by an angle ϑ , i.e., we let $V_\vartheta(x, y, z) = V(x, y, z)$ in $\{(x, y, z); x \geq 0\}$ and $V_\vartheta(x, y, z) = V(M_{-\vartheta}(x, y), z)$ in $\{(x, y, z); x < 0\}$, in which case our methods should apply. However, in \mathbb{R}^3 there are many other rotations for which our methods may or may not work.

(4) Of course, taking the limit $\vartheta \rightarrow 0$ is a mathematical idealization. In real crystals or alloys the lattice and its rotated version have to match up according to certain rules. This is usually only possible for a small number of angles. Related questions in higher dimensions are studied under the name of *coincidence site lattices* (CSL); cf. [B, Z].

6. APPENDIX

In this brief appendix we prove a—rather general—result on decoupling by Dirichlet boundary conditions placed on a set inside a high barrier. The method of proof is fairly standard but we have been unable to find a suitable reference which would have covered our situation. Let $d \in \mathbb{N}$. For some open set $\Omega \subset \mathbb{R}^d$ and a closed set $S \subset \Omega$ of measure zero, we consider for $n \in \mathbb{N}$ the Schrödinger operators $H_n := -\Delta + n\chi_\Omega$, acting in $L_2(\mathbb{R}^d)$, and $H_{n,S} := -\Delta + n\chi_\Omega$ in $L_2(\mathbb{R}^d \setminus S) = L_2(\mathbb{R}^d)$, where $H_{n,S}$ is assumed to obey Dirichlet boundary conditions on the set S . In other words, the associated quadratic forms h_n and $h_{n,S}$ have $C_c^\infty(\mathbb{R}^d)$ and $C_c^\infty(\mathbb{R}^d \setminus S)$ as form cores. We then show that the resolvent difference $(H_n + 1)^{-1} - (H_{n,S} + 1)^{-1}$ goes to zero in norm, as $n \rightarrow \infty$, provided the set S has a positive distance to the boundary of Ω ; note that the set S need not be bounded.

A.1. Theorem. *Let H_n and $H_{n,S}$ as above and suppose that $\text{dist}(S, \partial\Omega) > 0$. Then*

$$\|(H_n + 1)^{-1} - (H_{n,S} + 1)^{-1}\| \rightarrow 0, \quad n \rightarrow \infty. \quad (\text{A.1})$$

Proof. For $f \in L_2(\mathbb{R}^d)$ with $\|f\| \leq 1$, let $u_n := (H_n + 1)^{-1}f$ and $v_n := (H_{n,S} + 1)^{-1}f$. We then have the trivial estimates $\|u_n\| \leq 1$ and

$$\|\nabla u_n\|^2 + n \int_{\Omega} |u_n|^2 dx = h_n[u_n] \leq \langle (H_n + 1)u_n, u_n \rangle = \langle f, u_n \rangle \leq 1, \quad (\text{A.2})$$

so that $\|\nabla u_n\| \leq 1$ and $\int_{\Omega} |u_n|^2 dx \leq 1/n$, for all $n \in \mathbb{N}$; analogous estimates hold for v_n . We let \mathcal{F} denote the set of all functions $\varphi \in C^\infty(\mathbb{R}^d)$ that satisfy $\text{supp } \varphi \subset \Omega$, $\text{dist}(\text{supp } \varphi, \partial\Omega) > 0$, $0 \leq \varphi \leq 1$, and $\nabla\varphi, \Delta\varphi$ bounded. For any $\varphi \in \mathcal{F}$, we then have $\varphi u_n \in D(H_n)$ and $(H_n + 1)(\varphi u_n) = \varphi f - 2\nabla u_n \nabla \varphi - u_n \Delta \varphi$. We immediately see from the above estimates that there is a constant $c_\varphi \geq 0$ such that

$$\|H_n(\varphi u_n)\| \leq c_\varphi. \quad (\text{A.3})$$

We now derive a crucial estimate for $\|\nabla(\varphi u_n)\|$: Since $\|\nabla(\varphi u_n)\|^2 \leq h_n[\varphi u_n] = \langle H_n(\varphi u_n), \varphi u_n \rangle$, for any $\varepsilon > 0$ there are constants $C_\varepsilon, C'_\varepsilon \geq 0$ such that

$$\|\nabla(\varphi u_n)\|^2 \leq \varepsilon \|H_n(\varphi u_n)\|^2 + C_\varepsilon \|\varphi u_n\|^2 \leq \varepsilon c_\varphi^2 + C'_\varepsilon/n. \quad (\text{A.4})$$

Analogous estimates hold for $H_{n,S}$ and v_n .

We let $\rho := \text{dist}(S, \partial\Omega)$, $U_\rho := \{x \in \Omega \mid \text{dist}(x, S) < \rho/2\}$ and fix some function $\psi \in \mathcal{F}$ with $\psi = 1$ on U_ρ . Let $\eta := 1 - \psi$ and choose another function $\varphi_\psi \in \mathcal{F}$ satisfying $\varphi_\psi = 1$ on the support of $\nabla\psi$. Writing $w_n = u_n - v_n = (H_n + 1)^{-1}f - (H_{n,S} + 1)^{-1}f$ it is then clear that $\psi w_n \rightarrow 0$, as $n \rightarrow \infty$, uniformly for all f with $\|f\| \leq 1$, and it remains to consider ηw_n . Since η vanishes on U_ρ , we have $\eta w_n \in D(H_n)$ and

$$(H_n + 1)(\eta w_n) = -2\nabla\eta \cdot \nabla w_n - (\Delta\eta)w_n =: z_n, \quad (\text{A.5})$$

so that, in particular,

$$\eta w_n = (H_n + 1)^{-1}z_n, \quad \|\eta w_n\| \leq \|z_n\|. \quad (\text{A.6})$$

We finally show that $z_n \rightarrow 0$, uniformly for $\|f\| \leq 1$. Here we know already that $(\Delta\eta)w_n \rightarrow 0$ since $\Delta\eta$ is supported inside Ω . Applying the estimate (A.4) we find that

$$\|\nabla\eta \cdot \nabla w_n\| \leq C_\eta \|\nabla(\varphi_\psi w_n)\| \leq \varepsilon c_\psi + C'_{\varepsilon, \psi}/n, \quad (\text{A.7})$$

with c_ψ as in (A.3), (A.4). For $\delta > 0$ given, we can find $\varepsilon > 0$ s.th. $\varepsilon c_\psi < \delta/2$ and then $n_0 \in \mathbb{N}$ s.th. $C'_{\varepsilon, \psi}/n < \delta/2$ for all $n \geq n_0$, and we are done. \square

Remarks. (a) The assumption that S has measure zero is made chiefly for simplicity of notation.

(b) The same result holds for Schrödinger operators $-\Delta + nV$ where $V: \mathbb{R}^d \rightarrow \mathbb{R}$ is a non-negative, bounded potential satisfying $V \geq \chi_\Omega$. Our method of proof yields a bound on the norm of the resolvent difference which is independent of V .

(c) It is well-known that H_n and $H_{n,s}$ converge to a suitably defined Dirichlet Laplacians in the sense of strong resolvent convergence, cf. [HZh]. For periodic Ω the convergence would even be in the norm resolvent sense, cf. [HH].

We finally discuss uniform norm resolvent convergence for Schrödinger operators on domains of the type encountered at the end of Section 3. Here we will use the well-known fact that a monotonic sequence $(A_n)_{n \in \mathbb{N}}$ of compact operators which converges strongly to a (compact) operator A converges in norm, i.e., $\|A_n - A\| \rightarrow 0$.

For simplicity of notation, we consider an equivalent geometric situation with a family of domains in the first quadrant given as follows:

We fix $0 < \vartheta < \pi/4$ and $0 < r, d < 1/2$. Then, for $s \in \mathbb{R}$, we consider two parallel lines $\ell_{1,s}, \ell_{2,s}$ defined, respectively, by the equations $y = (x - s) \cot \vartheta$ and $y = (x - s) \cot \vartheta + d/\sin \vartheta$ so that $\ell_{1,s}$ and $\ell_{2,s}$ have distance d . We let

$$G_s := [0, 1]^2 \cap \{(x, y) \mid y > \ell_{1,s}(x)\}, \quad D_{r,s} := B_r(\tfrac{1}{2}, \tfrac{1}{2}) \cap \{(x, y) \mid y > \ell_{2,s}(x)\}; \quad (\text{A.8})$$

cf. Figure 3. Note that $D_{r,s}$ (if it is non-empty) has distance $\min\{d, 1 - r\}$ to the boundary of G_s .




Figure 3: A typical domain $D_{r,s}$ (shaded).

We next define quadratic forms on the domains G_s : for $n \in \mathbb{N}$, we let $V_{r,s,n} := 0$ on $D_{r,s}$ and $V_{r,s,n} := n$ on $G_s \setminus D_{r,s}$. As s increases, the sets G_s and $D_{r,s}$ both increase and we therefore see that the quadratic forms

$$h_{r,s,n}[\varphi] := \|\nabla \varphi\|^2 + \int_{G_s} V_n |\varphi|^2 dx, \quad \varphi \in C_c^\infty(G_s), \quad (\text{A.9})$$

depend monotonically on s . It is easy to include the case $n = \infty$ by setting

$$h_{r,s,\infty}[\varphi] := \|\nabla \varphi\|^2, \quad \varphi \in C_c^\infty(D_{r,s}). \quad (\text{A.10})$$

The self-adjoint operators $H_{r,s,n}$ and $H_{r,s,\infty}$ associated with (the closure of) these quadratic forms have compact resolvent. We have the following result:

A.2. Lemma. *With the above definitions and assumptions, we have*

$$\sup_s \|H_{r,s,n}^{-1} - H_{r,s,\infty}^{-1}\| \rightarrow 0, \quad n \rightarrow \infty. \quad (\text{A.11})$$

Proof. Obviously, we may restrict our attention to parameters s from a compact interval $J \subset \mathbb{R}$. Writing

$$f_n(s) := \|H_{r,s,n}^{-1} - H_{r,s,\infty}^{-1}\|, \quad s \in J, \quad (\text{A.12})$$

monotonicity and compactness imply that the functions f_n are continuous with $f_n(s) \rightarrow 0$ monotonically as $n \rightarrow \infty$, for each fixed s . Now the desired result follows by Dini's Theorem. \square

REFERENCES

- [AADH] S Alama, M Avellanda, PA Deift, and R Hempel, *On the existence of eigenvalues of a divergence form operator $A + \lambda B$ in a gap of $\sigma(A)$* , *Asymptotic Analysis* **8** (1994), 311–344
- [B] M Baake, *Solution of the coincidence problem in dimensions $d \leq 4$* , [arXiv:math/0605222v1](https://arxiv.org/abs/math/0605222v1) [[math.MG](https://arxiv.org/abs/math/0605222v1)]
- [CFS] IP Cornfield, SV Fomin, and YG Sinai, *Ergodic theory*, Springer, New York, 1982
- [DS] EB Davies and B Simon, *Scattering theory for systems with different spatial asymptotics on the left and right*, *Commun. Math. Phys.* **63** (1978), 277–301
- [E] MSP Eastham, *The Spectral Theory of Periodic Differential Equations*, Scottish Academic Press, Edinburgh-London, 1973
- [EKSchS] H Englisch, W Kirsch, M Schröder, and B Simon, *Random Hamiltonians ergodic in all but one direction*, *Commun. Math. Phys.* **128** (1990), 613–625
- [HH] R Hempel and I Herbst, *Strong magnetic fields, Dirichlet boundaries, and spectral gaps*, *Commun. Math. Phys.* **169** (1995), 237–259
- [HHK] R Hempel, A Hinz, and H Kalf, *On the essential spectrum of Schrödinger operators with spherically symmetric potentials*, *Math. Ann.* **277** (1987), 197–208
- [HK] R Hempel and M Kohlmann, *A variational approach to dislocation problems for periodic Schrödinger operators*, [arXiv:1009.3581v1](https://arxiv.org/abs/1009.3581v1) [[math-ph](https://arxiv.org/abs/1009.3581v1)]
- [HZh] I Herbst and Z Zhao, *Sobolev spaces, Kac-regularity, and the Feynman-Kac-formula*, *Seminar on Stochastic Processes* (Princeton, NJ), Birkhäuser, Boston 1988, pp. 171–191
- [K] T Kato, *Perturbation theory for linear operators*, Springer, New York, 1966
- [Ki] Ch Kittel, *Introduction to Solid State Physics*, Wiley, 2004
- [Kor1] E Korotyaev, *Lattice dislocations in a 1-dimensional model*, *Commun. Math. Phys.* **213** (2000), 471–489
- [Kor2] ———, *Schrödinger operators with a junction of two 1-dimensional periodic potentials*, *Asymptotic Anal.* **45** (2005), 73–97
- [KS] V Kostykin and R Schrader, *Regularity of the surface density of states*, *J. Funct. Anal.* **187** (2001), 227–246
- [Ku] P Kuchment, *Floquet Theory for Partial Differential Equations*, Birkhäuser, Basel-Boston-Berlin, 1993
- [RS-I] M Reed and B Simon, *Methods of Modern Mathematical Physics. Vol. I. Analysis of Operators*, Revised and enlarged edition, Academic Press, New York, 1979
- [RS-IV] M Reed and B Simon, *Methods of Modern Mathematical Physics. Vol. IV. Analysis of Operators*, Academic Press, New York, 1978

- [V] I Veselic, *Existence and regularity properties of the integrated density of states of random Schrödinger operators*, Springer Lecture Notes in Mathematics, vol. **1917**, Springer, Berlin, 2008
- [Z] P Zeiner, *Multiple CSLs for the body centered cubic lattice*. J Physics, Conference series **30** (2006), 163–167

INSTITUTE FOR COMPUTATIONAL MATHEMATICS, TECHNISCHE UNIVERSITÄT BRAUNSCHWEIG,
POCKELSTRASSE 14, 38106 BRAUNSCHWEIG, GERMANY

E-mail address: r.hempel@tu-bs.de

INSTITUTE FOR APPLIED MATHEMATICS, LEIBNIZ UNIVERSITÄT HANNOVER, WELFENGARTEN 1,
30167 HANNOVER, GERMANY

E-mail address: kohlmann@ifam.uni-hannover.de