
Inexact Newton regularization methods in Hilbert scales

Qinian Jin · Ulrich Tautenhahn

Abstract We consider a class of inexact Newton regularization methods for solving nonlinear inverse problems in Hilbert scales. Under certain conditions we obtain the order optimal convergence rate result.

1 Introduction

In this paper we consider the nonlinear inverse problems

$$F(x) = y, \quad (1.1)$$

where $F : D(F) \subset X \mapsto Y$ is a nonlinear Fréchet differentiable operator between two Hilbert spaces X and Y whose norms and inner products are denoted as $\|\cdot\|$ and (\cdot, \cdot) respectively. We assume that (1.1) has a solution x^\dagger in the domain $D(F)$ of F , i.e. $F(x^\dagger) = y$. We use $F'(x)$ to denote the Fréchet derivative of F at $x \in D(F)$ and $F'(x)^*$ the adjoint of $F'(x)$. A characteristic property of such problems is their ill-posedness in the sense that their solutions do not depend continuously on the data. Let y^δ be the only available approximation of y satisfying

$$\|y^\delta - y\| \leq \delta \quad (1.2)$$

with a given small noise level $\delta > 0$. Due to the ill-posedness, the regularization techniques should be employed to produce from y^δ a stable approximate solution of (1.1).

Many regularization methods have been considered in the last two decades. In particular, the nonlinear Landweber iteration [6], the Levenberg-Marquardt method [4,9], and the exponential Euler iteration [7] have been applied to solve nonlinear inverse problems. These methods take the form

$$x_{n+1} = x_n - g_{\alpha_n} (F'(x_n)^* F'(x_n)) F'(x_n)^* (F(x_n) - y^\delta), \quad (1.3)$$

Qinian Jin
Department of Mathematics, Virginia Tech, Blacksburg, VA 24060, USA
E-mail: qnjin@math.vt.edu

Ulrich Tautenhahn
Department of Mathematics, University of Applied Sciences Zittau/Görlitz, PO Box 1454, 02754 Zittau, Germany
E-mail: u.tautenhahn@hs-zigr.de

where x_0 is an initial guess of x^\dagger , $\{\alpha_n\}$ is a sequence of positive numbers, and $\{g_\alpha\}$ is a family of spectral filter functions. The scheme (1.3) can be derived by applying the linear regularization method defined by $\{g_\alpha\}$ to the equation

$$F'(x_n)(x - x_n) = y^\delta - F(x_n). \quad (1.4)$$

which follows from (1.1) by replacing y by y^δ and $F(x)$ by its linearization $F(x_n) + F'(x_n)(x - x_n)$ at x_n . It is easy to see that

$$F(x_n) - y^\delta + F'(x_n)(x_{n+1} - x_n) = r_{\alpha_n}(F'(x_n)F'(x_n)^*)(F(x_n) - y^\delta),$$

where

$$r_\alpha(\lambda) = 1 - \lambda g_\alpha(\lambda) \quad (1.5)$$

which is called the residual function associated with g_α . For well-posed problems where $F'(x_n)$ is invertible, usually one has $\|r_{\alpha_n}(F'(x_n)F'(x_n)^*)\| \leq \mu_n < 1$ and consequently

$$\|F(x_n) - y^\delta + F'(x_n)(x_{n+1} - x_n)\| \leq \mu_n \|F(x_n) - y^\delta\|. \quad (1.6)$$

Thus the methods belong to the class of inexact Newton methods [2]. For ill-posed problems, however, there only holds $\|r_{\alpha_n}(F'(x_n)F'(x_n)^*)\| \leq 1$ in general. In [4] the Levenberg-Marquardt scheme was considered with $\{\alpha_n\}$ chosen adaptively so that (1.6) holds and the discrepancy principle was used to terminate the iteration. The order optimal convergence rates were derived recently in [5]. The general methods (1.3) with $\{\alpha_n\}$ chosen adaptively to satisfy (1.6) were considered later in [14, 11], but only suboptimal convergence rates were derived in [15] and the convergence analysis is far from complete. On the other hand, one may consider the method (1.3) with $\{\alpha_n\}$ given a priori. This has been done for the Levenberg-Marquardt method in [9] and the exponential Euler method in [7] for instance.

In this paper we will consider the inexact Newton methods in Hilbert scales which are more general than (1.3). Let L be a densely defined self-adjoint strictly positive linear operator in X . For each $r \in \mathbb{R}$, we define X_r to be the completion of $\cap_{k=0}^{\infty} D(L^k)$ with respect to the Hilbert space norm

$$\|x\|_r := \|L^r x\|.$$

This family of Hilbert spaces $(X_r)_{r \in \mathbb{R}}$ is called the Hilbert scales generated by L . Let $x_0 \in D(F)$ be an initial guess of x^\dagger . The inexact Newton method in Hilbert scales defines the iterates $\{x_n\}$ by

$$x_{n+1} = x_n - g_{\alpha_n} (L^{-2s} F'(x_n)^* F'(x_n)) L^{-2s} F'(x_n)^* (F(x_n) - y^\delta), \quad (1.7)$$

where $s \in \mathbb{R}$ is a given number to be specified later, and $\{\alpha_n\}$ is an a priori given sequence of positive numbers with suitable properties. We will terminate the iteration by the discrepancy principle

$$\|F(x_{n_\delta}) - y^\delta\| \leq \tau \delta < \|F(x_n) - y^\delta\|, \quad 0 \leq n < n_\delta \quad (1.8)$$

with a given number $\tau > 1$ and consider the approximation property of x_{n_δ} to x^\dagger as $\delta \rightarrow 0$. We will establish for a large class of spectral filter functions $\{g_\alpha\}$ the order optimal convergence rates for the method defined by (1.7) and (1.8).

Regularization in Hilbert scales has been introduced in [12] for the linear Tikhonov regularization with the major aim to prevent the saturation effect. Such technique has been extended in various ways, in particular, a general class of regularization methods in Hilbert scales has been considered in [16] with the regularization parameter chosen by the Morozov's discrepancy principle. Regularization in Hilbert scales have

also been applied for solving nonlinear ill-posed problems. The nonlinear Tikhonov regularization in Hilbert scales has been considered in [10,3], a general continuous regularization scheme for nonlinear problems in Hilbert scales has been considered in [17], the general iteratively regularized Gauss-Newton methods in Hilbert scales has been considered in [8], and the nonlinear Landweber iteration in Hilbert scales has been considered in [13].

This paper is organized as follows. In Section 2 we first briefly review the relevant properties of Hilbert scales, and then formulate the necessary condition on $\{\alpha_n\}$, $\{g_\alpha\}$ and F together with some crucial consequences. In Section 3 we obtain the main result concerning the order optimal convergence property of the method given by (1.7) and (1.8). Finally we present in Section 4 several examples of the method (1.7) for which $\{g_\alpha\}$ satisfies the technical conditions in Section 2.

2 Assumptions

We first briefly review the relevant properties of the Hilbert scales $(X_r)_{r \in \mathbb{R}}$ generated by a densely defined self-adjoint strictly positive linear operator L in X , see [3]. It is well known that X_r is densely and continuously embedded into X_q for any $-\infty < q < r < \infty$, i.e.

$$\|x\|_q \leq \theta^{r-q} \|x\|_r, \quad x \in X_r, \quad (2.1)$$

where $\theta > 0$ is a constant such that

$$\|x\|^2 \leq \theta(Lx, x), \quad x \in D(L). \quad (2.2)$$

Moreover there holds the important interpolation inequality, i.e. for any $-\infty < p < q < r < \infty$ there holds for any $x \in X_r$ that

$$\|x\|_q \leq \|x\|_p^{\frac{r-q}{r-p}} \|x\|_r^{\frac{q-p}{r-p}}. \quad (2.3)$$

Let $T : X \mapsto Y$ be a bounded linear operator satisfying

$$m\|h\|_{-a} \leq \|Th\| \leq M\|h\|_{-a}, \quad h \in X$$

for some constants $M \geq m > 0$ and $a \geq 0$. Then the operator $A := TL^{-s} : X \mapsto Y$ is bounded for $s \geq -a$ and the adjoint of A is given by $A^* = L^{-s}T^*$, where $T^* : Y \mapsto X$ is the adjoint of T . Moreover, for any $|\nu| \leq 1$ there hold

$$R((A^*A)^{\nu/2}) = X_{\nu(a+s)} \quad (2.4)$$

and

$$\underline{c}(\nu)\|h\|_{-\nu(a+s)} \leq \|(A^*A)^{\nu/2}h\| \leq \bar{c}(\nu)\|h\|_{-\nu(a+s)} \quad (2.5)$$

on $D((A^*A)^{\nu/2})$, where

$$\underline{c}(\nu) := \min\{m^\nu, M^\nu\} \quad \text{and} \quad \bar{c}(\nu) = \max\{m^\nu, M^\nu\}.$$

If $g : [0, \|A\|^2] \mapsto \mathbb{R}$ is a continuous function, then

$$g(A^*A)L^s = L^s g(L^{-2s}T^*T). \quad (2.6)$$

In order to carry out the convergence analysis on the method defined by (1.7) and (1.8), we need to impose suitable conditions on $\{\alpha_n\}$, $\{g_\alpha\}$ and F . For the sequence $\{\alpha_n\}$ of positive numbers, we set

$$s_{-1} = 0, \quad s_n := \sum_{j=0}^n \frac{1}{\alpha_j}, \quad n = 0, 1, \dots \quad (2.7)$$

We will assume that there are constants $c_0 > 1$ and $c_1 > 0$ such that

$$\lim_{n \rightarrow \infty} s_n = \infty, \quad s_{n+1} \leq c_0 s_n \quad \text{and} \quad 0 < \alpha_n \leq c_1, \quad n = 0, 1, \dots. \quad (2.8)$$

We will also assume that, for each $\alpha > 0$, the function g_α is defined on $[0, 1]$ and satisfies the following structure condition, where \mathbb{C} denotes the complex plane.

Assumption 1 *For each $\alpha > 0$, the function*

$$\varphi_\alpha(\lambda) := g_\alpha(\lambda) - \frac{1}{\alpha + \lambda}$$

extends to a complex analytic function defined on a domain $D_\alpha \subset \mathbb{C}$ such that $[0, 1] \subset D_\alpha$, and there is a contour $\Gamma_\alpha \subset D_\alpha$ enclosing $[0, 1]$ such that

$$|z| \geq \frac{1}{2}\alpha \quad \text{and} \quad \frac{|z| + \lambda}{|z - \lambda|} \leq b_0, \quad \forall z \in \Gamma_\alpha, \alpha > 0 \quad \text{and} \quad \lambda \in [0, 1], \quad (2.9)$$

where b_0 is a constant independent of $\alpha > 0$. Moreover, there is a constant b_1 such that

$$\int_{\Gamma_\alpha} |\varphi_\alpha(z)| |dz| \leq b_1 \quad (2.10)$$

for all $0 < \alpha \leq c_1$.

By using the spectral integrals for self-adjoint operators, it follows easily from (2.9) in Assumption 1 that for any bounded linear operator A with $\|A\| \leq 1$ there holds

$$\|(zI - A^*A)^{-1}(A^*A)^\nu\| \leq \frac{b_0}{|z|^{1-\nu}} \quad (2.11)$$

for $z \in \Gamma_\alpha$ and $0 \leq \nu \leq 1$.

Moreover, since Assumption 1 implies $\varphi_\alpha(z)$ is analytic in D_α for each $\alpha > 0$, there holds the Riesz-Dunford formula (see [1])

$$\varphi_\alpha(A^*A) = \frac{1}{2\pi i} \int_{\Gamma_\alpha} \varphi_\alpha(z)(zI - A^*A)^{-1} dz$$

for any linear operator A satisfying $\|A\| \leq 1$.

Assumption 2 *Let $\{\alpha_n\}$ be a sequence of positive numbers, let $\{s_n\}$ be defined by (2.7). There is a constant $b_2 > 0$ such that*

$$0 \leq \lambda^\nu \prod_{k=j}^n r_{\alpha_k}(\lambda) \leq (s_n - s_{j-1})^{-\nu}, \quad (2.12)$$

$$0 \leq \lambda^\nu g_{\alpha_j}(\lambda) \prod_{k=j+1}^n r_{\alpha_k}(\lambda) \leq b_2 \frac{1}{\alpha_j} (s_n - s_{j-1})^{-\nu} \quad (2.13)$$

for $0 \leq \nu \leq 1$, $0 \leq \lambda \leq 1$ and $j = 0, 1, \dots, n$, where $r_\alpha(\lambda)$ is defined by (1.5).

In Section 4 we will give several important examples of $\{g_\alpha\}$ satisfying Assumptions 1 and 2. These examples of $\{g_\alpha\}$ include the ones arising from (iterated) Tikhonov regularization, asymptotical regularization, Landweber iteration and Lardy method.

Lemma 1 *The inequality (2.12) implies for $0 \leq \nu \leq 1$ and $\alpha > 0$ that*

$$0 \leq \lambda^\nu (\alpha + \lambda)^{-1} \prod_{k=j+1}^n r_{\alpha_k}(\lambda) \leq 2\alpha^{\nu-1} (1 + \alpha(s_n - s_j))^{-\nu} \quad (2.14)$$

for all $0 \leq \lambda \leq 1$ and $j = 0, 1, \dots, n$.

Proof For $0 \leq \nu \leq 1$ and $\alpha > 0$ it follows from (2.12) that

$$\begin{aligned} 0 \leq \lambda^\nu (\alpha + \lambda)^{-1} \prod_{k=j+1}^n r_{\alpha_k}(\lambda) &\leq \min \{ \alpha^{\nu-1}, \alpha^{-1} (s_n - s_j)^{-\nu} \} \\ &= \alpha^{\nu-1} \min \{ 1, \alpha^{-\nu} (s_n - s_j)^{-\nu} \} \\ &\leq 2^\nu \alpha^{\nu-1} (1 + \alpha (s_n - s_j))^{-\nu} \end{aligned}$$

for all $0 \leq \lambda \leq 1$ and $j = 0, 1, \dots, n$. \square

Assumption 3 (a) *There exist constants $a \geq 0$ and $0 < m \leq M < \infty$ such that*

$$m \|h\|_{-a} \leq \|F'(x)h\| \leq M \|h\|_{-a}, \quad h \in X$$

for all $x \in B_\rho(x^\dagger)$.

(b) *F is properly scaled so that $\|F'(x)L^{-s}\|_{X \rightarrow Y} \leq \min\{1, \sqrt{\alpha_0}\}$ for all $x \in B_\rho(x^\dagger)$, where $s \geq -a$.*

(c) *There exist $0 < \beta \leq 1$, $0 \leq b \leq a$ and $K_0 \geq 0$ such that*

$$\|F'(x)^* - F'(x^\dagger)^*\|_{Y \rightarrow X_b} \leq K_0 \|x - x^\dagger\|^\beta \quad (2.15)$$

for all $x \in B_\rho(x^\dagger)$.

The number a in condition (a) can be interpreted as the degree of ill-posedness of $F'(x)$ for $x \in B_\rho(x^\dagger)$. When F satisfies the condition

$$F'(x) = R_x F'(x^\dagger) \quad \text{and} \quad \|I - R_x\| \leq K_0 \|x - x^\dagger\|, \quad (2.16)$$

which has been verified in [6] for several nonlinear inverse problems, condition (a) is equivalent to

$$m \|h\|_{-a} \leq \|F'(x^\dagger)h\| \leq M \|h\|_{-a}, \quad h \in X$$

From (a) and (2.1) it follows for $s \geq -a$ that $\|F'(x)L^{-s}\|_{X \rightarrow Y} \leq M\theta^{a+s}$ for all $x \in B_\rho(x^\dagger)$. Thus $\|F'(x)L^{-s}\|_{X \rightarrow Y}$ is uniformly bounded over $B_\rho(x^\dagger)$. By multiplying (1.1) by a sufficiently small number, we may assume that F is properly scaled so that condition (b) is satisfied. Furthermore, condition (a) implies that $F'(x)^*$ maps Y into X_b for $b \leq a$ and $\|F'(x)^*\|_{Y \rightarrow X_b} \leq M\theta^{a-b}$ for all $x \in B_\rho(x^\dagger)$. Condition (c) says that $F'(x)^*$ is locally Hölder continuous around x^\dagger with exponent $0 < \beta \leq 1$ when considered as operators from Y to X_b . It is equivalent to

$$\|L^b[F'(x)^* - F'(x^\dagger)^*]\|_{Y \rightarrow X} \leq K_0 \|x - x^\dagger\|^\beta, \quad x \in B_\rho(x^\dagger)$$

or

$$\|[F'(x) - F'(x^\dagger)]L^b\|_{X \rightarrow Y} \leq K_0 \|x - x^\dagger\|^\beta, \quad x \in B_\rho(x^\dagger).$$

Condition (c) was used first in [13] for the convergence analysis of Landweber iteration in Hilbert scales. It is easy to see that when $b = 0$ and $\beta = 1$, this is exactly the Lipschitz condition on $F'(x)$. When F satisfies (2.16), (c) holds with $b = a$ and $\beta = 1$. In [13] it has been shown that (c) implies

$$\|F(x) - y - F'(x^\dagger)(x - x^\dagger)\| \leq K_0 \|x - x^\dagger\|^\beta \|x - x^\dagger\|_{-b} \quad (2.17)$$

which follows easily from the identity

$$F(x) - y - F'(x^\dagger)(x - x^\dagger) = \int_0^1 [F'(x^\dagger + t(x - x^\dagger)) - F'(x^\dagger)] L^b L^{-b}(x - x^\dagger) dt.$$

In this paper we will derive, under the above assumptions on $\{\alpha_n\}$, $\{g_\alpha\}$ and F , the rate of convergence of x_{n_δ} to x^\dagger as $\delta \rightarrow 0$ when $e_0 := x_0 - x^\dagger$ satisfies the smoothness condition

$$x_0 - x^\dagger \in X_\mu \quad \text{with} \quad \frac{a-b}{\beta} < \mu \leq b+2s, \quad (2.18)$$

where n_δ is the integer determined by the discrepancy principle (1.8) with $\tau > 1$.

The following consequence of the above assumptions on F and $\{g_\alpha\}$ plays a crucial role in the convergence analysis.

Lemma 2 *Let $\{g_\alpha\}$ satisfy Assumptions 1 and 2, let F satisfy Assumption 3, and let $\{\alpha_n\}$ be a sequence of positive numbers. Let $A = F'(x^\dagger)L^{-s}$ and for any $x \in B_\rho(x^\dagger)$ let $A_x = F'(x)L^{-s}$. Then for $-\frac{b+s}{2(a+s)} \leq \nu \leq 1/2$ there holds ¹*

$$\begin{aligned} & \left\| (A^*A)^\nu \prod_{k=j+1}^n r_{\alpha_k}(A^*A) [g_{\alpha_j}(A^*A)A^* - g_{\alpha_j}(A_x^*A_x)A_x^*] \right\| \\ & \lesssim \frac{1}{\alpha_j} (s_n - s_{j-1})^{-\nu - \frac{b+s}{2(a+s)}} K_0 \|x - x^\dagger\|^\beta \end{aligned}$$

for $j = 0, 1, \dots, n$.

Proof Let $\eta_\alpha(\lambda) = (\alpha + \lambda)^{-1}$ and $\varphi_\alpha(\lambda) = g_\alpha(\lambda) - (\alpha + \lambda)^{-1}$. We can write

$$(A^*A)^\nu \prod_{k=j+1}^n r_{\alpha_k}(A^*A) [g_{\alpha_j}(A^*A)A^* - g_{\alpha_j}(A_x^*A_x)A_x^*] = J_1 + J_2 + J_3,$$

where

$$\begin{aligned} J_1 &:= (A^*A)^\nu \prod_{k=j+1}^n r_{\alpha_k}(A^*A) g_{\alpha_j}(A^*A) [A^* - A_x^*], \\ J_2 &:= (A^*A)^\nu \prod_{k=j+1}^n r_{\alpha_k}(A^*A) [\eta_{\alpha_j}(A^*A) - \eta_{\alpha_j}(A_x^*A_x)] A_x^*, \\ J_3 &:= (A^*A)^\nu \prod_{k=j+1}^n r_{\alpha_k}(A^*A) [\varphi_{\alpha_j}(A^*A) - \varphi_{\alpha_j}(A_x^*A_x)] A_x^*. \end{aligned}$$

It suffices to show that the desired estimates hold for the norms of J_1 , J_2 and J_3 .

From (2.5), (2.13) in Assumption 2 and Assumption 3 it follows that

$$\begin{aligned} \|J_1\| &\lesssim \left\| (A^*A)^\nu \prod_{k=j+1}^n r_{\alpha_k}(A^*A) g_{\alpha_j}(A^*A) (A^*A)^{\frac{b+s}{2(a+s)}} \right\| \\ &\quad \times \left\| (A^*A)^{-\frac{b+s}{2(a+s)}} [A_x^* - A^*] \right\| \\ &\lesssim \sup_{0 \leq \lambda \leq 1} \left(\lambda^{\nu + \frac{b+s}{2(a+s)}} g_{\alpha_j}(\lambda) \prod_{k=j+1}^n r_{\alpha_k}(\lambda) \right) \|L^b[F'(x)^* - F'(x^\dagger)^*]\|_{Y \rightarrow X} \\ &\lesssim \frac{1}{\alpha_j} (s_n - s_{j-1})^{-\nu - \frac{b+s}{2(a+s)}} K_0 \|x - x^\dagger\|^\beta \end{aligned}$$

¹ Throughout this paper we will always use C to denote a generic constant independent of δ and n . We will also use the convention $\Phi \lesssim \Psi$ to mean that $\Phi \leq C\Psi$ for some generic constant C .

which is the desired estimate.

In order to estimate $\|J_2\|$, we note that

$$\begin{aligned} \eta_{\alpha_j}(A^*A) - \eta_{\alpha_j}(A_x^*A_x) &= (\alpha_j I + A^*A)^{-1}A^*(A_x - A)(\alpha_j I + A_x^*A_x)^{-1} \\ &\quad + (\alpha_j I + A^*A)^{-1}(A_x^* - A^*)A_x(\alpha_j I + A_x^*A_x)^{-1}. \end{aligned}$$

Therefore $J_2 = J_2^{(1)} + J_2^{(2)}$, where

$$\begin{aligned} J_2^{(1)} &= (A^*A)^\nu \prod_{k=j+1}^n r_{\alpha_k}(A^*A)(\alpha_j I + A^*A)^{-1}A^*(A_x - A)(\alpha_j I + A_x^*A_x)^{-1}A_x^*, \\ J_2^{(2)} &= (A^*A)^\nu \prod_{k=j+1}^n r_{\alpha_k}(A^*A)(\alpha_j I + A^*A)^{-1}(A_x^* - A^*)A_xA_x^*(\alpha_j I + A_x^*A_x)^{-1}. \end{aligned}$$

With the help of Assumption 3 and (2.5) we have for any $w \in Y$ that

$$\begin{aligned} \|(A_x - A)(\alpha_j I + A_x^*A_x)^{-1}A_x^*w\| &= \|[F'(x) - F'(x^\dagger)]L^bL^{-(b+s)}(\alpha_j I + A_x^*A_x)^{-1}A_x^*w\| \\ &\leq K_0\|x - x^\dagger\|^\beta \|(\alpha_j I + A_x^*A_x)^{-1}A_x^*w\|_{-(b+s)} \\ &\lesssim K_0\|x - x^\dagger\|^\beta \|(A_x^*A_x)^{\frac{b+s}{2(a+s)}}(\alpha_j I + A_x^*A_x)^{-1}A_x^*w\| \\ &\lesssim K_0\|x - x^\dagger\|^\beta \alpha_j^{-\frac{1}{2} + \frac{b+s}{2(a+s)}} \|w\|. \end{aligned}$$

This implies

$$\|(A_x - A)(\alpha_j I + A_x^*A_x)^{-1}A_x^*\| \lesssim K_0\|x - x^\dagger\|^\beta \alpha_j^{-\frac{1}{2} + \frac{b+s}{2(a+s)}}. \quad (2.19)$$

Thus, by using Lemma 1, we derive

$$\begin{aligned} \|J_2^{(1)}\| &\leq \sup_{0 \leq \lambda \leq 1} \left(\lambda^{\nu + \frac{1}{2}} (\alpha_j + \lambda)^{-1} \prod_{k=j+1}^n r_{\alpha_k}(\lambda) \right) \|(A_x - A)(\alpha_j I + A_x^*A_x)^{-1}A_x^*\| \\ &\lesssim \alpha_j^{\nu - 1 + \frac{b+s}{2(a+s)}} (1 + \alpha_j(s_n - s_j))^{-\nu - \frac{1}{2}} K_0\|x - x^\dagger\|^\beta. \end{aligned}$$

By using Assumption 3, Lemma 1 and a similar argument in estimating J_1 we can derive

$$\begin{aligned} \|J_2^{(2)}\| &\lesssim \sup_{0 \leq \lambda \leq 1} \left(\lambda^{\nu + \frac{b+s}{2(a+s)}} (\alpha_j + \lambda)^{-1} \prod_{k=j+1}^n r_{\alpha_k}(\lambda) \right) \|L^b[F'(x)^* - F'(x^\dagger)^*]\|_{Y \rightarrow X} \\ &\lesssim \alpha_j^{\nu - 1 + \frac{b+s}{2(a+s)}} (1 + \alpha_j(s_n - s_j))^{-\nu - \frac{b+s}{2(a+s)}} K_0\|x - x^\dagger\|^\beta. \end{aligned}$$

Combining the above estimates on $J_2^{(1)}$ and $J_2^{(2)}$ and noting $\frac{b+s}{2(a+s)} \leq \frac{1}{2}$, it follows that

$$\begin{aligned} \|J_2\| &\lesssim \alpha_j^{\nu - 1 + \frac{b+s}{2(a+s)}} (1 + \alpha_j(s_n - s_j))^{-\nu - \frac{b+s}{2(a+s)}} K_0\|x - x^\dagger\|^\beta \\ &= \frac{1}{\alpha_j} (s_n - s_{j-1})^{-\nu - \frac{b+s}{2(a+s)}} K_0\|x - x^\dagger\|^\beta. \end{aligned}$$

It remains to estimate J_3 . Since Assumption 1 implies that $\varphi_{\alpha_j}(z)$ is analytic in D_{α_j} , we have from the Riesz-Dunford formula that

$$J_3 = \frac{1}{2\pi i} \int_{\Gamma_{\alpha_j}} \varphi_{\alpha_j}(z) T_j(z) dz, \quad (2.20)$$

where

$$T_j(z) := (A^*A)^\nu \prod_{k=j+1}^n r_{\alpha_k}(A^*A) [(zI - A^*A)^{-1} - (zI - A_x^*A_x)^{-1}] A_x^*.$$

We can write $T_j(z) = T_j^{(1)}(z) + T_j^{(2)}(z)$, where

$$\begin{aligned} T_j^{(1)}(z) &:= (A^*A)^\nu \prod_{k=j+1}^n r_{\alpha_k}(A^*A) (zI - A^*A)^{-1} A^*(A - A_x)(zI - A_x^*A_x)^{-1} A_x^*, \\ T_j^{(2)}(z) &:= (A^*A)^\nu \prod_{k=j+1}^n r_{\alpha_k}(A^*A) (zI - A^*A)^{-1} (A^* - A_x^*)A_x A_x^* (zI - A_x A_x^*)^{-1}. \end{aligned}$$

We will estimate the norms of $T_j^{(1)}(z)$ and $T_j^{(2)}(z)$ for $z \in \Gamma_{\alpha_j}$. With the help of Assumption 3, (2.5) and (2.11), similar to the derivation of (2.19) we have

$$\|(A - A_x)(zI - A_x^*A_x)^{-1} A_x^*\| \lesssim K_0 \|x - x^\dagger\|^\beta |z|^{-\frac{1}{2} + \frac{b+s}{2(a+s)}}.$$

Since $|z| \geq \alpha_j/2$ and $|z - \lambda|^{-1} \leq b_0(|z| + \lambda)^{-1}$ for $z \in \Gamma_{\alpha_j}$, we have from (2.14) in Lemma 1 that

$$\begin{aligned} \|T_j^{(1)}(z)\| &\lesssim K_0 \|x - x^\dagger\|^\beta |z|^{-\frac{1}{2} + \frac{b+s}{2(a+s)}} \sup_{0 \leq \lambda \leq 1} \left(\lambda^{\nu + \frac{1}{2}} |z - \lambda|^{-1} \prod_{k=j+1}^n r_{\alpha_k}(\lambda) \right) \\ &\lesssim K_0 \|x - x^\dagger\|^\beta |z|^{-\frac{1}{2} + \frac{b+s}{2(a+s)}} \sup_{0 \leq \lambda \leq 1} \left(\lambda^{\nu + \frac{1}{2}} (|z| + \lambda)^{-1} \prod_{k=j+1}^n r_{\alpha_k}(\lambda) \right) \\ &\lesssim K_0 \|x - x^\dagger\|^\beta |z|^{\nu - 1 + \frac{b+s}{2(a+s)}} (1 + (s_n - s_j)|z|)^{-\nu - 1/2} \\ &\lesssim K_0 \|x - x^\dagger\|^\beta \alpha_j^{\nu - 1 + \frac{b+s}{2(a+s)}} (1 + (s_n - s_j)\alpha_j)^{-\nu - 1/2}. \end{aligned}$$

Next, by using (2.14) in Lemma 1, (2.5), Assumption 3(a) and (2.11), we have for $z \in \Gamma_{\alpha_j}$ that

$$\begin{aligned} \|T_j^{(2)}(z)\| &\leq \left\| (A^*A)^\nu \prod_{k=j+1}^n r_{\alpha_k}(A^*A) (zI - A^*A)^{-1} (A^*A)^{\frac{b+s}{2(a+s)}} \right\| \\ &\quad \times \left\| (A^*A)^{-\frac{b+s}{2(a+s)}} (A^* - A_x^*)A_x A_x^* (zI - A_x A_x^*)^{-1} \right\| \\ &\lesssim \sup_{0 \leq \lambda \leq 1} \left(\lambda^{\nu + \frac{b+s}{2(a+s)}} |z - \lambda|^{-1} \prod_{k=j+1}^n r_{\alpha_k}(\lambda) \right) \|L^b(F'(x^\dagger)^* - F'(x)^*)\| \\ &\lesssim K_0 \|x - x^\dagger\|^\beta \sup_{0 \leq \lambda \leq 1} \left(\lambda^{\nu + \frac{b+s}{2(a+s)}} (|z| + \lambda)^{-1} \prod_{k=j+1}^n r_{\alpha_k}(\lambda) \right) \\ &\lesssim K_0 \|x - x^\dagger\|^\beta |z|^{\nu - 1 + \frac{b+s}{2(a+s)}} (1 + (s_n - s_j)|z|)^{-\nu - \frac{b+s}{2(a+s)}} \\ &\lesssim K_0 \|x - x^\dagger\|^\beta \alpha_j^{\nu - 1 + \frac{b+s}{2(a+s)}} (1 + (s_n - s_j)\alpha_j)^{-\nu - \frac{b+s}{2(a+s)}}. \end{aligned}$$

Combining the above estimates on $T_j^{(1)}(z)$ and $T_j^{(2)}(z)$ and noting $\frac{b+s}{2(a+s)} \leq \frac{1}{2}$, it follows for $z \in \Gamma_{\alpha_j}$ that

$$\begin{aligned} \|T_j(z)\| &\lesssim K_0 \|x - x^\dagger\|^\beta \alpha_j^{\nu - 1 + \frac{b+s}{2(a+s)}} (1 + (s_n - s_j)\alpha_j)^{-\nu - \frac{b+s}{2(a+s)}} \\ &= \frac{1}{\alpha_j} (s_n - s_{j-1})^{-\nu - \frac{b+s}{2(a+s)}} K_0 \|x - x^\dagger\|^\beta \end{aligned}$$

Therefore, it follows from (2.20) and Assumption 1 that

$$\begin{aligned} \|J_3\| &\lesssim \frac{1}{\alpha_j} (s_n - s_{j-1})^{-\nu - \frac{b+s}{2(a+s)}} K_0 \|x - x^\dagger\|^\beta \int_{\Gamma_{\alpha_j}} |\varphi_{\alpha_j}(z)| |dz| \\ &\lesssim \frac{1}{\alpha_j} (s_n - s_{j-1})^{-\nu - \frac{b+s}{2(a+s)}} K_0 \|x - x^\dagger\|^\beta. \end{aligned}$$

The proof is therefore complete. \square

3 Convergence analysis

We begin with the following lemma.

Lemma 3 *Let $\{\alpha_n\}$ be a sequence of positive numbers satisfying $\alpha_n \leq c_1$, and let s_n be defined by (2.7). Let $p \geq 0$ and $q \geq 0$ be two numbers. Then we have*

$$\sum_{j=0}^n \frac{1}{\alpha_j} (s_n - s_{j-1})^{-p} s_j^{-q} \leq C_0 s_n^{1-p-q} \begin{cases} 1, & \max\{p, q\} < 1, \\ \log(1 + s_n), & \max\{p, q\} = 1, \\ s_n^{\max\{p, q\}-1}, & \max\{p, q\} > 1, \end{cases}$$

where C_0 is a constant depending only on c_1 , p and q .

Proof This result is essentially contained in [5, Lemma 4.3] and its proof. For completeness, we include here the proof with a simplified argument. We first rewrite

$$\sum_{j=0}^n \frac{1}{\alpha_j} (s_n - s_{j-1})^{-p} s_j^{-q} = s_n^{1-p-q} \sum_{j=0}^n \frac{1}{\alpha_j s_n} \left(1 - \frac{s_{j-1}}{s_n}\right)^{-p} \left(\frac{s_j}{s_n}\right)^{-q}.$$

Observe that when $0 \leq s_{j-1}/s_n \leq 1/2$ we have

$$\left(1 - \frac{s_{j-1}}{s_n}\right)^{-p} \left(\frac{s_j}{s_n}\right)^{-q} \leq 2^p \left(\frac{s_j}{s_n}\right)^{-q}$$

while when $s_{j-1}/s_n \geq 1/2$ we have

$$\left(1 - \frac{s_{j-1}}{s_n}\right)^{-p} \left(\frac{s_j}{s_n}\right)^{-q} \leq 2^q \left(1 - \frac{s_{j-1}}{s_n}\right)^{-p}.$$

Consequently there holds with $C_{p,q} = \max\{2^p, 2^q\}$

$$\begin{aligned} &\sum_{j=0}^n \frac{1}{\alpha_j} (s_n - s_{j-1})^{-p} s_j^{-q} \\ &\leq C_{p,q} s_n^{1-p-q} \left(\sum_{j=0}^n \frac{1}{\alpha_j s_n} \left(\frac{s_j}{s_n}\right)^{-q} + \sum_{j=0}^n \frac{1}{\alpha_j s_n} \left(1 - \frac{s_{j-1}}{s_n}\right)^{-p} \right). \end{aligned} \quad (3.1)$$

Note that $s_j - s_{j-1} = 1/\alpha_j$, we have with $h = \frac{1}{2\alpha_0 s_n}$

$$\begin{aligned} \int_{s_0/s_n-h}^1 t^{-q} dt &= \sum_{j=1}^n \int_{s_{j-1}/s_n}^{s_j/s_n} t^{-q} dt + \int_{s_0/s_n-h}^{s_0/s_n} t^{-q} dt \\ &\geq \sum_{j=1}^n \left(\frac{s_j}{s_n}\right)^{-q} \frac{s_j - s_{j-1}}{s_n} + \frac{1}{2\alpha_0 s_n} \left(\frac{s_0}{s_n}\right)^{-q} \\ &\geq \frac{1}{2} \sum_{j=0}^n \frac{1}{\alpha_j s_n} \left(\frac{s_j}{s_n}\right)^{-q}. \end{aligned}$$

Therefore

$$\sum_{j=0}^n \frac{1}{\alpha_j s_n} \left(\frac{s_j}{s_n} \right)^{-q} \leq 2 \int_{s_0/s_n-h}^1 t^{-q} dt \leq \begin{cases} \frac{2}{1-q}, & q < 1, \\ 2 \log(2\alpha_0 s_n), & q = 1, \\ \frac{2}{q-1} (2\alpha_0 s_n)^{q-1}, & q > 1. \end{cases} \quad (3.2)$$

By a similar argument we have with $h = \frac{1}{2\alpha_n s_n}$

$$\sum_{j=0}^n \frac{1}{\alpha_j s_n} \left(1 - \frac{s_{j-1}}{s_n} \right)^{-p} \leq 2 \int_0^{\frac{s_{n-1}}{s_n}+h} (1-t)^{-p} dt \leq \begin{cases} \frac{2}{1-p}, & p < 1, \\ 2 \log(2\alpha_n s_n), & p = 1, \\ \frac{2}{p-1} (2\alpha_n s_n)^{p-1}, & p > 1. \end{cases} \quad (3.3)$$

Combining (3.1), (3.2) and (3.3) and using the condition $\alpha_n \leq c_1$, we obtain the desired inequalities. \square

In order to derive the necessary estimates on $x_n - x^\dagger$, we need some useful identities. For simplicity of presentation, we set

$$e_n := x_n - x^\dagger, \quad A := F'(x^\dagger)L^{-s} \quad \text{and} \quad A_n := F'(x_n)L^{-s}.$$

It follows from (1.7) and (2.6) that

$$e_{n+1} = e_n - L^{-s} g_{\alpha_n} (A_n^* A_n) A_n^* (F(x_n) - y^\delta).$$

Let

$$u_n := F(x_n) - y - F'(x^\dagger)(x_n - x^\dagger).$$

Then we can write

$$\begin{aligned} e_{n+1} &= e_n - L^{-s} g_{\alpha_n} (A^* A) A^* (F(x_n) - y^\delta) \\ &\quad - L^{-s} [g_{\alpha_n} (A_n^* A_n) A_n^* - g_{\alpha_n} (A^* A) A^*] (F(x_n) - y^\delta) \\ &= L^{-s} r_{\alpha_n} (A^* A) L^s e_n - L^{-s} g_{\alpha_n} (A^* A) A^* (y - y^\delta + u_n) \\ &\quad - L^{-s} [g_{\alpha_n} (A_n^* A_n) A_n^* - g_{\alpha_n} (A^* A) A^*] (F(x_n) - y^\delta). \end{aligned} \quad (3.4)$$

By telescoping (3.4) we can obtain

$$\begin{aligned} e_{n+1} &= L^{-s} \prod_{j=0}^n r_{\alpha_j} (A^* A) L^s e_0 \\ &\quad - L^{-s} \sum_{j=0}^n \prod_{k=j+1}^n r_{\alpha_k} (A^* A) g_{\alpha_j} (A^* A) A^* (y - y^\delta + u_j) \\ &\quad - L^{-s} \sum_{j=0}^n \prod_{k=j+1}^n r_{\alpha_k} (A^* A) [g_{\alpha_j} (A_j^* A_j) A_j^* - g_{\alpha_j} (A^* A) A^*] (F(x_j) - y^\delta). \end{aligned} \quad (3.5)$$

By multiplying (3.5) by $T := F'(x^\dagger)$ and noting that $A = TL^{-s}$ and

$$I - \sum_{j=0}^n \prod_{k=j+1}^n r_{\alpha_k} (AA^*) g_{\alpha_j} (AA^*) AA^* = \prod_{j=0}^n r_{\alpha_j} (AA^*),$$

we can obtain

$$\begin{aligned}
& Te_{n+1} - y^\delta + y \\
&= A \prod_{j=0}^n r_{\alpha_j}(A^*A)L^s e_0 + \prod_{j=0}^n r_{\alpha_j}(AA^*)(y - y^\delta) \\
&\quad - \sum_{j=0}^n \prod_{k=j+1}^n r_{\alpha_k}(AA^*)g_{\alpha_j}(AA^*)AA^*u_j \\
&\quad - \sum_{j=0}^n A \prod_{k=j+1}^n r_{\alpha_k}(A^*A) [g_{\alpha_j}(A_j^*A_j)A_j^* - g_{\alpha_j}(A^*A)A^*] (F(x_j) - y^\delta). \quad (3.6)
\end{aligned}$$

Based on (3.5) and (3.6) we will derive the order optimal convergence rate of x_{n_δ} to x^\dagger when $e_0 := x_0 - x^\dagger$ satisfies the smoothness condition (2.18). Under such condition we have $L^s e_0 \in X_{\mu-s}$ and $|\frac{\mu-s}{a+s}| \leq 1$. Thus, with the help of Assumption 3(a), it follows from (2.4) and (2.5) that there exists $\omega \in X$ such that

$$L^s e_0 = (A^*A)^{\frac{\mu-s}{2(a+s)}} \omega \quad \text{and} \quad c_2 \|\omega\| \leq \|e_0\|_\mu \leq c_3 \|\omega\| \quad (3.7)$$

for some generic constants $c_3 \geq c_2 > 0$. We will first derive the crucial estimates on $\|e_n\|_\mu$ and $\|Te_n\|$. To this end, we introduce the integer \tilde{n}_δ satisfying

$$s_{\tilde{n}_\delta}^{-\frac{a+\mu}{2(a+s)}} \leq \frac{(\tau-1)\delta}{2c_0\|\omega\|} < s_n^{-\frac{a+\mu}{2(a+s)}}, \quad 0 \leq n < \tilde{n}_\delta, \quad (3.8)$$

where $c_0 > 1$ is the constant appearing in (2.8). Such \tilde{n}_δ is well-defined since $s_n \rightarrow \infty$ as $n \rightarrow \infty$.

Proposition 1 *Let F satisfy Assumptions 3, let $\{g_\alpha\}$ satisfy Assumptions 1 and 2, and let $\{\alpha_n\}$ be a sequence of positive numbers satisfying (2.8). If $e_0 \in X_\mu$ for some $(a-b)/\beta < \mu \leq b+2s$ and if $K_0\|\omega\|^\beta$ is suitably small, then there exists a generic constant $C_* > 0$ such that*

$$\|e_n\|_\mu \leq C_* \|\omega\| \quad \text{and} \quad \|Te_n\| \leq C_* s_n^{-\frac{a+\mu}{2(a+s)}} \|\omega\| \quad (3.9)$$

and

$$\|Te_n - y^\delta + y\| \leq (c_0 + C_* K_0 \|\omega\|^\beta) s_n^{-\frac{a+\mu}{2(a+s)}} \|\omega\| + \delta \quad (3.10)$$

for all $0 \leq n \leq \tilde{n}_\delta$.

Proof We will show (3.9) by induction. By using (3.7) and $\|A\| \leq \sqrt{\alpha_0}$ we have

$$\|Te_0\| = \|AL^s e_0\| = \|(A^*A)^{1/2} L^s e_0\| = \|(A^*A)^{\frac{a+\mu}{2(a+s)}} \omega\| \leq \alpha_0^{\frac{a+\mu}{2(a+s)}} \|\omega\|.$$

This together with (3.7) shows (3.9) for $n = 0$ if $C_* \geq \max\{1, c_3\}$. Next we assume that (3.9) holds for all $0 \leq n \leq l$ for some $l < \tilde{n}_\delta$ and we are going to show (3.9) holds for $n = l+1$.

With the help of (2.5) and (3.7) we can derive from (3.5) that

$$\begin{aligned}
& \|e_{l+1}\|_\mu \\
& \lesssim \left\| \prod_{j=0}^l r_{\alpha_j}(A^*A)\omega \right\| + \left\| \sum_{j=0}^l (AA^*)^{\frac{a+2s-\mu}{2(a+s)}} g_{\alpha_j}(AA^*) \prod_{k=j+1}^l r_{\alpha_k}(AA^*)(y - y^\delta + u_j) \right\| \\
& + \left\| \sum_{j=0}^l (A^*A)^{\frac{s-\mu}{2(a+s)}} \prod_{k=j+1}^l r_{\alpha_k}(A^*A) [g_{\alpha_j}(A_j^*A_j)A_j^* - g_{\alpha_j}(A^*A)A^*] (F(x_j) - y^\delta) \right\|.
\end{aligned}$$

Since $(a-b)/\beta < \mu \leq b+2s$ and $0 \leq b \leq a$, we have

$$0 \leq \frac{a+2s-\mu}{2(a+s)} < 1 \quad \text{and} \quad -\frac{b+s}{2(a+s)} \leq \frac{s-\mu}{2(a+s)} < \frac{1}{2}.$$

Thus we may use Assumption 2 and Lemma 2 to conclude

$$\begin{aligned} \|e_{l+1}\|_\mu &\lesssim \|\omega\| + \sum_{j=0}^l \frac{1}{\alpha_j} (s_l - s_{j-1})^{-\frac{a+2s-\mu}{2(a+s)}} (\delta + \|u_j\|) \\ &\quad + \sum_{j=0}^l \frac{1}{\alpha_j} (s_l - s_{j-1})^{-\frac{b+2s-\mu}{2(a+s)}} K_0 \|e_j\|^\beta \|F(x_j) - y^\delta\|. \end{aligned} \quad (3.11)$$

Moreover, by using (3.7), Assumption 2 and Lemma 2, we have from (3.6) that

$$\begin{aligned} \|Te_{l+1} - y^\delta + y\| &\leq s_l^{-\frac{a+\mu}{2(a+s)}} \|\omega\| + \delta + b_2 \sum_{j=0}^l \frac{1}{\alpha_j} (s_l - s_{j-1})^{-1} \|u_j\| \\ &\quad + c_4 \sum_{j=0}^l \frac{1}{\alpha_j} (s_l - s_{j-1})^{-\frac{b+a+2s}{2(a+s)}} K_0 \|e_j\|^\beta \|F(x_j) - y^\delta\|, \end{aligned} \quad (3.12)$$

where $c_4 > 0$ is a generic constant.

By using the interpolation inequality (2.3), Assumption 3(a) and the induction hypotheses, it follows for all $0 \leq j \leq l$ that

$$\|e_j\| \leq \|e_j\|_{-a}^{\frac{\mu}{a+\mu}} \|e_j\|_\mu^{\frac{a}{a+\mu}} \lesssim \|Te_j\|_{-a}^{\frac{\mu}{a+\mu}} \|e_j\|_\mu^{\frac{a}{a+\mu}} \lesssim \|\omega\| s_j^{-\frac{\mu}{2(a+s)}}. \quad (3.13)$$

With the help of (2.17) and the interpolation inequality (2.3), we have

$$\|u_j\| \leq K_0 \|e_j\|^\beta \|e_j\|_{-b} \leq K_0 \|e_j\|_{-a}^{\frac{b+\mu+\mu\beta}{a+\mu}} \|e_j\|_\mu^{\frac{a+\alpha\beta-b}{a+\mu}}. \quad (3.14)$$

We then obtain from Assumption 3(a) and the induction hypotheses that

$$\|u_j\| \lesssim K_0 \|Te_j\|_{-a}^{\frac{b+\mu+\mu\beta}{a+\mu}} \|e_j\|_\mu^{\frac{a+\alpha\beta-b}{a+\mu}} \lesssim K_0 \|\omega\|^{1+\beta} s_j^{-\frac{b+\mu+\mu\beta}{2(a+s)}}. \quad (3.15)$$

On the other hand, since (2.1) and the induction hypotheses implies

$$\|e_j\|_{-a} \lesssim \|e_j\|_\mu \lesssim \|\omega\|, \quad 0 \leq j \leq l$$

and since $\mu > (a-b)/\beta$, we have from (3.14) and Assumption 3(a) that

$$\|u_j\| \lesssim K_0 \|e_j\|_{-a} \|e_j\|_{-a}^{\frac{b-a+\mu\beta}{a+\mu}} \|e_j\|_\mu^{\frac{a+\alpha\beta-b}{a+\mu}} \lesssim K_0 \|\omega\|^\beta \|Te_j\|. \quad (3.16)$$

Therefore, by using the fact

$$\delta \leq \frac{2c_0}{\tau-1} \|\omega\| s_j^{-\frac{a+\mu}{2(a+s)}}, \quad 0 \leq j \leq l \quad (3.17)$$

and the induction hypotheses we have

$$\|F(x_j) - y^\delta\| \leq \delta + \|Te_j\| + \|u_j\| \lesssim \|\omega\| s_j^{-\frac{a+\mu}{2(a+s)}}. \quad (3.18)$$

In view of the estimates (3.13), (3.15), (3.18) and the inequality

$$\sum_{j=0}^l \frac{1}{\alpha_j} (s_l - s_{j-1})^{-\frac{a+2s-\mu}{2(a+s)}} \lesssim s_l^{\frac{a+\mu}{2(a+s)}}$$

which follows from Lemma 3, we have from (3.11) and (3.12) that

$$\begin{aligned} \|e_{l+1}\|_\mu &\leq c_5 \|\omega\| + c_5 s_l^{\frac{a+\mu}{2(a+s)}} \delta \\ &\quad + CK_0 \|\omega\|^{1+\beta} \sum_{j=0}^l \frac{1}{\alpha_j} (s_l - s_{j-1})^{-\frac{a+2s-\mu}{2(a+s)}} s_j^{-\frac{b+\mu+\mu\beta}{2(a+s)}} \\ &\quad + CK_0 \|\omega\|^{1+\beta} \sum_{j=0}^l \frac{1}{\alpha_j} (s_l - s_{j-1})^{-\frac{b+2s-\mu}{2(a+s)}} s_j^{-\frac{a+\mu+\mu\beta}{2(a+s)}} \end{aligned}$$

and

$$\begin{aligned} \|Te_{l+1} - y^\delta + y\| &\leq \|\omega\| s_l^{-\frac{a+\mu}{2(a+s)}} + \delta \\ &\quad + CK_0 \|\omega\|^{1+\beta} \sum_{j=0}^l \frac{1}{\alpha_j} (s_l - s_{j-1})^{-1} s_j^{-\frac{b+\mu+\mu\beta}{2(a+s)}} \\ &\quad + CK_0 \|\omega\|^{1+\beta} \sum_{j=0}^l \frac{1}{\alpha_j} (s_l - s_{j-1})^{-\frac{b+a+2s}{2(a+s)}} s_j^{-\frac{a+\mu+\mu\beta}{2(a+s)}}, \end{aligned}$$

where c_5 and C are two positive generic constants.

With the help of Lemma 3, $\mu > (a-b)/\beta$, (3.17) and (2.8) we have

$$\|e_{l+1}\|_\mu \leq \left(c_5 + \frac{2}{\tau-1} c_0 c_5 + CK_0 \|\omega\|^\beta \right) \|\omega\|,$$

and

$$\begin{aligned} \|Te_{l+1} - y^\delta + y\| &\leq \delta + (1 + CK_0 \|\omega\|^\beta) \|\omega\| s_l^{-\frac{a+\mu}{2(a+s)}} \\ &\leq \delta + c_0 (1 + CK_0 \|\omega\|^\beta) \|\omega\| s_{l+1}^{-\frac{a+\mu}{2(a+s)}}. \end{aligned} \quad (3.19)$$

Consequently $\|e_{l+1}\|_\mu \leq C_* \|\omega\|$ if $C_* \geq 2c_5 + \frac{2}{\tau-1} c_0 c_5$ and $K_0 \|\omega\|^\beta$ is suitably small. Moreover, from (3.19), (3.17) and (2.8) we also have

$$\begin{aligned} \|Te_{l+1}\| &\leq 2\delta + c_0 (1 + CK_0 \|\omega\|^\beta) \|\omega\| s_{l+1}^{-\frac{a+\mu}{2(a+s)}} \\ &\leq \left(\frac{4c_0^2}{\tau-1} + c_0 + CK_0 \|\omega\|^\beta \right) \|\omega\| s_{l+1}^{-\frac{a+\mu}{2(a+s)}} \\ &\leq C_* \|\omega\| s_{l+1}^{-\frac{a+\mu}{2(a+s)}} \end{aligned}$$

if $C_* \geq 2c_0 + \frac{4c_0^2}{\tau-1}$ and $K_0 \|\omega\|^\beta$ is suitably small. We therefore complete the proof of (3.9). In the meanwhile, (3.19) gives the proof of (3.10). \square

From Proposition 1 and its proof it follows that $x_n \in B_\rho(x^\dagger)$ for $0 \leq n \leq \tilde{n}_\delta$ if $\|\omega\|$ is sufficiently small. Furthermore, from (3.15) and (3.16) we have

$$\|F(x_n) - y - Te_n\| \lesssim K_0 \|\omega\|^{1+\beta} s_n^{-\frac{b+\mu+\mu\beta}{2(a+s)}} \quad (3.20)$$

and

$$\|F(x_n) - y - Te_n\| \lesssim K_0 \|\omega\|^\beta \|Te_n\| \quad (3.21)$$

for $0 \leq n \leq \tilde{n}_\delta$.

In the following we will show that $n_\delta \leq \tilde{n}_\delta$ for the integer n_δ defined by (1.8) with $\tau > 1$. Consequently, the method given by (1.7) and (1.8) is well-defined.

Lemma 4 *Let all the conditions in Proposition 1 hold. Let $\tau > 1$ be a given number. If $e_0 \in X_\mu$ for some $(a-b)/\beta < \mu \leq b+2s$ and if $K_0\|e_0\|_\mu^\beta$ is suitably small, then the discrepancy principle (1.8) defines a finite integer n_δ satisfying $n_\delta \leq \tilde{n}_\delta$.*

Proof From Proposition 1, (3.20) and $\mu > (a-b)/\beta$ it follows for $0 \leq n \leq \tilde{n}_\delta$ that

$$\begin{aligned} \|F(x_n) - y^\delta\| &\leq \|F(x_n) - y - Te_n\| + \|Te_n - y^\delta + y\| \\ &\leq CK_0\|\omega\|^{1+\beta} s_n^{-\frac{b+\mu+\mu\beta}{2(a+s)}} + (c_0 + CK_0\|\omega\|^\beta) s_n^{-\frac{a+\mu}{2(a+s)}} \|\omega\| + \delta \\ &\leq (c_0 + CK_0\|\omega\|^\beta) s_n^{-\frac{a+\mu}{2(a+s)}} \|\omega\| + \delta. \end{aligned}$$

By setting $n = \tilde{n}_\delta$ in the above inequality and using the definition of \tilde{n}_δ we obtain

$$\|F(x_{\tilde{n}_\delta}) - y^\delta\| \leq \left(1 + \frac{\tau-1}{2} + CK_0\|\omega\|^\beta\right) \delta \leq \tau\delta$$

if $K_0\|\omega\|^\beta$ is suitably small. According to the definition of n_δ we have $n_\delta \leq \tilde{n}_\delta$. \square

Now we are ready to prove the main result concerning the order optimal convergence rates for the method defined by (1.7) and (1.8) with $\tau > 1$.

Theorem 1 *Let F satisfy Assumptions 3, let $\{g_\alpha\}$ satisfy Assumptions 1 and 2, and let $\{\alpha_n\}$ be a sequence of positive numbers satisfying (2.8). If $e_0 \in X_\mu$ for some $(a-b)/\beta < \mu \leq b+2s$ and if $K_0\|e_0\|_\mu^\beta$ is suitably small, then for all $r \in [-a, \mu]$ there holds*

$$\|x_{n_\delta} - x^\dagger\|_r \leq C\|e_0\|_\mu^{\frac{a+r}{a+\mu}} \delta^{\frac{\mu-r}{a+\mu}}$$

for the integer n_δ determined by the discrepancy principle (1.8) with $\tau > 1$, where $C > 0$ is a generic constant.

Proof It follows from (3.21) that if $K_0\|\omega\|^\beta$ is suitably small then

$$\|F(x_n) - y - Te_n\| \leq \frac{1}{2}\|Te_n\|$$

which implies $\|Te_n\| \leq 2\|F(x_n) - y\|$ for $0 \leq n \leq \tilde{n}_\delta$. Since Lemma 4 implies $n_\delta \leq \tilde{n}_\delta$, it follows from Assumption 3(a) and the definition of n_δ that

$$\|e_{n_\delta}\|_{-a} \leq \frac{1}{m}\|Te_{n_\delta}\| \leq \frac{2}{m}(\|F(x_{n_\delta}) - y^\delta\| + \delta) \leq \frac{2(1+\tau)}{m}\delta.$$

But from Proposition 1 we have $\|e_{n_\delta}\|_\mu \leq C_*\|\omega\|$. The desired estimate then follows from the interpolation inequality (2.3) and (3.7). \square

Remark 1 If F satisfies (2.16) and $\{x_n\}$ is defined by (1.7) with $s > -a/2$, then the order optimal convergence rate holds for $x_0 - x^\dagger \in X_\mu$ with $0 < \mu \leq a+2s$. On the other hand, if $F'(x)$ satisfies the Lipschitz condition

$$\|F'(x) - F'(x^\dagger)\| \leq K_0\|x - x^\dagger\|, \quad x \in B_\rho(x^\dagger)$$

and $\{x_n\}$ is defined by (1.7) with $s > a/2$, then the order optimal convergence rate holds for $x_0 - x^\dagger \in X_\mu$ with $a < \mu \leq 2s$.

4 Examples

In this section we will give several important examples of $\{g_\alpha\}$ that satisfy Assumptions 1 and 2. Thus, Theorem 1 applies to the corresponding methods if F satisfies Assumption 3 and $\{\alpha_n\}$ satisfies (2.8). For all these examples, the functions g_α are analytic at least in the domain

$$D_\alpha := \{z \in \mathbb{C} : z \neq -\alpha, -1\}.$$

Moreover, for each $\alpha > 0$, we always take the closed contour Γ_α to be (see [1])

$$\Gamma_\alpha = \Gamma_\alpha^{(1)} \cup \Gamma_\alpha^{(2)} \cup \Gamma_\alpha^{(3)} \cup \Gamma_\alpha^{(4)},$$

with

$$\begin{aligned} \Gamma_\alpha^{(1)} &:= \{z = \frac{\alpha}{2}e^{i\phi} : \phi_0 \leq \phi \leq 2\pi - \phi_0\}, \\ \Gamma_\alpha^{(2)} &:= \{z = Re^{i\phi} : -\phi_0 \leq \phi \leq \phi_0\}, \\ \Gamma_\alpha^{(3)} &:= \{z = te^{i\phi_0} : \alpha/2 \leq t \leq R\}, \\ \Gamma_\alpha^{(4)} &:= \{z = te^{-i\phi_0} : \alpha/2 \leq t \leq R\}, \end{aligned}$$

where $R > \max\{1, \alpha\}$ and $0 < \phi_0 < \pi/2$ are fixed numbers. Clearly $\Gamma_\alpha \subset D_\alpha$ and $[0, 1]$ lies inside Γ_α . It is straightforward to check that (2.9) is satisfied.

Example 1 We first consider for $\alpha > 0$ the function g_α given by

$$g_\alpha(\lambda) = \frac{(\alpha + \lambda)^N - \alpha^N}{\lambda(\alpha + \lambda)^N}$$

where $N \geq 1$ is a fixed integer. This function arises from the iterated Tikhonov regularization of order N for linear ill-posed problems. The corresponding method (1.7) becomes

$$\begin{aligned} u_{n,0} &= x_n, \\ u_{n,l+1} &= u_{n,l} - (\alpha_n L^{2s} + T_n^* T_n)^{-1} T_n^* (F(x_n) - y^\delta - T_n(x_n - u_{n,l})), \\ &\quad l = 0, \dots, N-1, \\ x_{n+1} &= u_{n,N}, \end{aligned}$$

where $T_n := F'(x_n)$. When $N = 1$, this is the Levenberg-Marquardt method in Hilbert scales. The corresponding residual function is $r_\alpha(\lambda) = \alpha^N (\alpha + \lambda)^{-N}$. In order to verify Assumption 2, we recall the inequality (see [9, Lemma 3])

$$\lambda \prod_{k=j}^n \frac{\alpha_k}{\alpha_k + \lambda} \leq (s_n - s_{j-1})^{-1} \quad \text{for all } \lambda \geq 0.$$

Then for $0 \leq \nu \leq 1$ and $\lambda \geq 0$ we have

$$\lambda^\nu \prod_{k=j}^n r_{\alpha_k}(\lambda) \leq \left(\lambda \prod_{k=j}^n \frac{\alpha_k}{\alpha_k + \lambda} \right)^\nu \leq (s_n - s_{j-1})^{-\nu}$$

and

$$\begin{aligned}
\lambda^\nu g_{\alpha_j}(\lambda) \prod_{k=j+1}^n r_{\alpha_k}(\lambda) &= \frac{(\alpha_j + \lambda)^N - \alpha_j^N}{\alpha_j^N \lambda^{1-\nu}} \prod_{k=j}^n \left(\frac{\alpha_k}{\alpha_k + \lambda} \right)^N \\
&= \sum_{l=0}^{N-1} \binom{N}{l} \alpha_j^{l-N} \lambda^{N+\nu-l-1} \prod_{k=j}^n \left(\frac{\alpha_k}{\alpha_k + \lambda} \right)^N \\
&\leq \sum_{l=0}^{N-1} \binom{N}{l} \alpha_j^{l-N} \left(\lambda \prod_{k=j}^n \frac{\alpha_k}{\alpha_k + \lambda} \right)^{N+\nu-l-1} \\
&\leq \sum_{l=0}^{N-1} \binom{N}{l} \alpha_j^{l-N} (s_n - s_{j-1})^{-N-\nu+l+1} \\
&\leq C_N \frac{1}{\alpha_j} (s_n - s_{j-1})^{-\nu},
\end{aligned}$$

where $C_N = 2^N - 1$ and we used the fact $\alpha_j^{-1} \leq s_n - s_{j-1}$. We therefore obtain (2.12) and (2.13) in Assumption 2.

Next we will verify (2.10) in Assumption 1. Note that

$$\varphi_\alpha(z) = \frac{\alpha(\alpha + z)^{N-1} - \alpha^N}{z(\alpha + z)^N} = \frac{1}{z(\alpha + z)^N} \sum_{j=0}^{N-2} \binom{N-1}{j} \alpha^{j+1} z^{N-1-j}.$$

It is easy to check $|\varphi_\alpha(z)| \lesssim \alpha^{-1}$ on $\Gamma_\alpha^{(1)}$ and $|\varphi_\alpha(z)| \lesssim 1$ on $\Gamma_\alpha^{(2)}$. Moreover, on $\Gamma_\alpha^{(3)} \cup \Gamma_\alpha^{(4)}$ there holds

$$|\varphi_\alpha(z)| \lesssim \frac{1}{t(\alpha + t)^N} \sum_{j=0}^{N-2} \alpha^{j+1} t^{N-1-j} \lesssim \sum_{j=0}^{N-2} \alpha^{j+1} t^{-2-j}.$$

Therefore

$$\begin{aligned}
\int_{\Gamma_\alpha} |\varphi_\alpha(z)| |dz| &= \int_{\Gamma_\alpha^{(1)}} |\varphi_\alpha(z)| |dz| + \int_{\Gamma_\alpha^{(2)}} |\varphi_\alpha(z)| |dz| + \int_{\Gamma_\alpha^{(3)} \cup \Gamma_\alpha^{(4)}} |\varphi_\alpha(z)| |dz| \\
&\lesssim \alpha^{-1} \int_{\phi_0}^{2\pi - \phi_0} \alpha d\phi + \int_{-\phi_0}^{\phi_0} d\phi + \sum_{j=0}^{N-2} \alpha^{j+1} \int_{\alpha/2}^R t^{-2-j} dt \\
&\lesssim 1.
\end{aligned}$$

Assumption 1 is therefore verified.

Example 2 We consider the method (1.7) with g_α given by

$$g_\alpha(\lambda) = \frac{1}{\lambda} \left(1 - e^{-\lambda/\alpha} \right)$$

which arises from the asymptotic regularization for linear ill-posed problems. In this method, the iterative sequence $\{x_n\}$ is equivalently defined as $x_{n+1} := x(1/\alpha_n)$, where $x(t)$ is the unique solution of the initial value problem

$$\begin{aligned}
\frac{d}{dt} x(t) &= L^{-2s} F'(x_n)^* (y^\delta - F(x_n) + F'(x_n)(x_n - x(t))), \quad t > 0, \\
x(0) &= x_n.
\end{aligned}$$

The corresponding residual function is $r_\alpha(\lambda) = e^{-\lambda/\alpha}$. We first verify Assumption 2. It is easy to see

$$\lambda^\nu \prod_{k=j}^n r_{\alpha_k}(\lambda) = \lambda^\nu e^{-\lambda(s_n - s_{j-1})} \leq \nu^\nu e^{-\nu(s_n - s_{j-1})^{-\nu}} \leq (s_n - s_{j-1})^{-\nu}$$

for $0 \leq \nu \leq 1$ and $\lambda \geq 0$. This shows (2.12). By using the elementary inequality $e^{-p\lambda} - e^{-q\lambda} \leq (q-p)/q$ for $0 < p \leq q$ and $\lambda \geq 0$ and observing that $0 \leq r_\alpha(\lambda) \leq 1$ and $0 \leq g_\alpha(\lambda) \leq 1/\alpha$, we have for $0 \leq \nu \leq 1$ and $\lambda \geq 0$ that

$$\begin{aligned} \lambda^\nu g_{\alpha_j}(\lambda) \prod_{k=j+1}^n r_{\alpha_k}(\lambda) &\leq \frac{1}{\alpha_j^{1-\nu}} \left(\lambda g_{\alpha_j}(\lambda) \prod_{k=j+1}^n r_{\alpha_k}(\lambda) \right)^\nu \\ &= \frac{1}{\alpha_j^{1-\nu}} \left(e^{-(s_n - s_j)\lambda} - e^{-(s_n - s_{j-1})\lambda} \right)^\nu \\ &\leq \frac{1}{\alpha_j} (s_n - s_{j-1})^{-\nu} \end{aligned}$$

which gives (2.13).

In order to verify (2.10) in Assumption 1, we note that

$$\varphi_\alpha(z) = \frac{1 - e^{-z/\alpha}}{z} - \frac{1}{\alpha + z} = \frac{\alpha - (\alpha + z)e^{-z/\alpha}}{z(\alpha + z)}.$$

It is easy to see that $|\varphi_\alpha(z)| \lesssim \alpha^{-1}$ on $\Gamma_\alpha^{(1)}$, $|\varphi_\alpha(z)| \lesssim 1$ on $\Gamma_\alpha^{(2)}$ and

$$|\varphi_\alpha(z)| \lesssim \frac{\alpha + (\alpha + t)e^{-\frac{t}{\alpha} \cos \phi_0}}{t(\alpha + t)} \lesssim \alpha t^{-2}$$

on $\Gamma_\alpha^{(3)} \cup \Gamma_\alpha^{(4)}$. Therefore

$$\int_{\Gamma_\alpha} |\varphi_\alpha(z)| |dz| \lesssim 1 + \int_{\alpha/2}^R \alpha t^{-2} dt \lesssim 1.$$

Example 3 We consider for $0 < \alpha \leq 1$ the function g_α given by

$$g_\alpha(\lambda) = \sum_{l=0}^{[1/\alpha]-1} (1-\lambda)^l = \frac{1 - (1-\lambda)^{[1/\alpha]}}{\lambda}$$

which arises from the linear Landweber iteration, where $[1/\alpha]$ denotes the largest integer not greater than $1/\alpha$. The method (1.7) then becomes

$$\begin{aligned} u_{n,0} &= x_n, \\ u_{n,l+1} &= u_{n,l} - L^{-2s} T_n^* (F(x_n) - y^\delta - T_n(x_n - u_{n,l})), \quad 0 \leq l \leq [1/\alpha_n] - 1, \\ x_{n+1} &= u_{n,[1/\alpha_n]}, \end{aligned}$$

where $T_n := F'(x_n)$. When $\alpha_n = 1$ for all n , this method reduces to the Landweber iteration in Hilbert scales proposed in [13]. The corresponding residual function is $r_\alpha(\lambda) = (1-\lambda)^{[1/\alpha]}$. We first verify Assumption 2 when the sequence $\{\alpha_n\}$ is given by $\alpha_n = 1/k_n$ for some integers $k_n \geq 1$. Then for $0 \leq \nu \leq 1$ and $0 \leq \lambda \leq 1$ we have

$$\lambda^\nu \prod_{k=j}^n r_{\alpha_k}(\lambda) = \lambda^\nu (1-\lambda)^{s_n - s_{j-1}} \leq \nu^\nu (s_n - s_{j-1})^{-\nu} \leq (s_n - s_{j-1})^{-\nu}.$$

We thus obtain (2.12). Observing that $0 \leq r_{\alpha_j}(\lambda) \leq 1$ and $0 \leq g_{\alpha_j}(\lambda) \leq 1/\alpha_j$ for $0 \leq \lambda \leq 1$, we have

$$\begin{aligned} \lambda^\nu g_{\alpha_j}(\lambda) \prod_{k=j+1}^n r_{\alpha_k}(\lambda) &\leq \frac{1}{\alpha_j^{1-\nu}} \left(\lambda g_{\alpha_j}(\lambda) \prod_{k=j+1}^n r_{\alpha_k}(\lambda) \right)^\nu \\ &= \frac{1}{\alpha_j^{1-\nu}} \left((1-\lambda)^{s_n-s_j} - (1-\lambda)^{s_n-s_{j-1}} \right)^\nu. \end{aligned}$$

Thus, (2.13) follows from the elementary inequality $t^p - t^q \leq (q-p)/q$ for $0 < p \leq q$ and $0 \leq t \leq 1$.

In order to verify (2.10) in Assumption 1, in the definition of Γ_α we pick $R > 1$ and $0 < \phi_0 < \pi/2$ such that $R < 2 \cos \phi_0$. Note that

$$\varphi_\alpha(z) = \frac{1 - (1-z)^{[1/\alpha]}}{z} - \frac{1}{\alpha+z} = \frac{\alpha - (\alpha+z)(1-z)^{[1/\alpha]}}{z(\alpha+z)}.$$

By using the fact $(1+\alpha)^{1/\alpha} \leq e$ we can see

$$|\varphi_\alpha(z)| \lesssim \alpha^{-1} (1+\alpha/2)^{1/\alpha} \lesssim \alpha^{-1} \quad \text{on } \Gamma_\alpha^{(1)}.$$

According to the choice of R and ϕ_0 , we have $1 + R^2 - 2R \cos \phi_0 < 1$. Thus

$$|\varphi_\alpha(z)| \lesssim \frac{\alpha + (\alpha+R)(1+R^2-2R \cos \phi_0)^{[1/\alpha]/2}}{R(R+\alpha)} \lesssim 1 \quad \text{on } \Gamma_\alpha^{(2)}.$$

Furthermore, on $\Gamma_\alpha^{(3)} \cup \Gamma_\alpha^{(4)}$ we have

$$|\varphi_\alpha(z)| \lesssim \frac{\alpha + (\alpha+t)(1+t^2-2t \cos \phi_0)^{1/(2\alpha)}}{t(\alpha+t)}.$$

Therefore

$$\begin{aligned} \int_{\Gamma_\alpha} |\varphi_\alpha(z)| |dz| &\lesssim 1 + \int_{\alpha/2}^R \frac{\alpha + (\alpha+t)(1+t^2-2t \cos \phi_0)^{1/(2\alpha)}}{t(\alpha+t)} dt \\ &= 1 + \int_{1/2}^{R/\alpha} \frac{1 + (1+t)(1+\alpha^2 t^2 - 2\alpha t \cos \phi_0)^{1/(2\alpha)}}{t(1+t)} dt \\ &\lesssim 1 + \int_{1/2}^{R/\alpha} (1+\alpha^2 t^2 - 2\alpha t \cos \phi_0)^{1/(2\alpha)} dt. \end{aligned}$$

Observe that for $1/2 \leq t \leq R/\alpha$ there holds

$$(1+\alpha^2 t^2 - 2\alpha t \cos \phi_0)^{1/(2\alpha)} \leq (1-\mu_0 \alpha t)^{1/(2\alpha)} \leq e^{-\mu_0 t/2}$$

with $\mu_0 := 2 \cos \phi_0 - R > 0$. Thus

$$\int_{\Gamma_\alpha} |\varphi_\alpha(z)| |dz| \lesssim 1 + \int_{1/2}^\infty e^{-\mu_0 t/2} dt \lesssim 1.$$

Example 4 We consider for $0 < \alpha \leq 1$ the function g_α given by

$$g_\alpha(\lambda) = \sum_{i=1}^{[1/\alpha]} (1+\lambda)^{-i} = \frac{1 - (1+\lambda)^{-[1/\alpha]}}{\lambda}$$

which arises from the Lardy method for linear inverse problems. Then the method (1.7) becomes

$$\begin{aligned} u_{n,0} &= x_n, \\ u_{n,l+1} &= u_{n,l} - (L^{2s} + T_n^* T_n)^{-1} T_n^* (F(x_n) - y^\delta - T_n(x_n - u_{n,l})), \\ &\quad l = 0, \dots, [1/\alpha_n] - 1, \\ x_{n+1} &= u_{n,[1/\alpha_n]}, \end{aligned}$$

where $T_n = F'(x_n)$. The residual function is $r_\alpha(\lambda) = (1 + \lambda)^{-[1/\alpha]}$. Assumption 1 and Assumption 2 can be verified similarly as in Example 3 when the sequence $\{\alpha_n\}$ is given by $\alpha_n = 1/k_n$ for some integers $k_n \geq 1$.

References

1. A. B. Bakushinsky and M. Yu. Kokurin, *Iterative Methods for Approximate Solutions of Inverse Problems*, Mathematics and its applications, Springer, 2004.
2. R. S. Dembo, S. C. Eisenstat and T. Steihaug, *Inexact Newton methods*, SIAM J. Numer. Anal., 19 (1982), 400–408.
3. H. W. Engl, M. Hanke and A. Neunauer, *Regularization of Inverse Problems*, Kluwer, Dordrecht, 1996.
4. M. Hanke, *A regularizing Levenberg-Marquardt scheme with applications to inverse groundwater filtration problems*, Inverse Problems, 13(1997), 79–95.
5. M. Hanke, *The regularizing Levenberg-Marquardt scheme is of optimal order*, J. Integral Equations and Applications, 22 (2010), no. 2, 259–283.
6. M. Hanke, A. Neubauer and O. Scherzer, *A convergence analysis of the Landweber iteration for nonlinear ill-posed problems*, Numer. Math., 72 (1995), 21–37.
7. M. Hochbruck, M. Hönl and A. Ostermann, *A convergence analysis of the exponential Euler iteration for nonlinear ill-posed problems*, Inverse Problems, 25 (2009), no.7, article no. 075009.
8. Q. Jin, *Error estimates of some Newton-type methods for solving nonlinear inverse problems in Hilbert scales*, Inverse Problems, 16 (2000), no. 1, 187–197.
9. Q. Jin, *On a regularized Levenberg-Marquardt method for solving nonlinear inverse problems*, Numer. Math., 115 (2010), no. 2, 229–259.
10. J. Köhler and U. Tautenhahn, *Error bounds for regularized solutions of nonlinear ill-posed problems*, J. Inv. Ill-Posed Problems 3 (1995), 47–74.
11. A. Lechleiter and A. Rieder, *Towards a general convergence theory for inexact Newton regularizations*, Numer. Math. 114 (2010), no. 3, 521–548.
12. F. Natterer, *Error bounds for Tikhonov regularization in Hilbert scales*, Appl. Anal., 18 (1984), 29–37.
13. A. Neubauer, *On Landweber iteration for nonlinear ill-posed problems in Hilbert scales*, Numer. Math., 85 (2000), 309–328.
14. A. Rieder, *On the regularization of nonlinear ill-posed problems via inexact Newton iterations*, Inverse Problems, 15(1999), 309–327.
15. A. Rieder, *On convergence rates of inexact Newton regularizations*, Numer. Math. 88(2001), 347–365.
16. U. Tautenhahn, *Error estimates for regularization methods in Hilbert scales*, SIAM J. Numer. Anal., 33 (1996), 2120–2130.
17. U. Tautenhahn, *On a general regularization scheme for nonlinear ill-posed problems: II. regularization in Hilbert scales*, Inverse Problems, 14 (1998), 1607–1616.