

# Effect of Sun and Planet-Bound Dark Matter on Planet and Satellite Dynamics in the Solar System

L. Iorio\*

\**INFN-Sezione di Pisa. Address for correspondence: Viale Unità di Italia 68 70125 Bari (BA), Italy.*

*e-mail: lorenzo.iorio@libero.it*

## Abstract

We apply our recent results on orbital dynamics around a mass-varying central body to the phenomenon of accretion of Dark Matter—assumed not self-annihilating—on the Sun and the major bodies of the solar system due to its motion throughout the Milky Way halo. We inspect its consequences on the orbits of the planets and their satellites over timescales of the order of the age of the solar system. It turns out that a solar Dark Matter accretion rate of  $\approx 10^{-12} \text{ yr}^{-1}$ , inferred from the upper limit  $\Delta M/M = 0.02 - 0.05$  on the Sun's Dark Matter content, assumed somehow accumulated during last 4.5 Gyr, would have displaced the planets faraway by about  $10^{-2} - 10^1 \text{ au}$  4.5 Gyr ago. Another consequence is that the semimajor axis of the Earth's orbit, approximately equal to the Astronomical Unit, would undergo a secular increase of  $0.02 - 0.05 \text{ m yr}^{-1}$ , in agreement with the latest observational determinations of the Astronomical Unit secular increase of  $0.07 \pm 0.02 \text{ m yr}^{-1}$  and  $0.05 \text{ m yr}^{-1}$ . By assuming that the Sun will continue to accrete Dark Matter in the next billions year at the same rate as in the past, the orbits of its planets will shrink by about  $10^{-1} - 10^1 \text{ au}$  ( $\approx 0.2 - 0.5 \text{ au}$  for the Earth), with consequences for their fate, especially of the inner planets. Dark Matter accretion on planets has, instead, less relevant consequences for their satellites. Indeed, 4.5 Gyr ago their orbits would have been just  $10^{-2} - 10^1 \text{ km}$  wider than now. Dark Matter accretion is not able to explain the observed accelerations of the orbits of some of the Galilean satellites of Jupiter, the secular decrease of the semimajor axis of the Earth's artificial satellite LAGEOS and the secular increase of the Moon's orbit eccentricity.

Keywords: Experimental studies of gravity; Dark matter; Solar system objects; Celestial mechanics

PACS: 04.80.-y, 95.35.+d, 96.30.-t, 95.10.Ce

# 1 Introduction

Dark Matter (DM) was postulated long ago to explain the discrepancy between the observed kinematics of some components of astrophysical systems like clusters of galaxies [1] and spiral galaxies [2, 3], and the predicted one on the basis of the Newtonian dynamics and the matter directly detected from the emitted electromagnetic radiation (visible stars and gas clouds). Zwicky [4] postulated the existence of undetected, baryonic matter; today [5] it is believed that the hidden mass is constituted by non-baryonic, weakly interacting particles in order to cope with certain issues pertaining galaxy and galaxy clusters formation [6] and primordial nucleosynthesis [7]. On cosmological scales, DM accounts for about 23% of the mass-energy content of the Universe [8].

It is important to perform tests of the DM hypothesis in other scenarios, independently of the phenomena themselves for which its existence was originally postulated. Our solar system can be regarded as one of such different laboratories. The first arguments in support of invisible (standard) matter in solar system were presented in 1932 by Oort [9] in connection with the dynamical density in the vicinity of the Sun's Galactic orbit. Oort concluded that the total mass of nebulous or meteoric matter near the Sun is less than  $3 \times 10^{-24} \text{ g cm}^{-3}$ . Nowadays, it is evaluated that the upper bound on local DM density in the solar system

$$\rho_{\text{ss}} \cong 2 \times 10^{-19} \text{ g cm}^{-3} = 10^5 \left( \frac{\text{Gev}}{c^2} \right) \text{ cm}^{-3} \quad (1)$$

is several orders of magnitude higher than the mean Galactic value [10]

$$\rho_{\text{gal}} \cong 4 \times 10^{-25} \text{ g cm}^{-3} = 0.3 \left( \frac{\text{Gev}}{c^2} \right) \text{ cm}^{-3}, \quad (2)$$

so that it could be imagined that the solar system is surrounded by a local subhalo, a suggestion reinforced by recent simulations [11]. Several processes have been postulated to clump DM in the solar system [12, 13, 22, 15, 16, 17, 18, 19]; according to Ref. [20, 21], the existence of the solar system itself might be evidence for a local subhalo. However, it must be noted that, according to Ref. [22], densities of DM as large as that of eq. (1) could not be reached because of the inverse process of capturing DM from the Milky Way halo. According to it, whenever the density of DM bound to the solar system gets big, the process of throwing WIMPs out of it will be in equilibrium with capture from the Galactic halo. This fact has been proven

by using the Liouville’s theorem, which says that the phase space density (after a long time, when equilibrium has been reached) of the solar system bound WIMPs is the same as in the halo [22, 15]. Several studies have been dedicated to placing bounds on the local distribution of DM in our solar system from orbital motions of natural major and minor bodies and artificial probes<sup>1</sup> [23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 20, 35, 36], and electromagnetic wave propagation [37]. It has also been suggested that some proposed and approved space-based missions for fundamental physics may be used to search for local distributions of DM as well [38, 39, 40]. Apart from directly affecting orbital motions and electromagnetic waves propagation, DM may also have other effects on the solar system’s bodies: for example, it may concur to form main-sequence stars like the Sun [41] and accrete on the major planets [42, 21], and affect their internal heats [43, 44]. According to Ref. [42], during its journey along the Galaxy since its birth 4.5 Gyr ago the solar system would have encountered about  $203M_{\odot}$  of DM. In Ref. [41] it is stated that the upper limit to the amount of gravitating DM in the Sun would be 2 – 5% of the total solar mass; by assuming that it somehow accumulated during the Sun’s lifetime<sup>2</sup>, it is possible to infer a mass increase rate of

$$\frac{\dot{M}}{M} = 4.4 - 11 \times 10^{-12} \text{ yr}^{-1}. \quad (3)$$

Accretion of DM, in the form of Mirror Matter [45], by the Sun during its lifetime is envisaged in Ref. [38]. Indeed, according to eq. (7) of Ref. [38], reproduced below, a gravitating body of mass  $M$  traveling at speed  $v$  through a volumetric distribution of DM with density  $\varrho$  will accrete an amount  $\Delta M$  over a time span  $t$

$$\frac{\Delta M}{M} \simeq 10^{-5} \left( \frac{M}{M_{\odot}} \right) \left( \frac{v}{10 \text{ km s}^{-1}} \right)^{-3} \left( \frac{\varrho}{10^{-24} \text{ g cm}^{-3}} \right) \left( \frac{t}{10^{10} \text{ yr}} \right). \quad (4)$$

Thus, if we assume  $v = 30 \text{ km s}^{-1}$  and the upper limit of the local DM density in the solar system given by eq. (1), the Sun will increase its mass at a rate

$$\frac{\dot{M}}{M} \simeq 7 \times 10^{-12} \text{ yr}^{-1}, \quad (5)$$

---

<sup>1</sup>The bound of eq. (1) comes just from such investigations.

<sup>2</sup>In fact, in Ref. [41] it is argued that the DM content of the Sun cannot be due to a continuous accretion throughout its lifetime; however, the value of the amount of Galactic DM encountered by the Sun in the latest 4.5 Gyr in Ref. [41] radically differs from that in Ref. [42].

close to eq. (3) from Ref. [41]. For several possible observable manifestations of Mirror Matter at various astronomical scales, see Ref. [46].

Concerning the planets, according to Table II and Table III of Ref. [42], Jupiter would have captured the largest amount of DM during the latest 4.5 Gyr followed by Saturn, Uranus and Neptune. See Table 1 for the related relative mass variation rates  $\dot{M}/M$  of the eight planets.

Table 1: DM accretion rates  $\dot{M}/M$  on major bodies of the solar system according to Table II and Table III of Ref. [42].

Planet	$\dot{M}/M$ (yr <sup>-1</sup> )
Mercury	$2.7 \times 10^{-17}$
Venus	$1.6 \times 10^{-17}$
Earth	$1.4 \times 10^{-17}$
Mars	$4.1 \times 10^{-17}$
Jupiter	$0.6 \times 10^{-17}$
Saturn	$1.1 \times 10^{-17}$
Uranus	$2.9 \times 10^{-17}$
Neptune	$3.5 \times 10^{-17}$

The estimates in Ref. [21] are different but, as we will show, such discrepancies will not substantially affect the main result of our analysis.

We will investigate if the phenomenon of DM capture and accretion by the Sun and the planets has somewhat influenced the solar system dynamics throughout its 4.5 Gyr lifetime. We will address this problem in a specific way, i.e. we will look at the possible modifications of the orbital trajectories of the satellites of the major bodies which experienced the DM accretion, assumed not self-annihilating [44].

The paper is organized as follows. In Section 2 we will briefly review some basics of orbital motion around a mass-varying body. In Section 3 we will apply our results to concrete astronomical scenarios in the solar system. Section 4 is devoted to the Conclusions.

## 2 Orbital dynamics around a mass-varying body

The acceleration experienced by a test particle orbiting a mass-varying central body with  $\mu(t) \doteq GM(t)$  can be approximated with

$$\mathbf{A} = -\frac{\mu(t)}{r^2}\hat{\mathbf{r}} \approx -\frac{\mu}{r^2} \left[ 1 + \left( \frac{\dot{\mu}}{\mu} \right) (t - t_0) \right] \hat{\mathbf{r}}, \quad (6)$$

where  $t_0$  is a given epoch and  $\mu \doteq \mu|_{t=t_0}$ ,  $\dot{\mu} \doteq \dot{\mu}|_{t=t_0}$ . In our case  $t_0$  coincides with the present time, and  $t < t_0$ ; thus,  $t - t_0 < 0$ . Since in our case  $\dot{\mu}$  is due to DM accretion, we will assume that  $\dot{\mu} = \text{const.}$  throughout the solar system's lifetime, i.e. we will not deal with  $\ddot{\mu}$ ; the validity of such an approximation will be justified a-posteriori in Section 3. Moreover, the condition of validity of eq. (6), i.e.  $(\dot{\mu}/\mu)(t - t_0) \ll 1$ , is satisfied for all the major bodies of the solar system over  $\Delta t = -4.5$  Gyr, given the DM accretion rates quoted in Section 1.

In Ref. [47] the secular, i.e. averaged over one orbital revolution, orbital effects induced by eq. (6) on the usual Keplerian orbital elements<sup>3</sup> of a test particle have been analytically worked out by means of the Gauss equations for the variations of elements. They are

$$\langle \dot{a} \rangle = \frac{2ea}{(1-e)} \left( \frac{\dot{\mu}}{\mu} \right) \quad (7)$$

$$\langle \dot{e} \rangle = (1+e) \left( \frac{\dot{\mu}}{\mu} \right) \quad (8)$$

$$\langle \dot{I} \rangle = 0 \quad (9)$$

$$\langle \dot{\Omega} \rangle = 0 \quad (10)$$

$$\langle \dot{\omega} \rangle = 0 \quad (11)$$

$$\langle \dot{\mathcal{M}} \rangle = n + 2\pi \left( \frac{\dot{\mu}}{\mu} \right), \quad (12)$$

where  $a, e, I, \Omega, \omega, \mathcal{M}$  are the values at the epoch of the semimajor axis, the eccentricity, the inclination, the longitude of the ascending node, the argument of pericenter and the mean anomaly, respectively;  $n \doteq \sqrt{\mu/a^3}$  is the Keplerian mean motion. Such results are not in contrast with the orbital evolution effectively followed by the test particle, as shown in detail in Ref. [47]. Indeed, for, e.g.,  $\dot{\mu}/\mu < 0$ , i.e. for a mass decrease, the trajectory, which is not an ellipse, would expand in such a way that the osculating semimajor axis and eccentricity of the Keplerian ellipses approximating it at the pericentre, which remains fixed, get reduced revolution after revolution, in accordance to eq. (7) and eq. (8); see Figure 1 of Ref. [47] and the related discussion. Moreover, also the osculating Keplerian period  $P_b \doteq 2\pi/n$ , i.e. the time required to fully describe an osculating Keplerian ellipse, gets shorter, according to

$$\left\langle \frac{dP_b}{dt} \right\rangle = \frac{3}{2} P_b \frac{\langle \dot{a} \rangle}{a}, \quad (13)$$

---

<sup>3</sup>The type I, according to Ref. [48]

while the time of two consecutive pericentre crossings along the true path grows.

The variations of the radial  $\rho$  and transverse  $\tau$  components of the radius vector, taken at two consecutive pericentre passages, are [47]

$$\Delta\rho = -\frac{2\pi}{n}a(1-e)\left(\frac{\dot{\mu}}{\mu}\right), \quad (14)$$

$$\Delta\tau = \frac{4\pi^2}{n}a\left(\frac{\dot{\mu}}{\mu}\right)\sqrt{\frac{1+e}{1-e}}, \quad (15)$$

so that the overall pericenter-to-pericenter orbit variation is

$$\Delta r = \Delta\rho\sqrt{1 + 4\pi^2\frac{(1+e)}{(1-e)^3}}. \quad (16)$$

### 3 Consequences of DM accretion on the orbits of the satellites of the major planets

Here we will fruitfully apply the results of Section 2 to the problem of determining the impact of DM accretion on the Sun and the major planets on the orbital evolution of their planetary and satellite systems throughout the past 4.5 Gyr.

Concerning the Sun, by assuming for it  $\dot{M}/M = 4.4-11\times 10^{-12} \text{ yr}^{-1}$  [41], it turns out that 4.5 Gyr ago the orbits of the planets were larger than now by the amounts listed in Table 2. It can be noted that, while the distances of

Table 2: Increment  $\Delta r$  of the planetary orbits for  $\dot{M}/M = 4.4 - 11 \times 10^{-12} \text{ yr}^{-1}$  [41] and  $\Delta t = -4.5 \text{ Gyr}$  according to eq. (16).

Planet	$\Delta r$ (au)
Mercury	0.06 – 0.15
Venus	0.09 – 0.23
Earth	0.13 – 0.32
Mars	0.21 – 0.52
Jupiter	0.68 – 1.71
Saturn	1.26 – 3.17
Uranus	2.53 – 6.31
Neptune	3.82 – 9.55

the inner planets at their formation are not substantially altered, the giant gaseous planets are shifted by about 1 – 10 au with respect to their present-day locations. If the rate of DM accretion on the Sun will be the same for the remaining lifetime of the solar system, the consequences for its planets would be relevant because of the continuous shrinking of their orbits. Indeed, when the Sun will have reached the tip of the Red Giant Branch (RGB) in the Hertzsprung-Russel (HR) diagram in the next 7.58 Gyr [49], the orbits of the planets will be reduced by the amounts listed in Table 3. In particular, the

Table 3: Decrement  $\Delta r$  of the planetary orbits for  $\dot{M}/M = 4.4 - 11 \times 10^{-12} \text{ yr}^{-1}$  [41] and  $\Delta t = +7.58 \text{ Gyr}$  according to eq. (16).

Planet	$\Delta r$ (au)
Mercury	-(0.10 – 0.25)
Venus	-(0.15 – 0.38)
Earth	-(0.21 – 0.54)
Mars	-(0.35 – 0.88)
Jupiter	-(1.15 – 2.89)
Saturn	-(2.13 – 5.33)
Uranus	-(4.25 – 10.65)
Neptune	-(6.43 – 16.08)

heliocentric distance of the Earth may be reduced by  $\approx 0.2 - 0.5$  au. Figure 2 of Ref. [49] shows that at the beginning of RGB the solar photosphere will reach about 0.5–0.6 au; after entering the RG phase things will dramatically change because in only  $\approx 1$  Myr the Sun will reach the tip of the RGB phase expanding up to 1.20 au, while the Earth’s distance will be just 0.8 – 0.5 au or less. However, it must be pointed out that the results of Ref. [49] do not take into account possible DM accretion which, in principle, may have consequences on the lifetime cycle of main sequence stars depending on the type of DM involved [50]. Another interesting consequence of the Sun’s DM accretion is that the semimajor axis  $a_{\oplus}$  of the Earth’s orbit, which can be approximately taken equal to the Astronomical Unit amounting to<sup>4</sup>  $a_{\oplus} = 1.00000018$  au, undergoes a secular increment of just  $0.02 - 0.05 \text{ m yr}^{-1}$  according to eq. (7) and  $\dot{M}/M = 4.4 - 11 \times 10^{-12} \text{ yr}^{-1}$ . An anomalous increase of the Astronomical Unit of  $0.15 \pm 0.04 \text{ m yr}^{-1}$  was reported for the first time in Ref. [51]. Later results are  $0.07 \pm 0.02 \text{ m yr}^{-1}$  [52] and  $0.05 \text{ m yr}^{-1}$  [53]. Anyway, we point out that the figures presented here would

<sup>4</sup>See [http://ssd.jpl.nasa.gov/txt/p\\_elem\\_t2.txt](http://ssd.jpl.nasa.gov/txt/p_elem_t2.txt) on the WEB.

be radically smaller if eq. (2) for the Galactic mean density was used in eq. (4) of Ref. [38].

Let us, now, move to the planetary systems of satellites. In the case of the Earth and Moon, eq. (16) and  $\dot{M}/M = 1.4 \times 10^{-17} \text{ yr}^{-1}$  from Ref. [42] yields an expansion of the lunar orbit of just 0.16 km. The recently observed secular increase of the eccentricity of the geocentric Moon's orbit amounts to  $\langle \dot{e} \rangle = (0.9 \pm 0.3) \times 10^{-11} \text{ yr}^{-1}$  [54]. It is several orders of magnitude larger than the secular increase predicted by eq. (8). Concerning the observationally determined secular decrement  $\dot{a} = -0.4 \text{ m yr}^{-1}$  of the semimajor axis of the orbit of the geodetic satellite LAGEOS [55], DM accretion is unable to explain it because eq. (7) yields an increase of the order of  $10^{-12} \text{ m yr}^{-1}$ .

In Table 4-Table 7 we quote the expansion of the orbits of some satellites of Jupiter, Saturn, Uranus and Neptune 4.5 Gyr ago induced by the DM accretion according to Ref. [42]. It can be shown that the effect of such

Table 4: Jupiter: increment  $\Delta r$  of the Galilean satellites' orbits and of S/2003J2 for  $\dot{M}/M = 0.6 \times 10^{-17} \text{ yr}^{-1}$  [42] and  $\Delta t = -4.5 \text{ Gyr}$  according to eq. (16).

Satellite	$\Delta r$ (km)
Io	0.07
Europa	0.11
Ganymede	0.17
Callisto	0.31
S/2003J2	7.21

Table 5: Saturn: increment  $\Delta r$  of some satellites' orbits for  $\dot{M}/M = 1.1 \times 10^{-17} \text{ yr}^{-1}$  [42] and  $\Delta t = -4.5 \text{ Gyr}$  according to eq. (16).

Satellite	$\Delta r$ (km)
Mimas	0.06
Titan	0.39
Iapetus	1.15
Phoebe	4.78
Fornjot	9.73

form of  $\dot{M}/M$  is very tiny, amounting approximately to  $10^{-2} - 10^2 \text{ km}$ . This shows that the discrepancies between the evaluations of Ref. [42] and



Table 6: Uranus: increment  $\Delta r$  of some satellites' orbits for  $\dot{M}/M = 2.9 \times 10^{-17} \text{ yr}^{-1}$  [42] and  $\Delta t = -4.5 \text{ Gyr}$  according to eq. (16).

Satellite	$\Delta r$ (km)
Ariel	0.15
Oberon	0.48
Margaret	26.4
Setebos	28.18
Ferdinand	25.62

Table 7: Neptune: increment  $\Delta r$  of some satellites' orbits for  $\dot{M}/M = 3.5 \times 10^{-17} \text{ yr}^{-1}$  [42] and  $\Delta t = -4.5 \text{ Gyr}$  according to eq. (16).

Satellite	$\Delta r$ (km)
Naiad	0.05
Galatea	0.06
Triton	0.35
Nereid	14.46

Ref. [21] are negligible in our case. Moreover, a-posteriori we can well justify the assumption of neglecting  $\ddot{\mu}$ . Let us note that DM accretion on Jupiter cannot be the cause of the observationally determined acceleration of the orbits of Io, Europa and Ganymede [56], because the orbital period of Io is shortening, while those of Europa and Ganymede are lengthening; eq. (13), instead, predicts a uniform increase for  $P_b$ . We can conclude that DM accretion on the major planets of the solar system had negligible impact in sculpting the orbital patterns of their satellite systems.

## 4 Conclusions

In this paper we have investigated the impact that the phenomenon of DM capture by the Sun and the major bodies of the solar system may have on the orbital dynamics of the planets and their satellites on timescales comparable with the age of the solar system.

Concerning the planets, a solar mass increase by DM-assumed not-self-annihilating-as large as  $\approx 10^{-12} \text{ yr}^{-1}$  would displace their orbits outward by about  $10^{-2} - 10^1 \text{ au}$  at the epoch of the solar system's formation 4.5

Gyr ago; for the Earth the increase of its heliocentric distance would be in the range 0.1 – 0.3 au, while the largest shifts are for the giant gaseous planets. The solar DM accretion rate is assumed by postulating that the maximum amount of DM in the Sun, evaluated as 2 – 5% of the total solar mass, comes from a continuous accretion process. Anyway, we note that the real occurrence of such a process is controversial; moreover, also in this case, it would be orders of magnitude smaller if the local DM density was as large as the mean Galactic one. Anyway, one of the consequences of the assumed DM-induced solar accretion is that the osculating semimajor axis of the Earth’s shrinking orbit would experience a secular increase as large as the latest observational determinations of the Astronomical Unit rate of  $\approx 0.07 - 0.05 \text{ m yr}^{-1}$ . In fact, no real contradiction exists because it has been shown that the osculating semimajor axis of a gradually shrinking trajectory around a mass-increasing central body gets, in fact, larger. By assuming that the solar DM accretion will take place in the next billions year at the same rate as in the past, the orbits of the planets will shrink by  $\approx 10^{-1} - 10^1 \text{ au}$  (0.2 – 0.5 au for the Earth), with consequences for their fate, especially for the inner planets.

The DM accretion on the planets has, instead, no effects on the dynamics of their satellites whose planetocentric distances are altered by just  $10^{-2} - 10^1 \text{ km}$  over 4.5 Gyr. Concerning the Earth, both the observationally determined secular decrease of the semimajor axis of the LAGEOS satellite of  $0.4 \text{ m yr}^{-1}$  and the secular increase of the eccentricity of  $(0.9 \pm 0.3) \times 10^{-11} \text{ yr}^{-1}$  of the lunar orbit cannot be explained by the DM accretion on the Earth. The same occurs for the observed accelerations of the orbits of Io, Europa and Ganymede.

## Acknowledgments

I thank M.Yu. Khlopov for useful references and comments.

## References

- [1] F. Zwicky, “Die Rotverschiebung von extragalaktischen Nebeln”. *Helvetica Phisica Acta* **6**, 110-127 (1933).
- [2] A. Bosma, “21-cm line studies of spiral galaxies. I - Observations of the galaxies NGC 5033, 3198, 5055, 2841, and 7331. II - The distri-

- bution and kinematics of neutral hydrogen in spiral galaxies of various morphological types”. *Astron. J.* **86**, 1791-1846 (1981).
- [3] V.C. Rubin, W.K. Ford Jr., N. Thonnard and D. Burstein, “Rotational properties of 23 SB galaxies”. *Astrophys. J.* **261**, 439-456 (1982).
- [4] F. Zwicky, “On the Masses of Nebulae and of Clusters of Nebulae”. *Astrophys. J.* **86**, 217-246 (1937).
- [5] V.C. Rubin, “The rotation of spiral galaxies”. *Science* **220**(4604), 1339-1344 (1983).
- [6] W.A. Chiu, X. Fan and J.P. Ostriker, “Combining Wilkinson Microwave Anisotropy Probe and Sloan Digital Sky Survey Quasar Data on Reionization Constrains Cosmological Parameters and Star Formation Efficiency”. *Astrophys. J.* **599**(2), 759-772 (2003).
- [7] P. Gondolo, Lectures Delivered at the NATO Advanced Study Institute Frontiers of the Universe, 2003 September 820, Cargese, France. (2004). <http://arxiv.org/abs/astro-ph/0403064>
- [8] G. Hinshaw, J.L. Weiland, R.S. Hill, N. Odegard, D. Larson, C.L. Bennett, J. Dunkley, B. Gold, M.R. Greason, N. Jarosik, E. Komatsu, M.R. Nolta, L. Page, D.N. Spergel, E. Wollack, M. Halpern, A. Kogut, M. Limon, S.S. Meyer, G.S. Tucker and E.L. Wright, *Astrophys. J. Suppl.* **180**, 225-245 (2009).
- [9] J.H. Oort, “The force exerted by the stellar system in the direction perpendicular to the galactic plane and some related problems”. *Bulletin of the Astronomical Institutes of the Netherlands* **6**(238), 249-287 (1932).
- [10] G. Bertone, D. Hooper and J. Silk, “Particle dark matter: evidence, candidates and constraints”. *Physics Reports* **405**(5-6), 279-390 (2005).
- [11] J. Diemand, M. Kuhlen, P. Madau, M. Zemp, B. Moore, D. Potter and J. Stadel, “Clumps and streams in the local dark matter distribution”. *Nature* **454**(7205), 735-738 (2008).
- [12] W.H. Press and D.N. Spergel, “Capture by the sun of a galactic population of weakly interacting, massive particles”. *Astrophys. J.* **296**, 679-684 (1985).

- [13] T. Damour and L.M. Krauss, “A New Solar System Dark Matter Population of Weakly Interacting Massive Particles”. *Phys. Rev. Lett.* **81**(26), 5726-5729 (1998).
- [14] A. Gould and S.M.K. Alam, “Can Heavy WIMPS Be Captured by the Earth?” *Astrophys. J.* **549**, 72-75 (2001).
- [15] J. Lundberg and J. Edsjö, “Weakly interacting massive particle diffusion in the solar system including solar depletion and its effect on Earth capture rates” *Physical Review D* **69**(12), 123505 (2004).
- [16] A.H.G. Peter and S. Tremaine “Dynamics of WIMPs in the Solar System and Implications for Direct and Indirect Detection” *Proceedings of "Identification of dark matter 2008". August 18-22, 2008, Stockholm, Sweden* pp. 61-65 (2008).
- [17] A.H.G. Peter, “Dark matter in the Solar System. I. The distribution function of WIMPs at the Earth from solar capture” *Physical Review D* **79**(10), 103531 (2009).
- [18] A.H.G. Peter, “Dark matter in the Solar System. II. WIMP annihilation rates in the Sun” *Physical Review D* **79**(10), 103532 (2009).
- [19] A.H.G. Peter, “Dark matter in the Solar System. III. The distribution function of WIMPs at the Earth from gravitational capture” *Physical Review D* **79**(10), 103533 (2009).
- [20] J.-M. Frère, F.-S. Ling and G. Vertongen, “Bound on the dark matter density in the Solar System from planetary motions”. *Physical Review D* **77**(8), 083005 (2008).
- [21] I.B. Khriplovich and D.L. Shepelyansky, “Capture of dark matter by the Solar System”. *International Journal of Modern Physics D* **18**(12), 1903-1912 (2009).
- [22] A. Gould, “Gravitational diffusion of solar system WIMPs”. *Astrophysics Journal* **368**, 610-615 (1991).
- [23] L. Bracci and G. Fiorentini, “The missing-mass problem and the precession of perihelia”. *Il Nuovo Cimento C* **12**(1), 121-127 (1989).
- [24] J.D. Anderson, E.L. Lau, A.H. Taylor, D.A. Dicus, D.C. Teplitz, V.L. Teplitz, “Bounds on dark matter in solar orbit”. *Astrophys. J.* **342**, 539-544 (1989).

- [25] V.B. Braginsky, A.V. Gurevich and K.P. Zybin, “The influence of dark matter on the motion of planets and satellites in the solar system”. *Physics Letters A* **171**(5-6), 275-277 (1992).
- [26] S. Klioner and M. Soffel, “Note on the influence of dark matter on the motion of the solar system”. *Physics Letters A* **184**(1), 41-44 (1993).
- [27] K. Nordtvedt, “Cosmic acceleration of Earth and the Moon by dark matter”. *Astrophys. J.* **437**, 529-531 (1994).
- [28] J.D. Anderson, E.L. Lau, T.P. Krisher, D.A. Dicus, D.C. Rosenbaum and V.L. Teplitz, “Improved Bounds on Nonluminous Matter in Solar Orbit”. *Astrophys. J.* **448**, 885-892 (1995).
- [29] K. Nordtvedt, J. Müller and M. Soffel, “Cosmic acceleration of the Earth and Moon by dark matter”. *Astron. Astrophys.* **293**, L73-L74 (1995).
- [30] Ø. Gron and H.H. Soleng, “Experimental limits on the density of dark matter in the solar system ”. *Astrophys. J.* **456**, 445-448 (1996).
- [31] I.B. Khriplovich and E.V. Pitjeva, “Upper Limits on Density of Dark Matter in Solar System”. *International Journal of Modern Physics D* **15**(4), 615-618 (2006).
- [32] M. Sereno and Ph. Jetzer, “Dark matter versus modifications of the gravitational inverse-square law: results from planetary motion in the solar system”. *Monthly Notices of the Royal Astronomical Society* **371**(2), 626-632 (2006).
- [33] L. Iorio, “Solar system planetary orbital motions and dark matter”. *Journal of Cosmology and Astroparticle Physics* **5**, 2 (2006).
- [34] I.B. Khriplovich, “Density of Dark Matter in the Solar System and Perihelion Precession of Planets”. *International Journal of Modern Physics D* **16**(9), 1475-1478 (2007).
- [35] S. Adler, “FAST TRACK COMMUNICATION: Placing direct limits on the mass of earth-bound dark matter”. *Journal of Physics A: Mathematical and Theoretical* **41**(41), 412002 (2008).
- [36] S. Adler, “Can the flyby anomaly be attributed to earth-bound dark matter?”. *Physical Review D* **79**(2), 023505 (2009).

- [37] H. Arakida, “Influence of Dark Matter on Light Propagation in Solar System”. *Advances in Space Research* doi:10.1016/j.asr.2009.11.012 (2010).
- [38] S.I. Blinnikov and M.Yu. Khlopov, “Possible astronomical effects of mirror particles”. *Soviet Astronomy* **27**(4), 371-375 (1983).
- [39] W.-T. Ni, “Astrod and Astrod I Overview and Progress”. *International Journal of Modern Physics D* **17**(7), 921-940 (2008).
- [40] M. Cerdonio, R. De Pietri, Ph. Jetzer and M. Sereno, “Local dark matter searches with LISA”. *Classical and Quantum Gravity* **26**(9), 094022 (2009).
- [41] N.S. Kardashev, A.V. Tutukov and A.V. Fedorova, “Limits on the Mass of Dark Matter in the Sun from a Model for the Modern Sun and Its Previous Evolution”. *Astronomy Reports* **49**(2), 134-143 (2005).
- [42] X. Xu and E.R. Siegel, “Dark Matter in the Solar System”. <http://arxiv.org/abs/0806.3767>
- [43] S. Mitra, “Uranus’s anomalously low excess heat constrains strongly interacting dark matter”. *Physical Review D* **70**(10), 103517 (2004).
- [44] S.L. Adler, “Planet-bound dark matter and the internal heat of Uranus, Neptune, and hot-Jupiter exoplanets”. *Physics Letters B* **671**(2), 203-206 (2009).
- [45] M.Yu. Khlopov, *Cosmoparticle physics*, (World Scientific, Singapore, 1999).
- [46] M.Yu. Khlopov, G.M. Beslin, N.G. Bochkarev, L.A. Pustil’nik and S.A. Pustil’nik, “Observational physics of the mirror world”. *Soviet Astronomy* **35**(1), 21-29 (1991).
- [47] L. Iorio, “Classical and relativistic orbital motions around a mass-varying body”. *Scholarly Research Exchange Physics* **2010**, ID 261249, (2010).
- [48] K. V. Kholshevnikov and M. Fracassini, “Le Problème des deux corps avec G variable selon l’hypothèse de Dirac”. *Conferenze dell’ Osservatorio Astronomico di Milano-Merate*, Serie I, No. 9, pp. 5-50, (1968).

- [49] K.P. Schröder and R. C. Smith, “Distant future of the Sun and Earth revisited”, *Monthly Notices of the Royal Astronomy Society* **386**(1), 155-163 (2008).
- [50] M. Fairbairn, P. Scott and J. Edsjö, “The zero age main sequence of WIMP burners”. *Physical Review D* **77**(4), 047301 (2008).
- [51] G.A. Krasinsky and V.A. Brumberg, “Secular increase of astronomical unit from analysis of the major planet motions, and its interpretation”. *Celestial Mechanics and Dynamical Astronomy* **90**(3-4), 267-288 (2004).
- [52] E.M. Standish, “The Astronomical Unit now”, in *Transits of Venus: New Views of the Solar System and Galaxy. Proceedings of the IAU Colloquium No. 196*, ed. by D.W. Kurtz. (Cambridge University Press, Cambridge, 2005), p. 177.
- [53] E.V. Pitjeva, cited by E.M. Standish, “The Astronomical Unit now”, in *Transits of Venus: New Views of the Solar System and Galaxy. Proceedings of the IAU Colloquium No. 196*, ed. by D.W. Kurtz. (Cambridge University Press, Cambridge, 2005), p. 176.
- [54] J.D. Anderson, “Astrometric Solar-System Anomalies”. *Bulletin of the American Astronomical Society* **41**, p.882 (2009). <http://arxiv.org/abs/0907.2469>
- [55] D.P. Rubincam, “On the secular decrease in the semimajor axis of Lagesos’s orbit”. *Celestial Mechanics and Dynamical Astronomy* **26**, 361-382 (1982).
- [56] V. Lainey, J.-E. Arlot, Ö. Karatekin and T. van Hoolst, “Strong tidal dissipation in Io and Jupiter from astrometric observations”. *Nature* **459**(7249), 957-959 (2009).