

## Lichen Records from Afyonkarahisar and Isparta Provinces

Gülşah ÇOBANOĞLU, Mustafa YAVUZ

Marmara University, Faculty of Arts and Science, Department of Biology, Göztepe Campus, 34722, İstanbul – TURKEY

Received: 20.03.2006

Accepted: 18.09.2006

**Abstract:** A list of 89 lichen taxa is reported from the provinces of Afyonkarahisar and Isparta. Fifteen taxa from Afyonkarahisar and 38 from Isparta are new records for these provinces.

**Key Words:** Lichenized fungi, biodiversity, Turkey

### Afyonkarahisar ve Isparta İllerinden Liken Kayıtları

**Özet:** Afyonkarahisar ve Isparta illerinden toplam 89 lichen taksonu listesi rapor edilmiştir. Afyonkarahisar ili için 15 takson, Isparta ili için 38 takson yeni kayittır.

**Anahtar Sözcükler:** Likenleşmiş mantarlar, biyoçeşitlilik, Türkiye

### Introduction

The number of floristic studies on lichens in Turkey has increased rapidly in recent decades. However, there are still gaps in the knowledge of many provinces. There have been 6 papers referring to lichens in Isparta province (Szatala, 1960; Mayrhofer, 1984; Mayrhofer et al., 1990; Öztürk et al., 1998; Öztürk & Kaynak, 1999; Öztürk et al., 2005). In these papers, 99 lichen species are given from several regions of Isparta province. There is only one lichen record from Afyonkarahisar (Giralt et al., 1992). The present study aims to contribute to the first floristic records in these 2 provinces and to the lichen flora of Turkey.

The study areas in both provinces (Isparta and Afyonkarahisar) are situated in the north-western Mediterranean region of Turkey. According to data from the Isparta meteorological station, the climate is generally Mediterranean with low values of humidity in the north. The average annual mean temperature is 12.1 °C. The annual precipitation is about 600.4 mm. The characteristic Mediterranean maquis elements are prevalent in the study area. Although Dinar is a city in

Afyonkarahisar province, it is closer to Isparta and the study area in Dinar represents similar aspects (Figure 1).

### Materials and Methods

The lichen material was collected from 5 sites in Isparta province and 1 in Afyonkarahisar province, in 2002, 2004 and 2005. The geographic coordinates and elevations were recorded as well as the substrate types. The taxa were identified with the aid of flora books and identification keys (Clauzade & Roux, 1985; Purvis et al., 1994; Wirth, 1995; Schultz & Büdel, 2002; Orange, 2004). The lichen specimens are preserved in the Herbarium of the Faculty of Science and Arts, Marmara University, İstanbul (MUFE).

#### Collecting Sites

##### ISPARTA:

1. Eğirdir (Akrotiri, Eğridir); Gökçehöyük (Banus) Village, Erenler Hill, 1260-1280 m, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004.




Figure 1. The study area and collecting sites indicated with numbers.

2. Eğirdir (Akrotiri, Eğridir); Gökçehöyük (Banus) Village, Akçapınar Hill, 1310-1320 m, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004.
  3. Eğirdir (Akrotiri, Eğridir); Gökçehöyük (Banus) Village, Mezarönü District, 1226 m, 37° 52' 51.4" N - 31° 00' 02.0" E, 31.VII.2005.
  4. Eğirdir (Akrotiri, Eğridir); Gökçehöyük (Banus) Village, Sırçalı District, 1640 m, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005.
  5. Eğirdir (Akrotiri, Eğridir); Eyüpler Village, 970 m, 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002.
- AFYONKARAHİSAR:
6. Dinar (Apamea); Bülbülcalan Village, 960-1000 m, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002.

## Results

Eighty-nine lichen taxa from the provinces of Afyonkarahisar (36 taxa) and Isparta (53 taxa) are listed alphabetically. Fifteen taxa are new records for Afyonkarahisar, 38 taxa are new for Isparta and 4 taxa are new for both provinces. An asterisk (\*) indicates a new record for the province of Isparta and a number sign (#) indicates a new record for the province of Afyonkarahisar. The herbarium numbers pertain to Gülsah Çobanoğlu (G.Ç.) in MUFE.

The names of authors are abbreviated according to Brummitt & Powell (1992). The nomenclature follows the recent literature (Söchting et al., 2002; Blanco et al., 2004a, 2004b; Santesson et al., 2004). The taxonomical order follows Purvis et al. (1992).

Below are the abbreviations used for substrata data in the list of taxa:

BR: on brick                    CR: on calcareous rock  
 SO: on soil                    SR: on siliceous rock  
 WM: with moss

#### List of Taxa

##### Ascomycotina

###### Order: Graphidales

###### Family: Thelotremaeaceae

*Diploschistes ocellatus* (Vill.) Norman

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1553).

###### Order: Lecanorales

###### Family: Acarosporaceae

*Acarospora cervina* A.Massal.

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1671). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1535).

\**Acarospora fuscata* (Nyl.) Arnold

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, BR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1673).

#*Sarcogyne privigna* var. *calcicola* H.Magn.

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1562).

###### Family: Hymeneliaceae

*Aspicilia calcarea* (L.) Mudd

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1633). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1675). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, BR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1674). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m,

CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1541).

\**Aspicilia cinerea* (L.) Körb.

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, SR., 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1681).

*Aspicilia contorta* (Hoffm.) Kremp. subsp. *contorta*

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, BR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1676). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1534).

*Lobothallia radiososa* (Hoffm.) Hafellner

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1636). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1557).

*Megaspora verrucosa* (Ach.) Hafellner & V.Wirth

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1612). Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1578).

###### Family: Rhizocarpaceae

\**Rhizocarpon geographicum* (L.) DC.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, SR, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1624). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, SR 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1682).

###### Family: Squamarinaceae

*Squamarina cartilaginea* (With.) P.James

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1545).

*Squamarina lentigera* (Weber) Poelt

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1546).

**Family: Biatoraceae**

\**Lecania inundata* (Hepp ex Körb.) M.Mayrhofer

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, BR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1677). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1678).

#*Lecania turicensis* (Hepp) Müll.Arg.

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1538).

#*Toninia candida* (Weber) Th.Fr.

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1537).

*Toninia sedifolia* (Scop.) Timdal

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1538).

#*Toninia squalida* (Ach.) A.Massal.

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1539).

**Family: Candelariaceae**

*Candelariella aurella* (Hoffm.) Zahlbr,

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, CR, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1581). Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1638). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1663). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, BR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1664). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1550).

#\**Candelariella vitellina* (Hoffm.) Müll.Arg.

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, SR., 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1666). Eğirdir; Gökçehöyük Village,

Sırçalı District, 1640 m, BR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1665). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1552).

\**Candelariella xanthostigma* (Pers. ex Ach.) Lettau

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Cupressus sempervirens*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1586). Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1614). Eğirdir; Gökçehöyük Village, Mezarönü District, 1226 m, on *Prunus domesticus*, 37° 52' 51.4" N - 31° 00' 02.0" E, 31.VII.2005, (G.Ç. 1645). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, on *Cupressus sempervirens*, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1650). Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1567).

**Family: Lecanoraceae**

\**Lecanora argentata* (Ach.) Malme

Isparta: Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1573).

\**Lecanora carpinea* (L.) Vain.

Isparta: Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1576).

*Lecanora chlarotera* Nyl.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1598). Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1577).

*Lecanora crenulata* Hook.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, CR, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1582). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1547).

*Lecanora dispersa* (Pers.) Sommerf.

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, BR, 37° 51' 46.0" N - 31° 01' 18.6" E,

01.VIII.2005, (G.Ç. 1680). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR,  $37^{\circ} 51' 46.0''$  N -  $31^{\circ} 01' 18.6''$  E, 01.VIII.2005, (G.Ç. 1679). Afyonkarahisar: Dinar; Bülbülgan Village, 960-1000 m, CR,  $38^{\circ} 03' 04.1''$  N -  $30^{\circ} 11' 47.0''$  E, 22.IV.2002, (G.Ç. 1555).

\**Lecanora piniperda* Körb.

Isparta: Eğirdir; Gökçehöyük Village, Mezarönü District, 1226 m, on *Prunus domesticus*,  $37^{\circ} 52' 51.4''$  N -  $31^{\circ} 00' 02.0''$  E, 31.VII.2005, (G.Ç. 1643). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, on *Cupressus sempervirens*,  $37^{\circ} 51' 46.0''$  N -  $31^{\circ} 01' 18.6''$  E, 01.VIII.2005, (G.Ç. 1652).

\**Lecanora sambuci* (Pers.) Nyl.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Cupressus sempervirens*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1587). Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1626).

\**Lecanora varia* (Hoffm.) Ach.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1617). Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Sorbus takhtajanii*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1619).

*Lecidella elaeochroma* (Ach.) M.Choisy

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1615). Eğirdir; Eyüp Village, 970 m, on *Juniperus* sp.,  $37^{\circ} 46' 07.8''$  N -  $30^{\circ} 52' 04.7''$  E, 06.XII.2002, (G.Ç. 1566).

\**Lecidella stigmatea* (Ach.) Hertel & Leuckert

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR,  $37^{\circ} 53' 06''$  N -  $31^{\circ} 00' 54.7''$  E, 25.VII.2004, (G.Ç. 1639).

*Protoparmeliopsis muralis* (Schreb.) M.Choisy

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, CR,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1583). Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1623). Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR,  $37^{\circ} 53' 06''$  N -  $31^{\circ} 00' 54.7''$  E, 25.VII.2004,

(G.Ç. 1635). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, SR,  $37^{\circ} 51' 46.0''$  N -  $31^{\circ} 01' 18.6''$  E, 01.VIII.2005, (G.Ç. 1662). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR,  $37^{\circ} 51' 46.0''$  N -  $31^{\circ} 01' 18.6''$  E, 01.VIII.2005, (G.Ç. 1660). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, BR,  $37^{\circ} 51' 46.0''$  N -  $31^{\circ} 01' 18.6''$  E, 01.VIII.2005, (G.Ç. 1661). Afyonkarahisar: Dinar; Bülbülgan Village, 960-1000 m, CR,  $38^{\circ} 03' 04.1''$  N -  $30^{\circ} 11' 47.0''$  E, 22.IV.2002, (G.Ç. 1560).

#### Family: Parmeliaceae

\**Melanohalea exasperata* (De Not.) O.Blanco et al.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1592). Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Sorbus takhtajanii*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1608).

\**Melanohalea septentrionalis* (Lynge) O.Blanco et al.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1591).

\**Parmelina pastillifera* (Harm.) Hale

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, CR,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1584). Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Sorbus takhtajanii*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1610).

\**Parmelina tiliacea* (Hoffm.) Hale

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1593). Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Sorbus takhtajanii*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ} 00' 24.4''$  E, 22.VII.2004, (G.Ç. 1609). Eğirdir; Eyüp Village, 970 m, on *Juniperus* sp.,  $37^{\circ} 46' 07.8''$  N -  $30^{\circ} 52' 04.7''$  E, 06.XII.2002, (G.Ç. 1570).

*Pseudevernia furfuracea* (L.) Zopf var. *ceratea* (Ach.) D.Hawkes.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*,  $37^{\circ} 52' 21.8''$  N -  $31^{\circ}$

00° 24.4" E, 22.VII.2004, (G.Ç. 1602). Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1564).

\**Xanthoparmelia pulla* (Ach.) O.Blanco et al.

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, SR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1589).

#### Family: Physciaceae

*Anaptychia ciliaris* (L.) Körb. ex A.Massal.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1603).

\**Buellia disciformis* (Fr.) Mudd

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1616).

\**Buellia pulverulenta* (Anzi) Jatta

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1618).

*Diplotomma alboatrum* (Hoffm.) Flot.

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1634). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1554).

#*Phaeophyscia orbicularis* (Neck.) Moberg

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1597). Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Sorbus takhtajanii*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1620). Eğirdir; Gökçehöyük Village, Mezarönü District, 1226 m, on *Prunus domesticus*, 37° 52' 51.4" N - 31° 00' 02.0" E, 31.VII.2005, (G.Ç. 1647). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1544).

*Physcia adscendens* (Th.Fr.) H.Olivier

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, on *Sorbus takhtajanii*, 1260-1280 m, 37° 52' 21.8" N - 31°

00' 24.4" E, 22.VII.2004, (G.Ç. 1621). Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1565). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1558).

*Physcia aipolia* (Ehrh. ex Humb.) Fürnr.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1595).

*Physcia dubia* (Hoffm.) Lettau

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR, on *Protoparmeliopsis muralis*, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1659 & G.Ç. 1653). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1540).

*Physcia semipinnata* (J.F.Gmel.) Moberg

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1601). Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Sorbus takhtajanii*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1622). Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1569).

*Physcia stellaris* (L.) Nyl.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Sorbus takhtajanii*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1611). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, on *Cupressus sempervirens*, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1649).

*Physcia tenella* (Scop.) DC.

Isparta: Eğirdir; Gökçehöyük Village, Mezarönü District, 1226 m, on *Prunus domesticus*, 37° 52' 51.4" N - 31° 00' 02.0" E, 31.VII.2005, (G.Ç. 1646).

\**Physcia tribacioides* Nyl.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Cupressus sempervirens*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1588).

\**Physconia distorta* (With.) J.R.Laundon

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31°

00° 24.4" E, 22.VII.2004, (G.Ç. 1596). Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Sorbus takhtajanii*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1607). Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1568).

\**Physconia grisea* (Lam.) Poelt

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, CR, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1585).

*Physconia perisidiosa* (Erichsen) Moberg

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, SO, with *Cladonia* sp. and WM, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1605).

#*Rinodina bischoffii* (Hepp) A.Massal.

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1539).

#*Rinodina calcarea* (Arnold) Arnold

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1561).

*Rinodina exigua* (Ach.) Gray

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1613). Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1574).

\**Rinodina gennarii* Bagl.

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1642).

Family: Collemataceae

*Collema crispum* (L.) Weber ex F.H. Wigg. var. *crispum*

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, SO, with *Cladonia* sp. and WM, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1606). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1549).

*Collema cristatum* var. *marginale* (Huds.) Degel.

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1627).

\**Collema subflaccidum* Degel.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1604). Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1575).

*Leptogium lichenoides* (L.) Zahlbr.

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, WM, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1628). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, WM, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1556).

Family: Placynthiaceae

*Placynthium nigrum* (Huds.) Gray

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1629). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1537).

Family: Pertusariaceae

\**Ochrolechia parella* (L.) A.Massal.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 16594).

\**Ochrolechia tartarea* (L.) A.Massal.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1599). Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1571).

\**Pertusaria albescens* (Huds.) M.Choisy & Werner

Isparta: Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1579).

\**Pertusaria opthalmiza* (Nyl.) Nyl.

Isparta: Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1580).

Family: Teloschistaceae

\**Caloplaca alociza* (A.Massal.) Mig.

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1637).

*Caloplaca aurantia* (Pers.) Hellb.

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1542).

*Caloplaca cerina* (Ehrh. ex Hedw.) Th.Fr. var. *cerina*

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Quercus cerris*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1600). Eğirdir; Eyüpler Village, 970 m, on *Juniperus* sp., 37° 46' 07.8" N - 30° 52' 04.7" E, 06.XII.2002, (G.Ç. 1572).

*Caloplaca cerina* var. *chloroleuca* (Sm.) Th.Fr.

Isparta: Eğirdir; Gökçehöyük Village, Mezarönü District, 1226 m, on *Prunus domesticus*, 37° 52' 51.4" N - 31° 00' 02.0" E, 31.VII.2005, (G.Ç. 1648). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, on *Cupressus sempervirens*, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1651).

\**Caloplaca chalybaea* (Fr.) Müll.Arg.

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1632).

\**Caloplaca crenulatella* (Nyl.) H.Olivier

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, BR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1672).

#*Caloplaca decipiens* (Arnold) Blomb. & Forssell

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1535).

#*Caloplaca erythrocarpa* (Pers.) Zwackh

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1551).

\**Caloplaca flavescens* (Huds.) J.R.Laundon

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1669).

*Caloplaca holocarpa* (Hoffm.) A.E.Wade

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1536).

#*Caloplaca saxicola* (Hoffm.) Nordin

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1543).

*Caloplaca variabilis* (Pers.) Müll.Arg.

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1657). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, BR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1658).

#*Fulglesia bracteata* var. *bracteata* (Hoffm.) Räsänen

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1544).

Xanthomendoza fulva (Hoffm.) Söchting, Kärnefelt & S.Y.Kondr.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Cupressus sempervirens*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1625).

Xanthoria elegans (Link) Th.Fr.

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1668).

Xanthoria parietina (L.) Th.Fr.

Isparta: Eğirdir; Gökçehöyük Village, Erenler Hill, 1260-1280 m, on *Cupressus sempervirens*, 37° 52' 21.8" N - 31° 00' 24.4" E, 22.VII.2004, (G.Ç. 1590). Eğirdir; Gökçehöyük Village, Mezarönü District, 1226 m, on *Prunus domesticus*, 37° 52' 51.4" N - 31° 00' 02.0" E, 31.VII.2005, (G.Ç. 1644).

Incertae sedis:

*Leproloma membranaceum* (Dicks.) Vain.

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1640).

**Order: Verrucariales****Family: Verrucariaceae**

\* *Placidium rufescens* (Ach.) A.Massal.

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, WM, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1641).

***Placocarpus schaeferi* (Fr.) Breuss**

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1667). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1559).

**#*Verrucaria fuscella* (Turner) Winch**

Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1540).

**\**Verrucaria glaucina* Ach.**

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1655). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, BR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1656).

**\**Verrucaria hochstetteri* Fr.**

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, 37° 53' 06" N - 31° 00' 54.7" E, 25.VII.2004, (G.Ç. 1631).

**#\**Verrucaria muralis* Ach.**

Isparta: Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1670). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1548).

**#\**Verrucaria nigrescens* Pers.**

Isparta: Eğirdir; Gökçehöyük Village, Akçapınar Hill, 1310-1320 m, CR, 37° 53' 06" N - 31° 00' 54.7" E,

25.VII.2004, (G.Ç. 1630). Eğirdir; Gökçehöyük Village, Sırçalı District, 1640 m, CR, 37° 51' 46.0" N - 31° 01' 18.6" E, 01.VIII.2005, (G.Ç. 1654). Afyonkarahisar: Dinar; Büyüçalan Village, 960-1000 m, CR, 38° 03' 04.1" N - 30° 11' 47.0" E, 22.IV.2002, (G.Ç. 1543).

**Discussion**

The list includes 89 taxa (88 species and 1 subspecies) belonging to 38 genera, of which 19 taxa are common for both provinces. The most frequent 4 genera within the study area are *Caloplaca* (represented by 11 species), *Lecanora* (represented by 8 species), *Physcia* (represented by 7 species) and *Verrucaria* (represented by 5 species).

The lichen composition is rather rich at collecting sites 1, 4 and 6. Collecting site 1, a small hill on the western slopes of Mount Kekikdağ, has the highest percentage of taxa (29.2%) on a diversity of substrata. Site 4, a high plateau with some *Cupressus* sp. trees, historical ruins of an ancient basilica and bricks, also has a large number of taxa (22.1%). This is the only collecting site where siliceous rocks were found. At collecting site 6 (Afyonkarahisar), there were no corticolous lichen samples, due to a lack of plant vegetation; only calcicolous taxa were found (23.4%).

The saxicolous taxa are higher in number in all the study sites with 57.1%, 46.1% of which are on calcareous rocks, 7.1% on bricks and 3.9% on siliceous rocks. The terricolous taxa are 1.3% of the total. As no significant forest vegetation is present in the study area, the epiphytic taxa is rather poor (41.6%) and limited to scattered trees such as *Juniperus*, *Prunus*, *Quercus*, *Sorbus* and *Cupressus* spp. The number of corticolous taxa is higher on *Quercus* sp. (14.9%) and *Juniperus* sp. (11%), decreasing on *Cupressus* sp. (5.8%), on *Sorbus* sp. (5.8%) and on *Prunus* sp. (3.9%). Substrate types of all the taxa correspond with those in the literature.

**References**

Blanco O, Crespo A, Divakar PK, Esslinger TL, Hawksworth DL & Lumbsch TH (2004a). *Melanelia* and *Melanoalea*, two new genera segregated from *Melanelia* (Parmeliaceae) based on molecular and morphological data. *Mycol Res* 108: 873-884.

Blanco O, Crespo A, Elix JA, Hawksworth DL & Lumbsch HT (2004b). A molecular phylogeny and a new classification of parmelioid lichens containing *Xanthoparmelia*-type lichenan (Ascomycota: Lecanorales). *Taxon* 53: 959-975.

- Brummitt RK & Powell CE (1992). *Authors of Plant Names*. Royal Botanical Gardens, Kew: 1-732.
- Clauzade G & Roux C (1985). *Likenoj De Okcidenta Europo Illustrata Determinlibro*. Bulletin de la Société Botanique du Centre – Quest Nouvelle Série – Numéro Spécial: 7, Royan, France.
- Giralt M, Nimis PL & Poelt J (1992). Studien über den Formenkreis von *Caloplaca flavorubescens* in Europa. *Cryptogamie Bryol Lichénol* 13: 261-273.
- Mayrhofer H (1984). Die saxicolen Arten der Flechtengattungen *Rinodina* und *Rinodinella* in der Alten Welt. *J Hattori Bot Lab* 55: 327-493.
- Mayrhofer H, Scheidegger C & Sheard J (1990) *Rinodina lecanorina* and *R. saridata*, two closely related species on calciferous rocks *Biblioth. Lichenol* 38: 335.
- Orange A (2004). The *Verrucaria fuscella* group in Great Britain and Ireland. *Lichenologist* 36: 173-182.
- Öztürk Ş, Kaynak G. & Güvenç Ş (1998). New floristic records for the various grid squares from the lichen flora of Turkey. *OT Sistematisatik Botanik Dergisi* 5: 93-98.
- Öztürk Ş & Kaynak G. (1999). New records for the lichen flora of Turkey. *Turk J Bot* 23: 357-358.
- Öztürk Ş, Güvenç Ş & Aydin S (2005). Floristic lichen records from Isparta and Burdur provinces. *Turk J Bot* 29: 243-250.
- Purvis OW, Coppins BJ, Hawksworth DL, James PW & Moore DM (1992). *The Lichen Flora of Great Britain and Ireland*. London: Natural History Museum Publications in association with the British Lichen Society.
- Santesson R, Moberg R, Nordin A, Tønsberg T & Vitikainen O (2004). *Lichen-forming and lichenicolous fungi of Fennoscandia*. Göteborg: Museum of Evolution, Uppsala University, Majornas Copyprint AB.
- Schultz M & Büdel B (2002). Key to the genera of the Lichinaceae. *Lichenologist* 34: 39-62.
- Söchting U, Kärnefelt I & Kondratyuk SY (2002). Revision of *Xanthomendoza* (Teloschistaceae, Lecanorales) based on morphology, anatomy, secondary metabolites and molecular data. In: Mitteilungen aus dem Institut für Allgemeine Botanik in Hamburg 30-32: 225-240.
- Szatala Ö (1960). Lichenes Turciacae Asiaticae ab Victor Pietschmann collecti. *Sydomia* 14: 312-325.
- Wirth V (1995). *Die Flechten Baden-Württembergs*. Teil 1-2. Stuttgart: Ulmer.